# This Page Is Inserted by IFW Operations and is not a part of the Official Record

# **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

# IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.





## **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) WO 95/09634 (51) International Patent Classification 6: (11) International Publication Number: A1 A61K 31/69, C07F 5/02, 5/04 13 April 1995 (13.04.95) (43) International Publication Date: (74) Agents: KERR, Don, M. et al.; The du Pont Merck Pharmaceu-PCT/US94/11280 (21) International Application Number: tical Company, Legal/Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US). 6 October 1994 (06.10.94) (22) International Filing Date: (81) Designated States: AU, CA, CZ, FI, HU, JP, KR, NO, NZ, PL, SK, European patent (AT, BE, CH, DE, DK, ES, FR, (30) Priority Data: US 7 October 1993 (07.10.93) 08/133,251 GB, GR, IE, IT, LU, MC, NL, PT, SE). 20 October 1993 (20.10.93) US 08/139,445 (71) Applicant: THE DU PONT MERCK PHARMACEUTICAL Published With international search report. COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: GALEMMO, Robert, Anthony, Jr.; 3039 Stump

Hall Road, Collegeville, PA 19426 (US). ABELMAN, Matthew, Mark; Apartment 3312, 873 Stevens Avenue, Solana Beach, CA 92075 (US). AMPARO, Eugene, Cruz; 107 Garfield Avenue, West Chester, PA 19380-4509 (US). FEVIG, John, Matthew; 987 Church Road, Lincoln University, PA 19352-9349 (US). KNABB, Robert, Madara; 14 Sullivan Chase Drive, Avondale, PA 19311-9349 (US). MILLER, William, Henry; 1909 Coventry Lane, Glen Mills, PA 19342 (US). PACOFSKY, Gregory, James; 510 Polk Street, Raleigh, NC 27604 (US). WEBER, Patricia, Carol; 107 Marshall Bridge Road, Kennett Square, PA 19348-2705 (US).

## (54) Title: ELECTROPHILIC PEPTIDE ANALOGS AS INHIBITORS OF TRYPSIN-LIKE ENZYMES

### (57) Abstract

This invention relates to electrophilic dipeptide analogs conjugated to an N,N disubstituted α-amino acid as inhibitors of trypsin-like serine protease enzymes.

NSDOCID: <WO\_\_9509634A1\_I\_>

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE.	Niger
BE	Belgium	GR	Greece	NL.	Netherlands
BF	Burkina Faso	BU	Hungary	NO	Norway
BG	Bulgaria	Œ	Ireland	NZ.	New Zealand
BJ	Benin	17	Italy	PL	Poland
BR	Brazil	JP		PT	
BY	Belarus	KE	Kenya	RO	Portugal ·
CA	Canada	KG	Kyrgystan	RU	Romania
CF	Central African Republic	KP	Democratic People's Republic		Russian Federation
CG	Congo		of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	KZ	Kazakhstan	SI	Slovenia
CM	Cameroon	Li	Liechtenstein	SK	Slovakia
CN	China	LK		SN	Senegal
CS	Czechoslovakia	LU	Sri Lanka	TD	Chad
cz	Czech Republic	LV	Luxembourg	TG	Togo
DE	Germany		Latvia	TJ	Tajikistan
DK	Demnark	MC	Monaco	TT	Trinidad and Tobago
ES	Spain	MD	Republic of Moldova	UA.	Ukraine
FI	Finland	MG	Madagascar	. US	United States of America
FR		MIL.	Mali	UZ	Uzbekistan
GA	Prance Gabon	MIN	Mongolia	VN	Viet Nam

### Title

Electrophilic Peptide Analogs As Inhibitors of Trypsin-Like Enzymes

# Cross-reference to Earlier Filed Applications

This application is a continuation-in-part of U.S. Patent Application Serial Number 08/139,445 filed October 20, 1993.

10

15

20

25

5

# Field of the Invention

This invention relates generally to electrophilic peptide analogs as inhibitors of trypsin-like serine proteases. These compounds are dipeptides in which an electrophilic derivative of an  $\alpha$ -amino acid is conjugated with an N,N-disubstituted  $\alpha$ -amino acid. The N,N-disubstituted  $\alpha$ -amino acid conjugates of the electrophilic amino acid analog are derivatives of an amino acid where the  $\alpha$ -amino group is alkylated and acylated or diacylated to give alicyclic or cyclic substituents. The electrophilic functional groups used to derivatize these peptide analogs are: boronic acids and their esters,  $\alpha$ -mono- and  $\alpha$ -perhaloketones, aldehydes, vicinal di- and tricarbonyl compounds,  $\alpha$ -mono- and  $\alpha$ -dihalo- $\beta$ -ketoesters.

# Background of the Invention

containing the ((D-phenylalanyl)prolyl)- arginyl- sequence are well known as effective inhibitors of the trypsin-like serine protease thrombin. H-(D)Phe-Pro-ArgCH<sub>2</sub>Cl was first reported by Kettner and Shaw (Thromb. Res. 14, 969 (1979)) to be a selective but irreversible inhibitor of human thrombin. A number of studies looking for alternatives to the electrophilic P<sub>1</sub> argininechloromethylketones that would

10

20

yield a reversible protease inhibitor have been reported. Bajuez et al. (Folia Haematol. 109, s. 16 (1982)) found the corresponding aldehyde, p-phenylalanyl-prolyl-arginal, to be a reversible thrombin inhibitor with a  $K_i$  = 75 nM for human thrombin. The nitrile analog, p-phenylalanyl-prolyl- $NHCH((CH_2)_3NHC(=NH)NH_2)-CN$ , was found to be substantially less potent with a  $K_i = 700$  nM (Kaiser et al., Pharmazie 46, 128 (1991)). A retroamide inhibitor, with the Dphenylalanyl-prolyl- sequence and 2-(4guanidinophenylalanyl)-N-acetyl-2,2-difluoroethylamine substituting for an electrophilic arginine derivative, is a good inhibitor with a  $K_i$  of 70 nM for thrombin (Altenburger and Schirlin, Tetrahedron Lett. 32, 7255 (1991)). Cheng et al. claim that the substitution of racemic diphenyl 1-15 amino-4-methoxybutylphosphonate for an electrophilic

arginine derivative gives very good inhibitors with a  $K_1$  = 4.8 nM (Tetrahedron Lett. 32, 7333 (1991)). Iwanowicz et al. (Bioorgan. Med. Chem. Lett. 2, 1607 (1992)) has studied the efficacy of (D-phenylalanine)prolyl- conjugated to -NHCH[(CH $_2$ ) $_4$ NH $_2$ ]CH(OH)CO $_2$ Me) and -NHCH[(CH $_2$ ) $_4$ NH $_2$ ]C(=O)CO $_2$ Me derivatives. The most effective inhibitor of human thrombin reported to date is the boropeptide acetyl-Dphenylalanyl-prolyl-boro arginine with a  $K_i$  = 0.041 nM (Kettner et al., J. Biol. Chem. 265, 18289 (1990)).

25 Walker et al. (Biochem. J. 230, 645 (1985)) published a comparative study of irreversible thrombin inhibitors based on the D-phenylalanyl-prolyl-argininyl sequence confirming the earlier report by Kettner and Shaw (1979). H-(D)Phe-Pro-ArgCH2Cl was found to be the most effective inhibitor ( $K_i = 25 \text{ nM}$ ) while replacing the D-phenylalanine with 4-amino-D-phenylalanine or  $\omega$ -benzoyl-D-lysine gave less active analogs. Compounds in development include -(prolyl)arginal derivatives with a variety of unusual  $P_3$ amino acids including D-N-methylphenylglycine, Boc-Dfluorophenylglycine as well as constrained cyclized

derivatives of D-phenylglycine and D-phenylalanine (Shuman et. al., J. Med. Chem. 36, 314 (1993)). Balasubramanian et al. (J. Med. Chem. 36, 300 (1993)) has reported an extensive study of replacements for the P<sub>3</sub> D-phenylalanine of D-phenylalanyl-prolyl-arginal and found the dihydrocinnamoyl group to be effective, although somewhat less potent.

Patent disclosures in this area have centered around suitably protected peptides composed of natural and 10 unnatural amino acids. In U.S. Patent No. 5,187,157 DuPont Merck has disclosed peptides comprised of C-terminal boronic acid derivatives of lysine, ornithine and arginine as reversible inhibitors of trypsin-like serine proteases, as well as a series of boropeptides active as elastase 15 inhibitors in U.S. Patent No. 4,499,082. In European Patent Application EP 471 651 A2 Sandoz disclosed borolysine and boroarginine peptide analogs containing at least one unnatural hydrophobic  $\alpha$ -amino acid substituted with groups such as the trimethylsilyl- or naphthyl-. U.S. Patent No. 5,106,948 was disclosed a series of boropeptides that are effective as cytotoxic agents. PCT Application WO 92/07869, Thrombosis Research Institute has disclosed tripeptide analogs containing a P2 proline and an unnatural disubstituted amino acid at P3. A variety of electrophilic and non-electrophilic  $\alpha$ -amino acid analogs 25 were claimed as suitable P1 substituents. Tripeptide antithrombotic agents limited to  $\alpha\text{-alkyl}$  and  $\alpha$  aryl or heteroaryl substituted glycines at P3 conjugated to -(prolyl)arginal were been disclosed by Lilly (European Patent Application EP 479 489 A2). Marion Merrell Dow 30 disclosed a series of activated electrophilic ketone analogs of peptidase substrates useful for inhibiting serine-, carboxylic acid- and metallo- proteolytic enzymes; compounds are peptides composed of suitably protected  $\alpha$ amino acids conjugated to an electrophilic ketone . 35

derivative of an  $\alpha$ -amino acid (European Patent Applications EP 417 721 A2, EP 364 344 A2, EP 363 284 A2, EP 195 212 A2). Astra has disclosed a series of  $\alpha$ - ((trifluoroethyl)oxymethyl)-arginine tripeptides (European Patent Application EP 0 530 167 A). Georgia Tech Research Corporation disclosed peptidyl ketoamides, -ketoacids and -ketoesters as inhibitors of serine and cysteine proteases (WO 92/12140). Boehringer Ingelheim disclosed a series of trifluoromethyl- and  $\alpha\alpha$ -difluoromethyl- $\beta$ ketoesterpeptide derivatives as elastase inhibitors (EP 0 369 391 A2).

The present invention concerns dipeptides which contain an electrophilic derivative of an  $\alpha$ -amino acid at  $P_1$  conjugated with an N,N-disubstituted  $\alpha$ -amino acid at  $P_2$ . 15 The electrophilic functional groups used to derivatize the P<sub>1</sub> amino acid analog are: boronic acids and their esters,  $\alpha$ -mono- and  $\alpha$ -perhaloketones, vicinal di- and tricarbonyl compounds, and  $\alpha\alpha$ -dihalo- $\beta$ ketoesters. The N,Ndisubstituted  $\alpha$ -amino acids are derivatives of an amino 20 acid other than proline where the  $\alpha$ -amino group is alkylated and acylated or diacylated to give alicyclic or cyclic substituents. As a result these compounds are found to have the advantage of an improved toxicological profile as well as the selectivity and inhibition activity for 25 thrombin required for a useful therapeutic agent.

### Summary of the Invention

30 [1] There is provided by this invention a compound of the formula (I):

10

or a pharmaceutically acceptable salt, hydrate or prodrug thereof, wherein:

$$R^{1}$$
 is

a)  $-(C_1-C_{12} \text{ alkyl})-X$ ,

b)  $-(C_1-C_{12} \text{ alkenyl})-X$ , or

10 c)

$$-(CH_2)_q$$
  $(CH_2)_pX$ 

X is

a) halogen,

15 b) -CN,

 $c) -NO_2,$ 

d) -CF<sub>3</sub>,

e) -NH<sub>2</sub>,

f) -NHOR<sup>2</sup>,

g) -NHC(=NH) $\mathbb{R}^2$ ,

h) -NHC(=NH)NHOH,

i) -NHC(=NH)NHNH2;

j) -NHC(=NH)NHCN,

)  $-NHC (=NH) NHR^2$ ,

j) -NHC(=NH)NHCOR<sup>2</sup>,

k) -C (=NH) NHR<sup>2</sup>,

1)  $-C (=NH) NHCOR^2$ ,

m)  $-C (=0) NHR^2$ ,

```
n) - CO_2R^2
            o) -OR^2.
            p) -OCF3,
            q) -S(0)_{r}R^{2},
            r) -SC(=NH)NHR^2, or
   5
            s) -SC(=NH)NHC(=0)R^2;
       R^2 is
            a) hydrogen, or
 10
            b) C<sub>1</sub>-C<sub>4</sub> alkyl;
       R<sup>3</sup> is:
             a) -C(=0)-aryl,
             b) -C(=0) - (CH_2)_p - CR^6R^7 - (CH_2)_q - aryl,
 15
             c) -C(=0)-(C_2-C_5 \text{ alkenyl})-aryl,
             d) -C(=0)-W-CR^8R^9-aryl, with the proviso that W cannot
                   be a bivalent oxygen atom,
             e) -C(=0)-CR^8R^9-W-(CH_2)_r-aryl, with the proviso that W
                   cannot be -NR^4- or -NC(=0)R^4-,
 20
             f) -C(=0)-heteroaryl,
             g) -C(=0)-(CH<sub>2</sub>)_p-CR<sup>6</sup>R<sup>7</sup>-(CH<sub>2</sub>)_q-heteroaryl,
            h) -C(=0)-(C_2-C_5 \text{ alkenyl})-\text{heteroaryl},
            i) -C(=0)-W-CR8R9-heteroaryl,
            j) -C(=0)-CR^8R^9-W-(CH_2)_r-heteroaryl, with the proviso
25
                   that W cannot be -NR^4- or -NC(=0)R^4-,
            k) -C(=0)-heterocycle,
            1) -C(=0)-(CH_2)_p-CR^6R^7-(CH_2)_q-heterocycle,
            m) -C(=0)-(C_2-C_5 \text{ alkenyl})-\text{heterocycle},
            n) -C(=0)-W-CR8R9-heterocycle,
            o) -C(=0)-CR^8R^9-W-(CH_2)_r-heterocycle, with the proviso
30
                  that W cannot be -NR4- or -NCOR4-,
            p) -C(=0)-(CH_2)_t-adamantyl,
            q) -C(=0)-(CH_2)_{t}-(C_5-C_7 \text{ cycloalkyl}),
            r) -C(=0)-(CH_2)_t-W-(C_5-C_7 \text{ cycloalkyl}),
35
            s)
```

H or 
$$R^{13}$$
 (CH<sub>2</sub>)<sub>t</sub>-aryl, wherein aryl is limited to phenyl,

H or  $R^{13}$   $(CH_2)_p\text{-W-}(CH_2)_q\text{-aryl}, \text{ wherein aryl is}$  limited to phenyl,

H or R<sup>13</sup>

10 v) H or R<sup>13</sup>

w)

5

with the proviso that  $R^{13}$  cannot be  $-N(C_1-C_4 \text{ alkyl})_2$ when A is  $-C(=0)R^{14}$ ,

10

5

aa)

ee)

10

\_\_\_

5

hh)

10

ii) .

5

10

O R<sup>18</sup> R<sup>18</sup> O NHR<sup>17</sup>

00)  $-C (=0) - (CR^8R^9) - NHS (0) rR^8;$ pp)

H or R<sup>13</sup>
(CH<sub>2</sub>),-heteroaryl

15 qq)

H or R<sup>13</sup>

(CH<sub>2</sub>)<sub>n</sub>-W-(CH<sub>2</sub>)<sub>a</sub>-heteroaryl

 ${\bf R}^4$  and  ${\bf R}^5$  are independently selected at each occurrence from the group consisitng of:

PCT/US94/11280

```
a) hydrogen,
              b) C_1-C_4 alkyl,
              c) -(C_1-C_4 \text{ alkyl}) -aryl, or
              d) C5-C7 cycloalkyl;
  5
       {\rm R}^6,~{\rm R}^7,~{\rm R}^8 and {\rm R}^9 are independently selected at each
       occurrence from the group consisting of:
              a) hydrogen,
             b) C_1-C_4 alkyl,
 10
             c) C_1-C_4 alkoxy,
             d) arvl,
             e) -(C_1-C_4 \text{ alkyl})-\text{aryl},
             f) -(C_1-C_4 \text{ alkyl})-heterocycle,
             g) -0-aryl,
 15
             h) -(CH_2)_{p}-CO_2R^4,
             i) {\ensuremath{R^6}} and {\ensuremath{R^7}} can be taken together to form a (C2-C7)
                    alkyl, or
             j) R^8 and R^9 can be taken together to form a (C_2-C_7)
                   alkyl;
20
      R<sup>10</sup> is:
             phenyl, wherein phenyl is optionally substituted with
             one to three substituents selected from the group
             consisting of halogen, C_1-C_4 alkyl, C_1-C_4 alkoxy, C_7-
25
            C<sub>15</sub> alkylaryl, C<sub>7</sub>-C<sub>15</sub> alkoxyaryl, methylenedioxy,
             -NO_2, -CF_3, -SH, -S(O)_r - (C_1 - C_4 \text{ alkyl}), CN, -OH, -NH_2,
            -NH(C_1-C_4 \text{ alkyl}), -N(C_1-C_4 \text{ alkyl})_2, -NHCOR^4, -(CH_2)_{p-1}
            CO_2R^4, -C(=NH)NHR^4), -NHC(=NR^4)R^4, -NHC(=NH)NHR^4;
30
     R<sup>11</sup> is:
```

- - a)  $C_1-C_4$  alkyl,
  - b) C<sub>3</sub>-C<sub>6</sub> cycloalkyl,
  - c)  $-OR^4$ ,
  - d)  $-NR^{15}R^{16}$ ,
- e)  $-NC(=0)R^{15}R^{16}$ , 35

f)  $-NR^{15}C(=0)OR^4$ ,

g) aryl,

h)  $-(C_1-C_4 \text{ alkyl})-\text{aryl}$ ,

i) heteroaryl,

j) -( $C_1$ - $C_4$  alkyl)-heteroaryl,

 $k) - (C_1-C_4 \text{ alkyl}) - CO_2R^4$ 

1) heterocycle,

m)  $-(C_1-C_4 \text{ alkyl})$  heterocycle,

n)

0)

5

10

p)

15 q)

r)

 ${\bf R}^3$  and  ${\bf R}^{11}$ , when taken together to form a ring bonded to the nitrogen:

5 a)

10

d)

5

g)

10

h)

 ${\bf R}^{11}$ ' is independently selected at each occurrence from the group consisting of:

- 5 a) hydrogen;
  - b)  $C_1-C_4$  alkyl
  - $c) OR^4$
  - d)  $-NR^{15}R^{16}$
  - e)  $-NC(=0)R^{15}R^{16}$
- 10 f)  $-NR^{15}C(=0)OR^4$ 
  - g) aryl,
  - h)  $-(C_1-C_4 \text{ alkyl})-aryl,$ 
    - i) heteroaryl,
    - $j) (C_1-C_4 \text{ alkyl}) \text{heteroaryl},$
- 15 k)  $-(C_1-C_4 \text{ alkyl})-CO_2R^4$ 
  - 1) heterocycle,
  - m)  $-(C_1-C_4 \text{ alkyl})$  heterocycle,
- 20 R<sup>13</sup> is independently selected at each occurrence from the group consisting of:
  - a) hydrogen
  - b) halogen,
  - c)  $C_1-C_4$  alkyl,
- d)  $C_1-C_4$  alkoxy,
  - e) methylenedioxy,
  - $f) -NO_2$
  - g) -CF3,
  - h) -SH,
- i)  $-S(0)_r (C_1 C_4 \text{ alkyl})$ ,

```
j) -CN,
            k) -OH,
            1) - NH<sub>2</sub>
            m) - NH(C_1-C_4 alkyl),
            n) -N(C_1-C_4 \text{ alkyl})_2,
            o) -NHC(=0)R^4, or
            p) -(CH_2)_{D}-CO_2R^4;
            \alpha) -C(=NH)NHR<sup>4</sup>
            r) -NHC(=NR^4)R^4
            s) -NHC (=NH) NHR^4
10
     R^{14} is:
            a) -CF3,
            b) -CHF2,
            c) -CH2F,
15
            d) -CH<sub>2</sub>Cl,
             e) -C(=0)OR^4,
             f) -C (=0) NR^{15}R^{16},
            g) -C(=0)R^4,
            h) -C(=0)COOR^4,
20
             i) -C(=0)C(=0)NR^{15}R^{16},
            j) -C(=0)C(=0)R^4,
             k) -CY^3Y^4COOR^4,
             1) -CY^3Y^4C (=0)NR^{15}R^{16}, or
             m) -CY^3Y^4C(=0)R^4;
25
```

WO 95/09634

 $\rm R^{15}$  and  $\rm R^{16}$  are independently selected at each occurrence from the group consisting of:

- a) hydrogen,
- 30 b)  $C_1-C_4$  alkyl,
  - c)  $-(C_1-C_4 \text{ alkyl})$  -aryl, where aryl is defined above,
  - d) C5-C7 cycloalkyl, or
  - e) phenyl, unsubstituted or substituted by  $R^{13}$ ,
  - f) C1-C4 alkoxy;

35

WO 95/09634

PCT/US94/11280

\_\_\_\_

```
\ensuremath{\text{R}^{15}} and \ensuremath{\text{R}^{16}} taken together to form a ring can also include:
             a)
             b)
  5
      R<sup>17</sup> is:
             a) hydrogen,
            b) C_1-C_4 alkyl,
10
             c) aryl, wherein aryl is defined above,
            d) -(C_1-C_4 \text{ alkyl}) -aryl, wherein aryl is defined above,
                   or
             e) C5-C7 cycloalkyl;
15
      R<sup>18</sup> is:
           a) hydrogen,
           b) -(C_1-C_5) alkyl, or
         c) -(C_1-C_5) haloalkyl,
           d) -(C1-C5) alkoxy;
20
     R<sup>19</sup> is:
           a) hydrogen,
           b) -(C_1-C_5) alkyl,
           c) halo, or
25
           d) -(C_1-C_5) haloalkyl,
           e) -NO_2
           f) -NR^4R^5,
```

30

g) -CN,

 $R^{20}$  is

h)  $-(C_1-C_5)$  alkoxy;

- a) hydrogen; or
- b)  $-N^2$  with amine protecting;

A is:

- 5 a)  $-BY^{1}Y^{2}$ , or
  - b)  $-C(=0)R^{14}$ ,
  - c)  $C(OH) R^{14}R^{18}$ ;

W is:

- 10 a) -O-,
  - b)  $-S(0)_{r}$ -,
  - c)  $-NR^4-$ , or
  - d)  $-NC(=0)R^4-;$
- 15  $y^1$  and  $y^2$  are:
  - a) -OH,
  - b) -F,
  - c)  $-NR^4R^5$ ,
  - d)  $C_1-C_8$  alkoxy, or
- when taken together  $Y^1$  and  $Y^2$  form:
  - e) a cyclic boron ester where said chain or ring contains from 2 to 20 carbon atoms and, optionally, 1-3 heteroatoms which can be N, S, or O,
- f) a cyclic boron amide where said chain or ring contains from 2 to 20 carbon atoms and, optionally, 1-3 heteroatoms which can be N, S, or O,
- g) a cyclic boron amide-ester where said chain or ring contains from 2 to 20 carbon atoms and, optionally, 1-3 heteroatoms which can be N, S, or O:

 $y^3$  and  $y^4$  are

35 a) -OH or

```
b) -F;
      n is 0 or 1;
      p is 0 to 3;
     q is 0 to 4;
      r is 0 to 2;
      t is 1 to 3;
      u is 1 to 4;
      v is 1 to 17.
 10
           Specifically preferred compounds of this invention
      include:
     Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn-C10H16 HCl
 15
     Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn(CH=NH)-C10H16
           HCl
     Hydrocinnamoy1-[N-(Phenethy1)-Gly]-boroOrn(CH=NH)-OH HC1
     Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroArg(CH3)-C10H16 HCl
     Hydrocinnamoy1-[N-(N(CH3)2)-Gly]-boroLys-C10H16 HC1
     Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroLys-OH HCl
20
     Hydrocinnamoy1-[N-(N(CH3)2)-Gly]-boroOrn-C10H16 HC1
     Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroOrn(CH=NH)-C10H16 HCl
     Methanesulfonyl-Sar-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl
     Methanesulfonyl-Sar-[N-(Phenethyl)-Gly]-boroLys-OH HCl
    Methanesulfonyl-Gly-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl
25
     Hydrocinnamoyl-[N-(3-(Trifluoromethyl)-Phenethyl)-Gly]-
          boroLys-C10H16 HCl
    \label{eq:hydrocinnamoyl-[N-(3-(Trifluoromethyl)-Phenethyl)-Gly]-} Hydrocinnamoyl-[N-(3-(Trifluoromethyl)-Phenethyl)-Gly]-
          boroLys-OH HCl
    Hydrocinnamoyl-[N-(3-(Methyl)-Phenethyl)-Gly]-boroLys-
30
          C10H16 HC1
    Hydrocinnamoyl-[N-(3-(Methyl)-Phenethyl)-Gly]-boroLys-OH
    Succinyl-[N-(3-(Methyl)-Phenethyl)-Gly]-boroLys-C10H16 HCl
```

```
Hydrocinnamoy1-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroLys-
         OH HCl
    Hydrocinnamoy1-[N-(2-(Cyclopropy1)-Phenethyl)-Gly]-boroLys-
         C10H16 HCl
    Hydrocinnamoy1-[N-(2-(Cyclopropy1)-Phenethy1)-Gly]-boroLys-
5
    Hydrocinnamoyl-[N-(2-(Cyclopropyl)-Phenethyl)-Gly]-
         boroOrn(CH=NH)-OH HCl
    Hydrocinnamoy1-[N-(2,2-(Diethyl)-Phenethyl)-Gly]-boroLys-
10
         C10H16 HCl
    Hydrocinnamoyl-Sar-Lys[C(=0)-C(=0)-OH]
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly[CH2)3-Br]-
         C10H16
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly[CH2)4)-Br]-
15
         C10H16
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroIrg-C10H16 HBr
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly[CH2)3-N3]-
         C10H16
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-borohomoIrg-C10H16 HBr
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly[CH2)4)-N3]-
20
         C10H16
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroOrn-C10H16 HCl
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-borohomoArg-C10H16 HCl
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroArg-C10H16 HCl
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroOrn(CH=NH)-C10H16
25
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroLys(CH=NH)-C10H16
    2-Benzyl-(N-Benzyl)-Sar-boroLys-C10H16 HCl
    2-Thiophenyl-Benzoyl-Sar-boroLys(CH=NH)-C10H16
30
    2-(Thiophenyl)-Benzoyl-Sar-boroIrg-C10H16 HBr
    2-(Thiophenyl)-Benzoyl-Sar-boroOrn-C10H16 HCl
    2-(Thiophenyl)-Benzoyl-Sar-boroOrn(CH=NH)-C10H16 HCl
     Pinanediol N-{N-methyl-N-[2-(Thiophenyl)-Benzoyl]Sar}-1-
         amido-5-thiocyanatobutane boronate
35
```

```
(2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroLys-C10H16 HCl
     Acetyl-Gly[N-(2-(Benzyl)-Benzyl)]-boroLys-C10H16 HCl
     Pinanediol N-{N-methyl-N-[2-(pyrrol-1-ylmethyl)-
          Benzyl]glycyl]-1-amido-5-aminopentaneboronate,
  5
          hydrochloride salt
     N-{N-methyl-N-[2-(pyrrol-1-ylmethyl)-Benzyl]glycyl}-1-
          amido-5-aminopentaneboronic acid, hydrochloride salt
          2-(2-(Trifluoromethyl)-Benzyl)-Benzoyl-Sar-Lys-C(=0)-
          NHNH2 2 HC1
 10
     2-(Benzyl)-Benzoyl-Sar-Lys-C(=O)-NHNH2 2 HCl
     [3-(Trifluoromethyl)-Benzyl]-Benzoyl-Sar-boroLys-C10H16 HCl
     3-(3-(Chloro)-Benzyl)-Benzoyl-Sar-boroLys-C10H16 HCl
     Hydrocinnamoy1-Sar-Lys(Z)-C(=O)-O-(CH2)2-NH(Z)
     Hydrocinnamoyl-Sar-Lys-C(=O)-O-(CH2)2-NH2 2 HCl
15
     Hydrocinnamoyl-Sar-Lys(Z)-C(=O)-OCH3
     Hydrocinnamoyl-Sar-Lys-C(=0)-OCH3 HCl
     Hydrocinnamoyl-Sar-Lys-C(=O)-CH3 HCl
     Hydrocinnamoyl-Sar-Lys(2)-H
     Hydrocinnamoyl-Sar-NHCH(CH2OH)(CH2)4-NH(Z)
20
     Hydrocinnamoy1-Sar-NHCH(CH2OH)(CH2)4-NH2
     Hydrocinnamoyl-Sar-Lys[CH(OH)(OCH3)-C(=O)-OCH3] HCl
     Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn-C10H16 HCl
     Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl
     Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn(CH=NH)-OH HCl
    Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn(CH=NH)-C10H16
25
          HC1
    Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys(CH=NH)-C10H16
         HC1
    Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys(CH=NH)-OH HCl
    Phenoxyacetyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl
30
    Thiophenacetyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl
     Phenoxyacetyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl
    Thiophenacetyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl
    Hydrocinnamoyl-[N-(N-(Methyl)-Phenyl)-Gly]-boroLys-C10H16
35
         HC1
```

```
Hydrocinnamoyl-[N-(N-(Methyl)-Phenyl)-Gly]-boroLys-OH HCl
    Hydrocinnamoy1-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-C10H16
         HC1
    Glutaryl-[N-(Phenethyl)-Gly]-boroLys-OH HCl
    Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-C10H16
    Methyl Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-
         C10H16 HCl
    Glutary1(3,3-Dimethy1)-[N-(Phenethy1)-Gly]-boroLys-OH
    Methyl Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-
         OH HCl
10
    Boc-Asp-[N-(Phenethyl)-Gly]-boroLys-C10H16
    Boc-Glu-[N-(Phenethyl)-Gly]-boroLys-C10H16
    Boc-Glu(OCH3) - [N-(Phenethyl)-Gly]-boroLys-C10H16 HCl
    Boc-Glu-[N-(Phenethyl)-Gly]-boroLys-OH
    Hydrocinnamoy1-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-OH HCl
    Methanesylfonyl-Gly-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-
         C10H16 HCl
    Methanesulfonyl-Gly-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-OH
    Hydrocinnamoyl-[N-(Succinyl)-Gly]-boroLys-C10H16
20
    Hydrocinnamoy1-[N-(Methyl Succinyl)-Gly]-boroLys-C10H16 HCl
    Succinyl-[N-(Phenethyl)-Gly]-boroLys-OH
    Methyl Succinyl-[N-(Phenethyl)-Gly]-boroLys-OH HCl
    Glutaryl-[N-(Phenethyl)-Gly]-boroLys-C10H16
    Methyl Glutaryl-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl
25
    Methyl Glutaryl-[N-(Phenethyl)-Gly]-boroLys-OH HCl
    Hydrocinnamoy1-[N-(2-(Cyclopropy1)-Phenethy1)-Gly]-boroArg-
         C10H16 HCl
    Hydrocinnamoy1-[N-(2-(Cyclopentyl)-Phenethyl)-Gly]-boroLys-
30
         C10H16 HCl
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroArg-OH HCl
    Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroLys-
         C10H16 HCl
    Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroOrn-
         C10H16 HC1
35
```

```
Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroArg-
          C10H16 HC1
     Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
          boroOrn(CH=NH)-C10H16 HCl
     Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroArg-
          OH HCl
     Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
          boroOrn(CH=NH)-OH HCl
     Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroOrn-
10
          C10H16 HCl
     Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroLys-
          C10H16 HC1
     Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-
          boroOrn-C10H16 HCl
     Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-
15
          boroLys-C10H16 HCl
     Hydrocinnamoy1-{N-[2,2-(Dimethyl)-2-(3,5-dimethylphenyl)-}
          ethyl]-Gly}-boroArg-C10H16 HCl
     Hydrocinnamoyl-{N-[2,2-(Dimethyl)-2-(3,5-dimethylphenyl)-
20
          ethyl]-Gly}-boroOrn(CH=NH)-C10H16 HCl
     Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-
          boroArg-C10H16 HCl
    Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-
          boroOrn(CH=NH)-C10H16 HCl
25
    Hydrocinnamoyl-[N-(Cyclohexyl)-Gly]-boroLys-C10H16 HC1
    Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl
    Hydrocinnamoyl-[N-(Cyclohexyl)-Gly]-boroLys-OH HCl
    Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn(CH=NH)-C10H16
          HCl
30
    Hydrocinnamoyl-[N-(2-(Cyclopentyl)-Phenethyl)-Gly]-
          boroOrn(CH=NH)-C10H16 HCl
    Hydrocinnamoyl-[N-(2-(Cyclopentyl)-Phenethyl)-Gly]-boroArg-
         C10H16 HC1
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroLys-OH
35
    [N-(-C(O)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-boroLys-OH
```

```
[N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-boroLys-OH
     [N-(-C(O)N(CH_3)CH_2Ph)-N-Ph]Gly-boroLys-OH
     [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(0)Ph-3-SPh-2-OCH_3)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(O)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(O)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(O)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-boroLys-OH
10
     [N-(-C(O)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroLys-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-boroLys-OH
     [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroLys-OH
15
     [N-(-C(O)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroLys-OH
     [N-(-C(O)Ph)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-CH<sub>2</sub>Ph)]Gly-boroLys-OH
     [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-boroLys-OH
20
     [N-(-C(O)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-boroLys-OH
      [N-(-C(O)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-boroLys-C<sub>10</sub>H<sub>16</sub>
25
      [N-(-C(O)N(CH_3)CH_2Ph)-N-Ph]Gly-boroLys-C_{10}H_{16}
      [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0) Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(O)Ph-3-SPh-2-OCH_3)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
30
      [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0) Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
      [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
35
```

```
[N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)Ph)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
5
     [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph-4-Cl)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-bor\'eLys-C_{10}H_{16}
10
     [N-(-C(O)CH_3](D)-Phe[N-(CH_3)]Gly-boroLys-C_{10}H_{16}
     [N-(SO_2CH_3](D)-Phe[N-(CH_3)]Gly-boroLys-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-Lys-OCH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_3)]Gly-boroArg-OH
     [N-(-C(0)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-boroArg-OH
15
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroOrn(N-
20
           methylamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(N-methylamidino)-
           C10H16
      [N-(-C(0)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-boroOrn(N-
           methylamidino)-C10H16
25
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-
30
            OH
      [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-C_{10}H_{16}
      [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroOrn(formamidino)-
            C10H16
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3, 4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-
            boroOrn(formamidino)-C10H16
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-boroLys-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph)]-Gly-
 5
            boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-CH_3)]-Gly-boroLys-
            OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>Ph-3-CH<sub>3</sub>)]-Gly-
10
            boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph)]-Gly-
            boroArg-OH .
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-NO_2)]-Gly-boroLys-
            OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-NO_2)]-Gly-
15
            boroLys-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CH<sub>2</sub>Ph-3,5-(CH<sub>3</sub>)<sub>2</sub>)]-Gly-
            boroArg-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>Ph-3,5-(CH<sub>3</sub>)<sub>2</sub>)]-Gly-
20
            boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph)]-Gly-
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-CH_3)]-Gly-
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-CH_3)]-Gly-
25
            boroOrn(formamidino)-OH
             Illustrative of the preferred compounds of this
      invention are the following:
30
```

 $[N-(-C(O)(CH_2)_2Ph)-N-(C_2H_5)]Gly-boroLys-OH\\ [N-(-C(O)(CH_2)_2Ph)-N-(N-C_3H_7)]Gly-boroLys-OH\\ [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-boroLys-OH\\ [N-(-C(O)(CH_2)_2Ph)-N-(OCH_3)]Gly-boroLys-OH\\ [N-(-C(O)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-boroLys-OH$ 

-27**-**

35

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-boroLys-OH
       [N-(-C(O)(CH_2)_2Ph)-N-(NHBoc)]Gly-boroLys-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>H)]Gly-boroLys-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)]Gly-boroLys-OH
       [N-(-C(O)(OCH<sub>2</sub>)Ph)](D)-Phe[N-(CH<sub>3</sub>)]Gly-boroLys-OH
       [N-(-C(O)(OCH_2)Ph)](D)-Phe[N-(CH_3)]Ala-boroLys-OH
       [N-(-C(0)(OCH_2)Ph)](D)-Phe[N-(Ph)]Gly-boroLys-OH
       [N-(-C(O)(OCH_2)Ph)](D)-Phe[N-(CH_2Ph)]Gly-boroLys-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-boroLys-C<sub>10</sub>H<sub>16</sub>
 10
       [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-boroLys-C_{10}H_{16}
       [N-(-C(0)(CH_2)_2Ph)-N-(OH)]Gly-boroLys-C_{10}H_{16}
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-boroLys-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-boroLys-C_{10}H_{16}
       [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-boroLys-C_{10}H_{16}
15
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-boroLys-C_{10}H_{16}
       [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-boroLys-C_{10}H_{16}
       [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-boroLys-C_{10}H_{16}
       (N-CO_2CH_2Ph) [Leu-Ser(O<sup>t</sup>Bu)-Asn]<sub>4</sub>-[N-(CH_3)]Gly-boroLys-C_{10}H_{16}
              [Sequence No. 1]
20
       (H) - [Leu-Ser (O<sup>t</sup>Bu) -Asn]<sub>4</sub>-[N-(CH<sub>3</sub>)]Gly-boroLys-C<sub>10</sub>H<sub>16</sub>
              [Sequence No. 2]
       (H)-[Leu-Ser-Asn]4-[N-(CH3)]Gly-boroLys-C10H16 [Sequence No.
             31
       [N-(-C(0)(OCH_2Ph)](D)-Phe[N-(CH_3)]Gly-boroLys-C_{10}H_{16}
25
      [N-(-C(0)(CH_3)](D)-(\beta eyclohexyl)Ala[N-(CH_3)]Gly-boroLys-
             C10H16
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Lys-CF<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Lys-CF_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Lys-CF<sub>3</sub>
30
      [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Lys-CF_3
      [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Lys-CF_3
      [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Lys-CF_3
      [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Lys-CF_3
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Lys-CF_3
35
      [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Lys-CF_3
```

```
[N-(-C(O)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Lys-CF_3
     [N-(-C(O)Ph-3-CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Lys-CF<sub>3</sub>
     [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Lys-CF_3
     [N-(-C(O)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Lys-CF_3
     [N-(-C(O)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Lys-CF_3
5
     [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Lys-CF_3
     [N-(-C(O)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Lys-CF_3
     [N-(-C(O)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Lys-CF_3
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-Lys-CF<sub>3</sub>
     [N-(-C(O)(CH_2)_2Ph)-N-(OCH_3)]Gly-Lys-CF_3
10
     [N-(-C(O)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Lys-CF_3
     [N-(-C(O)(CH_2)_2Ph)-N-(NH_2)]Gly-Lys-CF_3
     [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Lys-CF_3
     [N-(-C(O)(CH_2)_2Ph)-N-(NHBoc)]Gly-Lys-CF_3
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Lys-CF_3
15
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Lys-CF_3
     [N-(-C(O)CH_2Ph)-N-(CH_3)]Gly-Lys-CF_3
     [N-(-C(O)(CH_2)_3Ph)-N-(CH_3)]Gly-Lys-CF_3
     [N-(-C(O)Ph)-N-(CH_3)]Gly-Lys-CF_3
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-CH<sub>2</sub>Ph)]Gly-Lys-CF<sub>3</sub>
20
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3,4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-Lys-CF<sub>3</sub>
      [N-(-C(O)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Lys-CF_3
      [N-(-C(O)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Lys-CF_3
      [N-(-C(O)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-Lys-CF_3
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Lys-OCH_3
25
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Lys-OCH<sub>3</sub>
      [N-(-C(O)N(CH_3)CH_2Ph)-N-Ph]Gly-Lys-OCH_3
      [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Lys-OCH_3
      [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Lys-OCH_3
      [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Lys-OCH_3
30
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Lys-OCH_3
      [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Lys-OCH_3
      [N-(-C(O)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Lys-OCH_3
      [N-(-C(O)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Lys-OCH_3
      [N-(-C(O)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Lys-OCH_3
35
```

```
[N-(-C(0)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-Lys-OCH_3
        [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Lys-OCH_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-Lys-OCH<sub>3</sub>
        [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Lys-OCH_3
  5
        [N-(-C(O)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Lys-OCH_3
        [N-(-C(0)(CH_2)_2Ph)-N-(OH)]Gly-Lys-OCH_3
        [N-(-C(0)(CH_2)_2Ph)-N-(OCH_3)]Gly-Lys-OCH_3
        [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Lys-OCH_3
        [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Lys-OCH_3
 10
        [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Lys-OCH_3
        [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Lys-OCH<sub>3</sub>
        [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Lys-OCH_3
        [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Lys-OCH_3
        [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Lys-OCH_3
 15
        [N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-Lys-OCH_3
        [N-(-C(O)Ph)-N-(CH_3)]Gly-Lys-OCH_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-CH<sub>2</sub>Ph)]Gly-Lys-OCH<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3, 4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-Lys-OCH<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-Lys-OCH<sub>3</sub>
20
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-CH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Lys-OCH<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Lys-OCH<sub>3</sub>
        [N-(-C(O)(CH_2)_2Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-Lys-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Lys-CO<sub>2</sub>CH<sub>3</sub>
25
       [N-(-C(O)N(CH_3)CH_2Ph)-N-Ph]Gly-Lys-CO_2CH_3
       [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Lys-CO_2CH_3
       [N-(-C(O)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-Lys-CO_2CH_3
35
       [N-(-C(O)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Lys-CO_2CH_3
```

PCT/US94/11280

```
[N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Lys-CO_2CH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Lys-CO_2CH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Lys-CO_2CH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-Lys-CO_2CH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(OCH_3)]Gly-Lys-CO_2CH_3
5
     [N-(-C(O)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Lys-CO_2CH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Lys-CO_2CH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Lys-CO_2CH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-Lys-CO_2CH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Lys-CO_2CH_3
10
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Lys-CO_2CH_3
     [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
     [N-(-C(O)(CH_2)_3Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
     [N-(-C(O)Ph)-N-(CH_3)]Gly-Lys-CO_2CH_3
     [N-(-C(O)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Lys-CO_2CH_3
15
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3,4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-Lys-CO<sub>2</sub>CH<sub>3</sub>
     [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Lys-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Lys-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-Lys-CO_2CH_3
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-boroArg-OH
20
      [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-boroArg-OH
      [N-(-C(O)N(CH_3)CH_2Ph)-N-Ph]Gly-boroArg-OH
      [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroArg-OH
      [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroArg-OH
      [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroArg-OH
25
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroArg-OH
      [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroArg-OH
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroArg-OH
      [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroArg-OH
      [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-boroArg-OH
30
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-boroArg-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(OH)]Gly-boroArg-OH
35
```

```
[N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-boroArg-OH
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-boroArg-OH
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-boroArg-OH
       [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-boroArg-OH
  5
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-boroArg-OH
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>H)]Gly-boroArg-OH
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)]Gly-boroArg-OH
        [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroArg-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroArg-OH
10
       [N-(-C(0)Ph)-N-(CH_3)]Gly-boroArg-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-CH<sub>2</sub>Ph)]Gly-boroArg-OH
       [N-(-C(0)(CH_2)_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-boroArg-OH
       [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-boroArg-OH
       [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-boroArg-OH
15
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-boroArg-OH
       [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-boroArg-C_{10}H_{16}
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)N(CH<sub>3</sub>)CH<sub>2</sub>Ph)-N-Ph]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
20
       [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
25
       [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(O)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(O)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(i-C<sub>3</sub>H<sub>7</sub>)]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
30
       [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroArg-C_{10}H_{16}
       [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-boroArg-C_{10}H_{16}
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
35
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(N(CH<sub>3</sub>)<sub>2</sub>)]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-boroArg-C<sub>10</sub>H<sub>16</sub>
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
 5
     [N-(-C(0) Ph)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
10
     [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(O)CH_3](D)-Phe[N-(CH_3)]Gly-boroArg-C_{10}H_{16}
     [N-(-C(O)CH_3](D)-Phe[N-(CH_3)]Gly-boroIrg-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-Arg-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Arg-CF_3
15
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Arg-CCF<sub>3</sub>
      [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Arg-CF_3
      [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Arg-CF_3
      [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Arg-CF_3
      [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Arg-CF_3
20
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Arg-CF_3
      [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Arg-CF_3
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Arg-CF_3
      [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Arg-CF_3
     [N-(-C(0) Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Arg-CF_3
25
      [N-(-C(0)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-Arg-CF_3
      [N-(-C(O)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Arg-CF_3
      [N-(-C(O)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Arg-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Arg-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Arg-CF_3
30
      [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-Arg-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_3)]Gly-Arg-CF_3
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-Arg-CF<sub>3</sub>
      [N-(-C(O)(CH_2)_2Ph)-N-(NH_2)]Gly-Arg-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Arg-CF_3
35
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Arg-CF<sub>3</sub>
        [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Arg-CF_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)]Gly-Arg-CF<sub>3</sub>
        [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Arg-CF_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Arg-CF<sub>3</sub>
   5
        [N-(-C(0)Ph)-N-(CH_3)]Gly-Arg-CF_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-CH<sub>2</sub>Ph)]Gly-Arg-CF<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3,4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-Arg-CF<sub>3</sub>
        [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Arg-CF_3
 10
        [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Arg-CF_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Arg-CF<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Arg-OCH<sub>3</sub>
        [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Arg-OCH_3
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Arg-OCH<sub>3</sub>
 15
       [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Arg-OCH_3
        [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Arg-OCH_3
        [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Arg-OCH_3
       [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Arg-OCH_3
       [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Arg-OCH_3
       [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Arg-OCH_3
 20
       [N-(-C(O)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Arg-OCH_3
       [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Arg-OCH_3
       [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Arg-OCH_3
       [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Arg-OCH_3
25
       [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Arg-OCH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Arg-OCH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Arg-OCH_3
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>6</sub>H<sub>12</sub>)]Gly-Arg-OCH<sub>3</sub>
      [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-Arg-OCH_3
30
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_3)]Gly-Arg-OCH_3
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Arg-OCH_3
      [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Arg-OCH_3
      [N-(-C(O)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Arg-OCH_3
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Arg-OCH<sub>3</sub>
35
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Arg-OCH_3
```

```
[N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Arg-OCH_3
      [N-(-C(O)CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Arg-OCH<sub>3</sub>
      [N-(-C(O)(CH_2)_3Ph)-N-(CH_3)]Gly-Arg-OCH_3
      [N-(-C(O)Ph)-N-(CH_3)]Gly-Arg-OCH_3
      [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Arg-OCH_3
 5
      [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Arg-OCH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-Arg-OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-CH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Arg-OCH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-Arg-OCH_3
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
10
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Arg-CO_2CH_3
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(O)N(CH<sub>3</sub>)CH<sub>2</sub>Ph)-N-Ph]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Arg-CO_2CH_3
      [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
15
       [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(O)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(O)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
20
       [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(O)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Arg-CO_2CH_3
       [N-(-C(O)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Arg-CO_2CH_3
       [N-(-C(O)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Arg-CO_2CH_3
25
       [N-(-C(O)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Arg-CO_2CH_3
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph)-N-(NH_2)]Gly-Arg-CO_2CH_3
30
       [N-(-C(O)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Arg-CO_2CH_3
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Arg-CO_2CH_3
       [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(O)CH_2Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
35
```

```
[N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(0)Ph)-N-(CH<sub>3</sub>)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-CH<sub>2</sub>Ph)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3, 4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-Arg-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-Arg-CO_2CH_3
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-boroPhe(mCN)-OH
       [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-boroPhe(mCN)-OH
10
       [N-(-C(O)N(CH_3)CH_2Ph)-N-Ph]Gly-boroPhe(mCN)-OH
       [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroPhe(mCN)-OH
15
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-OH
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-boroPhe(mCN)-OH
20
      [N-(-C(0)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-boroPhe(mCN)-OH
       [N-(-C(O)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-boroPhe(mCN)-OH
       [N-(-C(O)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroPhe(mCN)-OH
25
      [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-boroPhe(<math>mCN)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-boroPhe(mCN)-OH
30
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-boroPhe(mCN)-OH
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>H)]Gly-boroPhe(mCN)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)]Gly-boroPhe(mCN)-OH
       [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-OH
      [N-(-C(O)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-OH
35
      [N-(-C(0)Ph)-N-(CH_3)]Gly-boroPhe(mCN)-OH
```

```
[N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroPhe(mCN)-OH
     [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroPhe(mCN)-OH
     [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-boroPhe(mCN)-OH
     [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-boroPhe(mCN)-OH
     [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-boroPhe(mCN)-OH
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph] Gly-boroPhe(mCN)-C_{10}H_{16}
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-boroPhe(mCN)^{\frac{1}{2}}C<sub>10</sub>H<sub>16</sub>
     [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-boroPhe(mCN)-C_{10}H_{16}
     [N-(-C(O)Ph-3-CH=CHPh)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-C<sub>10</sub>H<sub>16</sub>
10
      [N-(-C(O)Ph-3-CH<sub>2</sub>CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-C<sub>10</sub>H<sub>16</sub>
      [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(O)Ph-4-CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-C<sub>10</sub>H<sub>16</sub>
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
15
      [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-boroPhe(mCN)-C<sub>10</sub>H<sub>16</sub>
20
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(OH)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-boroPhe(mCN)-C<sub>10</sub>H<sub>16</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)] \\ Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-boroPhe(mCN)-C_{10}H_{16}
25
      [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
30
      [N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(O)Ph)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-C<sub>10</sub>H<sub>16</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
      [N-(-C(0)(CH_2)_2Ph-4-Cl)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
35
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-CH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-boroPhe(mCN)-C_{10}H_{16}
             [N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CF_3
       [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Phe(mCN)-CF_3
  5
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Phe(mCN)-CF<sub>3</sub>
       [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Phe(mCN)-CF_3
       [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Phe(mCN)-CF_3
       [N-(-C(0) Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Phe(mCN)-CF_3
10
      [N-(-C(0) Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(O)Ph-4-CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Phe(\piCN)-CF<sub>3</sub>
       [N-(-C(0) Ph-2-CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CF_3
       [N-(-C(0) Ph-3-CH_2Ph) -N-(CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Phe(mCN)-CF_3
15
      [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(O)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Phe(mCN)-CF_3
20
      [N-(-C(0)(CH_2)_2Ph)-N-(OH)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-Phe(mCN)-CF<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-Phe(mCN)-CF<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Phe(\pi CN)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Phe(\pi CN)-CF_3
25
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Phe(mCN)-CF<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-Phe(mCN)-CF_3
30
      [N-(-C(0)Ph)-N-(CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Phe(mCN)-CF_3
      [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Phe(\pi CN)-CF_3
      [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Phe(mCN)-CF_3
35
      [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-Phe(mCN)-CF_3
```

```
[N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Phe(mCN)-OCH_3
     [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0) Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(O)Ph-4-CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Phe(mCN)-OCH<sub>3</sub>
     [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
10
     [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Phe(mCN)-OCH_3
15
     [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Phe(mCN)-OCH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-Phe(mCN)-OCH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(OCH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Phe(mCN)-OCH_3
20
     [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Phe(mCN)-OCH<sub>3</sub>
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Phe(mCN)-OCH_3
25
     [N-(-C(O)CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)Ph)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(O)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
30
     [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-Phe(mCN)-OCH_3
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Phe(mCN)-CO_2CH_3
35
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Phe(mCN)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)N(CH_3)CH_2Ph)-N-Ph]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0) Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
  5
       [N-(-C(0) Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
10
       [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Phe(mCN)-CO_2CH_3
15
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Phe(mCN)-CO_2CH_3
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-Phe(mCN)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(0)(CH_2)_2Ph)-N-(OCH_3)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-Phe(mCN)-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Phe(mCN)-CO_2CH_3
20
      [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Phe(mCN)-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(O)CH_2Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
25
      [N-(-C(O)(CH_2)_3Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(O)Ph)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3,4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-Phe(mCN)-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
30
      [N-(-C(O)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-Phe(mCN)-CO_2CH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-boroOrn(N-methylamidino)-
            OH
      [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-boroOrn(N-methylamidino)-OH
35
     [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-boroOrn(N-methylamidino)-OH
```

```
[N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroOrn(N-
          methylamidino)-OH
     [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-
          methylamidino)-OH
     [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroOrn(N-CH_3)
5
          methylamidino)-OH
     [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroOrn(N-
          methylamidino)-OH
     [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-
10
     [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-
     [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-
     [N-(-C(0) Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-boroOrn(N-1)
15
          methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-boroOrn(N-methylamidino)-
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(i-C<sub>3</sub>H<sub>7</sub>)]Gly-boroOrn(N-methylamidino)-
20
           OH
     [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(O)(CH_2)_2Ph)-N-(OCH_3)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-boroOrn(N-methylamidino)-
25
           OH
     [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-boroOrn(N-
           methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-boroOrn(N-methylamidino)-
30
           OH
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-boroOrn(N-
           methylamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)]Gly-boroOrn(N-
           methylamidino)-OH
35
```

```
[N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(0)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(O)Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroOrn(N-methylamidino)-OH
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3,4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-boroOrn(N-
 5
           methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph-4-Cl)-N-(CH_3)]Gly-boroOrn(N-
           methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-boroOrn(N-
           methylamidino) -OH
10
     [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-boroOrn(N-CH_3)
           methylamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-boroOrn(N-methylamidino)-C_{10}H_{16}
      [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-boroOrn(N-methylamidino)-
15
           C10H16
      [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroOrn(N-
           methylamidino)-C<sub>10</sub>H<sub>16</sub>
      [N-(-C(0) Ph-3-CH<sub>2</sub>CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(N-
           methylamidino)-C10H16
      [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroOrn(N-
20
           methylamidino)-C10H16
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroOrn(N-
           methylamidino)-C10H16
      [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-
25
           C10H16
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-
           C10H16
      [N-(-C(O)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-
           C10H16
      [N-(-C(O)Ph-3-CH_2Ph-2+CF_3)-N-(CH_3)]Gly-boroOrn(N-
30
           methylamidino)-C10H16
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-boroOrn(N-methylamidino)-
           C10H16
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-boroOrn(N-methylamidino)-
35
           C10H16
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(i-C<sub>3</sub>H<sub>7</sub>)]Gly-boroOrn(N-methylamidino)-
           C10H16
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>6</sub>H<sub>12</sub>)]Gly-boroOrn(N-methylamidino)-
           C10H16
     [N-(-C(O)(CH_2)_2Ph)-N-(OH)]Gly-boroOrn(N-methylamidino)-
           C_{10}H_{16}
     [N-(-C(0)(CH_2)_2Ph)-N-(OCH_3)]Gly-boroOrn(N-methylamidino)-
           C10H16
     [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-boroOrn(N-methylamidino)-
10
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-boroOrn(N-methylamidino)-
           C10H16
     [N-(-C(O)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-boroOrn(N-
           methylamidino)-C10H16
     [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-boroOrn(N-methylamidino)-
15
           C10H16
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-boroOrn(N-
           methylamidino)-C10H16
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-boroOrn(N-
20
           methylamidino)-C10H16
     [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-C_{10}H_{16}
     [N-(-C(0)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(N-methylamidino)-
           C10H16
     [N-(-C(O)Ph)-N-(CH_3)]Gly-boroOrn(N-methylamidino)-C<sub>10</sub>H<sub>16</sub>
     [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroOrn(N-methylamidino)-
25
           C10H16
     [N-(-C(0)(CH_2)_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-boroOrn(N-CH_3)
           methylamidino)-C10H16
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-boroOrn(N-
30
           methylamidino)-C<sub>10</sub>H<sub>16</sub>
      [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-boroOrn(N-
            methylamidino)-C10H16
      [N-(-C(0)(CH_2)_2Ph-4-OCH_3)-N-(CH_3)]Gly-boroOrn(N-
           methylamidino)-C10H16
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CF_3
35
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
        [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-Orn(N-methylamidino)-CF_3
        [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Orn(N-methylamidino)-CF_3
        [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Orn(N-methylamidino)-CF_3
  5
        [N-(-C(0)Ph-3-CH<sub>2</sub>CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
               CF<sub>3</sub>
        [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Orn(N-methylamidino)-
        [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-
 10
        [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CF_3
        [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CF_3
        [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CF_3
        [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Orn(N-
15
               methylamidino)-CF3
        [N-(-C(0)CH_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-Orn(N-methylamidino)-
               CF2
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
20
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(i-C<sub>3</sub>H<sub>7</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>6</sub>H<sub>12</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
25
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(N(CH<sub>3</sub>)<sub>2</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-Orn(N-methylamidino)-CF_3
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>H)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
30
              CF<sub>3</sub>
       [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CF_3
       [N-(-C(0)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-CF<sub>3</sub>
       [N-(-C(0)Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CF_3
       [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Orn(N-methylamidino)-CF_3
```

```
[N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(N-
            methylamidino)-CF3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
      [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Orn(N-methylamidino)-
 5
            CF3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
            CF<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
10
      [N-(-C(O)(CH_2)_2Ph)-N-Ph]Gly-Orn(N-methylamidino)-OCH_3
      [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Orn(N-methylamidino)-OCH_3
      [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Orn(N-methylamidino)-
             OCH<sub>3</sub>
      [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-
15
             OCH<sub>3</sub>
      [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Orn(N-methylamidino)-
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-
20
             OCH<sub>3</sub>
      [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-OCH_3
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-OCH_3
      [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-OCH_3
      [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Orn(N-
25
             methylamidino) - OCH3
      [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(N-methylamidino)-
             OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
30
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>6</sub>H<sub>12</sub>)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Orn(N-methylamidino)-OCH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
35
```

```
[N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Orn(N-methylamidino)-OCH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-Orn(N-methylamidino)-OCH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Orn(N-methylamidino)-OCH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Orn(N-methylamidino)-
 5
             OCH<sub>3</sub>
      [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-OCH_3
       [N-(-C(0)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-OCH<sub>3</sub>
       [N-(-C(O)Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-OCH_3
       [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Orn(N-methylamidino)-OCH_3
10
       [N-(-C(0)(CH_2)_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-Orn(N-
             methylamidino) - OCH3
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-CH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
15
             OCH<sub>3</sub>
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
             OCH<sub>3</sub>
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-Orn(N-methylamidino)-
20
             CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-Orn(N-methylamidino)-CO_2CH_3
      [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Orn(N-methylamidino)-CO_2CH_3
      [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Orn(N-methylamidino)-
             CO<sub>2</sub>CH<sub>3</sub>
25
      [N-(-C(0) Ph-3-CH<sub>2</sub>CH<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
      [N-(-C(0) Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Orn(N-methylamidino)-
             CO2CH2
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-
30
      [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-
            CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0) Ph-2-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-
            CO<sub>2</sub>CH<sub>3</sub>
```

```
[N-(-C(0) Ph-3-CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-
            CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Orn(N-CH_3)
            methylamidino)-CO2CH3
      [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(N-methylamidino)-
            CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Orn(N-methylamidino)-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Orn(N-methylamidino)-
             CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Orn(N-methylamidino)-
10
           CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Orn(N-methylamidino)-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph)-N-(OH)]Gly-Orn(N-methylamidino)-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_3)]Gly-Orn(N-methylamidino)-CO_2CH_3
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Orn(N-methylamidino)-
15
             CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Orn(N-methylamidino)-CO_2CH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(N(CH<sub>3</sub>)<sub>2</sub>)]Gly-Orn(N-methylamidino)-
             CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Orn(N-methylamidino)-
20
             CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Orn(N-methylamidino)-
       [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Orn(N-methylamidino)-
25
             CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CO_2CH_3
       [N-(-C(O)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)Ph)-N-(CH_3)]Gly-Orn(N-methylamidino)-CO_2CH_3
       [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Orn(N-methylamidino)-CO_2CH_3
       [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(N-
30
              methylamidino)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
              CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Orn(N-methylamidino)-
 35
              CO<sub>2</sub>CH<sub>3</sub>
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(N-methylamidino)-
             CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(0)(CH_2)_2Ph)-N-Ph]Gly-boroOrn(formamidino)-OH
       [N-(-C(0)N(CH<sub>3</sub>)CH<sub>2</sub>Ph)-N-Ph]Gly-boroOrn(formamidino)-OH
  5
       [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
       [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
       [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroOrn(formamidino)-
       [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-
 10
       [N-(-C(0)Ph-4-CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
       [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
       [N-(-C(0)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
       [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-
 15
             boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroOrn(formamidino)-OH
20
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-boroOrn(formamidino)-OH]
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_3)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-boroOrn(formamidino)-OH
25
      [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-boroOrn(formamidino)-OH
      [N-(-C(O)(CH_2)_3Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
      [N-(-C(O)Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-OH
30
      [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3,4-Cl<sub>2</sub>)-N-(CH<sub>3</sub>)]Gly-
           boroOrn(formamidino)-OH
     [N-(-C(O)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-boroOrn(formamidino)-OH
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-CH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-
35
            OH
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-boroOrn(formamidino)-
             C10H16
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
      [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-boroOrn(formamidino)-C_{10}H_{16}
      [N-(-C(O)Ph-3-CH=CHPh)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-
             C10H16
      [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-boroorn(formamidino)-
             C10H16
      [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-boroOrn(formamidino)-
10
             C10H16
      [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-
              C10H16
       [N-(-C(O)Ph-3-CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-C_{10}H_{16}
       [N-(-C(O)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-
15
              boroOrn(formamidino)-C10H16
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-boroOrn(formamidino)-
              C10H16
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(i-C<sub>3</sub>H<sub>7</sub>)]Gly-boroOrn(formamidino)-
20
              C10H16
       [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-boroOrn(formamidino)-C_{10}H_{16}
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>2</sub>Ph)]Gly-boroOrn(formamidino)-
25
              C10H16
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(N(CH<sub>3</sub>)<sub>2</sub>)]Gly-boroOrn(formamidino)-
              C10H16
       [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-boroOrn(formamidino)-C_{10}H_{16}
       [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-boroOrn(formamidino)-
30
              C<sub>10</sub>H<sub>16</sub>
       [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-boroOrn(formamidino)-
              C10H16
       [N-(-C(O)CH_2Ph)-N-(CH_3)]Gly-boroOrn(formamidino)-C_{10}H_{16}
       [N-(-C(0)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
35
```

```
[N-(-C(O)Ph)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-C<sub>10</sub>H<sub>16</sub>
        [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-boroOrn(formamidino)-C_{10}H_{16}
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-Cl)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-
               C10H16
   5
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-CH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-
              C<sub>10</sub>H<sub>16</sub>
        [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-boroOrn(formamidino)-
              C10H16
        [N-(-C(O)(CH_2)_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
        [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Orn(formamidino)-CF<sub>3</sub>
  10
        [N-(-C(O)(CH_2)_2Ph)-N-Ph]Gly-Orn(formamidino)-CF_3
        [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Orn(formamidino)-CF_3
        [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
       [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF<sub>3</sub>
       [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Orn(formamidino)-CF<sub>3</sub>
  15
       [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF<sub>3</sub>
       [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
       [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
       [N-(-C(O)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
 20
       [N-(-C(O)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Orn(formamidino)-CF<sub>3</sub>
       [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
       [N-(-C(0)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Orn(formamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Orn(formamidino)-CF<sub>3</sub>
       [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Orn(formamidino)-CF<sub>3</sub>
 25
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(NH_2)]Gly-Orn(formamidino)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Orn(formamidino)-CF<sub>3</sub>
30
      [N-(-C(0)(CH_2)_2Ph)-N-(NHBoc)]Gly-Orn(formamidino)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CO_2H)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(0)CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
35
      [N-(-C(0)(CH_2)_3Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
```

```
[N-(-C(0)Ph)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
      [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Orn(formamidino)-CF_3
      [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
      [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Orn(formamidino)-CF_3
      [N-(-C(0)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-CF<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2)_2Ph]Gly-Orn(formamidino)-OCH<sub>3</sub>
      [N-(-C(O)(CH_2)_2Ph)-N-Ph]Gly-Orn(formamidino)-OCH_3
      [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Orn(formamidino)-OCH_3
10
      [N-(-C(O)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
      [N-(-C(O)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
      [N-(-C(O)Ph-3-SPh-3-OCH_3)-N-(CH_3)] Gly-Orn(formamidino)-OCH_3
      [N-(-C(O)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)] \\ Gly-Orn(formamidino)-OCH_3
      [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
15
      [N-(-C(0)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-OCH<sub>3</sub>
      [N-(-C(O)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
      [N-(-C(0)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Orn(formamidino)-
             OCH<sub>3</sub>
      [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
20
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(C<sub>2</sub>H<sub>5</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(n-C<sub>3</sub>H<sub>7</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(i-C<sub>3</sub>H<sub>7</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
      [N-(-C(O)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Orn(formamidino)-OCH_3
      [N-(-C(O)(CH_2)_2\overline{Ph})-N-(OH)]Gly-Orn(formamidino)-OCH_3
25
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Orn(formamidino)-OCH_3
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(N(CH<sub>3</sub>)<sub>2</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Orn(formamidino)-OCH<sub>3</sub>
30
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>H)]Gly-Orn(formamidino)-OCH<sub>3</sub>
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
       [N-(-C(O)CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
       [N-(-C(O)(CH<sub>2</sub>)<sub>3</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
       [N-(-C(O)Ph)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
35
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-CH<sub>2</sub>Ph)]Gly-Orn(formamidino)-OCH<sub>3</sub>
         [N-(-C(0)(CH_2)_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(formamidino)-OCH_3
         [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-C1)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
         [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-CH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
   5
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-OCH<sub>3</sub>
         [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
        [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>)<sub>2</sub>Ph]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
        [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-Ph]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
        [N-(-C(0)N(CH_3)CH_2Ph)-N-Ph]Gly-Orn(formamidino)-CO_2CH_3
 10
        [N-(-C(0)Ph-3-CH=CHPh)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
        [N-(-C(0)Ph-3-CH_2CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
        [N-(-C(0)Ph-3-SPh-3-OCH_3)-N-(CH_3)]Gly-Orn(formamidino)-
               CO<sub>2</sub>CH<sub>3</sub>
        [N-(-C(0)Ph-2-CH_2Ph-2-Ph)-N-(CH_3)]Gly-Orn(formamidino)-
 15
               CO2CH3
        [N-(-C(O)Ph-4-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
        [N-(-C(O)Ph-2-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
        [N-(-C(O)Ph-3-CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
        [N-(-C(O)Ph-3-CH_2Ph-2-CF_3)-N-(CH_3)]Gly-Orn(formamidino)-
 20
              CO<sub>2</sub>CH<sub>3</sub>
        [N-(-C(0)CH_2Ph-3,4-Cl_2)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
       [N-(-C(O)(CH_2)_2Ph)-N-(C_2H_5)]Gly-Orn(formamidino)-CO_2CH_3
       [N-(-C(O)(CH_2)_2Ph)-N-(n-C_3H_7)]Gly-Orn(formamidino)-CO_2CH_3
       [N-(-C(0)(CH_2)_2Ph)-N-(i-C_3H_7)]Gly-Orn(formamidino)-CO_2CH_3
       [N-(-C(O)(CH_2)_2Ph)-N-(C_6H_{12})]Gly-Orn(formamidino)-CO_2CH_3
25
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OH)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(OCH<sub>3</sub>)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph)-N-(OCH_2Ph)]Gly-Orn(formamidino)-CO_2CH_3
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NH<sub>2</sub>)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph)-N-(N(CH_3)_2)]Gly-Orn(formamidino)-CO_2CH_3
30
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(NHBoc)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CO<sub>2</sub>H)]Gly-Orn(formamidino)-
              CO<sub>2</sub>CH<sub>3</sub>
       [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CO_2CH_3)]Gly-Orn(formamidino)-
35
             CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(O)CH_2Ph)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
```

```
[N-(-C(O)(CH_2)_3Ph)-N-(CH_3)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(O)Ph)-N-(CH_3)]Gly-Orn(formamidino)-CO_2CH_3
     [N-(-C(0)(CH_2)_2Ph)-N-CH_2Ph)]Gly-Orn(formamidino)-CO_2CH_3
     [N-(-C(0)(CH_2)_2Ph-3, 4-Cl_2)-N-(CH_3)]Gly-Orn(formamidino)-
5
            CO<sub>2</sub>CH<sub>3</sub>
     [N-(-C(0)(CH_2)_2Ph-4-C1)-N-(CH_3)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
     [N-(-C(O)(CH_2)_2Ph-4-CH_3)-N-(CH_3)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-
            CO<sub>2</sub>CH<sub>3</sub>
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph)]-Gly-
10
            boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-CH_3)]-Gly-
            boroLys-OH
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_2C(ethanediy1)Ph-3-CH_3)]-Gly-
15
            boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph-3-CH_3)]-
            Gly-boroLys-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(butane-1,4-diyl)Ph-3-CH<sub>3</sub>)]-
            Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3,5-(CH_3)_2)]-Gly-
20
            boroLys-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CH(i-C<sub>3</sub>H<sub>7</sub>)Ph-3,5-(CH<sub>3</sub>)<sub>2</sub>)]-Gly
            boroLys-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediy1)Ph-3,5-(CH<sub>3</sub>)<sub>2</sub>)]-
25
            Gly-boroLys-OH
      [N-(-C(O)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)_Ph-3,5-
             (CH3)2)]-Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3,5-(CH_3)]
30
             2)]-Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH_2Ph-3-NH_2)]-Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-NH_2)]-Gly-boroLys-
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-NH_2)]-Gly-
35
            boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-NH_2)]-Gly-
```

```
boroLys-OH
       [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediyl)Ph-3-NH<sub>2</sub>)]-Gly-
            boroLys-OH
       [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diy1)Ph-3-NH_2)]_
            Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3-NH_2)]-
            Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH_2Ph-3-NO_2)]-Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-NO_2)]-Gly-
 10
            boroLys-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediyl)Ph-3-NO<sub>2</sub>)]-Gly
            -boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diy1)Ph-3-NO_2)]-
            Gly-boroLys-OH
15
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3-NO_2)]-
            Gly-boroLys-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH_2Ph)]-Gly-boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-boroArg-OH
20
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph)]-Gly-
            boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH_2Ph-3-CH_3)]-Gly-boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-CH_3)]-Gly-boroArg-
           OH
25
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-CH_3)]-Gly-
           boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-CH_3)]-Gly-
           boroArg-OH
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediyl)Ph-3-CH<sub>3</sub>)]-Gly-
30
           boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph-3-CH_3)]-
           Gly-boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3-CH_3)]-
           Gly-boroArg-OH
35
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3,5-(CH_3)_2)]-Gly-
```

PCT/US94/11280

```
boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3,5-(CH_3)_2)]-Gly-
          boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(ethanediyl)Ph-3,5-(CH_3)_2)]-
          Gly-boroArg-OH
5
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,4-diyl)Ph-3,5-
           (CH3)2)]-Gly-boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3,5-(CH_3)]
           2)]-Gly-boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH_2Ph-3-NH_2)]-Gly-boroArg-OH
10
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-NH_2)]-Gly-boroArg-
           OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-NH_2)]-Gly-
           boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-NH_2)]-Gly-
15
           boroArg-OH
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediyl)Ph-3-NH<sub>2</sub>)]-Gly-
           boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph-3-NH_2)]-
20
           Gly-boroArg-OH
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(butane-1, 4-diyl)Ph-3-NH<sub>2</sub>)]-
           Gly-boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH_2Ph-3-NO_2)]-Gly-boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-NO_2)]-Gly-boroArg
25
           -OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-NO_2)]-Gly-
           boroArg-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-NO_2)]-Gly-
           boroArg-OH
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediy1)Ph-3-NO<sub>2</sub>)]-Gly-
30
           boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph-3-NO_2)]-
           Gly-boroArg-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3-NO_2)]-
35
           Gly-boroArg-OH
```

```
[N-(-C(O)(CH2)2Ph)-N-(CH2C(CH2CH3)2Ph)]-Gly-boroOrn
             (formamidino) -OH
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CH(i-C<sub>3</sub>H<sub>7</sub>)Ph)]-Gly-boroOrn
             (formamidino) - OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph)]-Gly-
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CH<sub>2</sub>Ph-3-CH<sub>3</sub>)]-Gly-boroOrn
             (formamidino) -OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-CH_3)]-Gly-
10
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediyl)Ph-3-CH<sub>3</sub>)]-Gly-
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,4-diyl)Ph-3-CH_3)]-
            Gly-boroOrn(formamidino)-OH
15
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3-CH_3)]-
            Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>CH<sub>2</sub>Ph-3,5-(CH<sub>3</sub>)<sub>2</sub>)]-Gly-
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3,5-(CH_3)_2)]-Gly-
20
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3,5-(CH_3)_2)]-Gly-
            boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3,5-(CH_3)_2)]-Gly-
            boroOrn(formamidino)-OH
25
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(ethanediy1)Ph-3,5-(CH_3)_2)]-
            Gly-boroOrn (formamidino) -OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph-3,5-
            (CH3)2)]-Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(butane-1, 4-diy1)Ph-3, 5-(CH<sub>3</sub>)]
30
            2)]-Gly-boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-NH_2)]-Gly-
           boroOrn(formamidino)-OH
      [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-NH_2)]-Gly-
           boroOrn(formamidino)-OH
```

```
[N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(ethanediy1)Ph-3-NH_2)]-Gly-
          boroOrn(formamidino)-OH
    [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diyl)Ph-3-NH_2)]-
          Gly-boroOrn(formamidino)-OH
    [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3-NH_2)]-
5
          Gly-boroOrn(formamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH_2Ph-3-NO_2)]-Gly-boroOrn
          (formamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph-3-NO_2)]-Gly-
          boroOrn(formamidino)-OH
10
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(CH_2CH_3)_2Ph-3-NO_2)]-Gly-
          boroOrn(formamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2CH(i-C_3H_7)Ph-3-NO_2)]-Gly-
          boroOrn(formamidino)-OH
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>2</sub>C(ethanediyl)Ph-3-NO<sub>2</sub>)]-Gly-
15
          boroOrn(formamidino)-OH
     [N-(-C(0)(CH_2)_2Ph)-N-(CH_2C(propane-1,3-diy1)Ph-3-NO_2)]-
          Gly-boroOrn(formamidino)-OH
     [N-(-C(O)(CH_2)_2Ph)-N-(CH_2C(butane-1,4-diyl)Ph-3-NO_2)]-
          Gly-boroOrn(formamidino)-OH
20
     [N-(-C(O)(CH_2)_2-2-pyridyl)-N-(CH_2CH_2Ph)]-Gly-boroLys-OH
     [N-(-C(0)(CH_2)_2-2-pyridyl)-N-(CH_2C(CH_3)_2Ph)]-Gly-
          boroLys-OH
     [N-(-C(0)(CH_2)_2-2-pyridy1)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-
25
          boroLys-OH
     [N-(-C(O)(CH_2)_2-2-pyridyl)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-
           boroLys-OH
     [N-(-C(O)(CH_2)_2-2-pyridy1)-N-(CH_2C(ethanediy1)Ph)]-Gly-
           boroLys-OH
     [N-(-C(0)(CH_2)_2-2-pyridy1)-N-(CH_2C(propane-1,3-diy1)Ph)]
30
           1-Gly-boroLys-OH
     [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>-2-pyridyl)-N-(CH<sub>2</sub>C(butane-1,4-diyl)Ph)]
           -Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2-3-pyridyl)-N-(CH_2CH_2Ph)]-Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2-3-pyridy1)-N-(CH_2C(CH_3)_2Ph)]-Gly-
35
```

```
boroLys-OH
        [N-(-C(O)(CH_2)_2-3-pyridy1)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-
             boroLys-OH
       [N-(-C(0)(CH_2)_2-3-pyridy1)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-
   5
             boroLys-OH
       [N-(-C(O)(CH_2)_2-3-pyridyl)-N-(CH_2C(ethanediyl)Ph)]-Gly-
             boroLys-OH
       [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>-3-pyridyl)-N-(CH<sub>2</sub>C(propane-1,3-diyl)Ph)
             ]-Gly-boroLys-OH
       [N-(-C(O)(CH_2)_2-3-pyridyl)-N-(CH_2C(butane-1,4-diyl)Ph)]
  10
             -Gly-boroLys-OH
       [N-(-C(O)(CH_2)_2Ph-3-OH)-N-(CH_2CH_2Ph)]-Gly-boroLys-OH
       [N-(-C(O)(CH_2)_2Ph-3-OH)-N-(CH_2C(CH_3)_2Ph)]-Gly-boroLys-OH
      [N-(-C(O)(CH_2)_2Ph-3-OH)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-boroLys-OH
 15
      [N-(-C(O)(CH_2)_2Ph-3-OH)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-boroLys-OH
      [N-(-C(O)(CH_2)_2Ph-3-OH)-N-(CH_2C(ethanediyl)Ph)]-Gly-
            boroLys-OH
      [N-(-C(O)(CH_2)_2Ph-3-OH)-N-(CH_2C(propane-1,3-diyl)Ph)]-Gly-
 20
            boroLys-OH
      [N-(-C(O)(CH_2)_2Ph-3-OH)-N-(CH_2C(butane-1,4-diyl)Ph)]-Gly-
           boroLys-OH
      [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-3-OMe)-N-(CH<sub>2</sub>CH<sub>2</sub>Ph)]-Gly-boroLys-OH
25
       [N-(-C(O)(CH_2)_2Ph-3-OMe)-N-(CH_2C(CH_3)_2Ph)]-Gly-boroLys-OH 
      [N-(-C(O)(CH_2)_2Ph-3-OMe)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-boroLys-
           OH
      [N-(-C(O)(CH_2)_2Ph-3-OMe)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-boroLys-
30
     [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-3-OMe)-N-(CH<sub>2</sub>C(ethanediyl)Ph)]-Gly-
           boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-OMe)-N-(CH_2C(propane-1,3-diyl)Ph)]-Gly-
          boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-OMe)-N-(CH_2C(butane-1,4-diyl)Ph)]-Gly-
35
          boroLys-OH
```

```
[N-(-C(0)(CH_2)_2Ph-3-NO_2)-N-(CH_2CH_2Ph)]-Gly-borolys-OH
     [N-(-C(O)(CH_2)_2Ph-3-NO_2)-N-(CH_2C(CH_3)_2Ph)]-Gly-boroLys-OH
     [N-(-C(0)(CH_2)_2Ph-3-NO_2)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-boroLys-
5
     [N-(-C(0)(CH_2)_2Ph-3-NO_2)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-boroLys-
          OH
     [N-(-C(0)(CH_2)_2Ph-3-NO_2)-N-(CH_2C(ethanediyl)Ph)]-Gly-
          boroLys-OH
     [N-(-C(0)(CH_2)_2Ph-3-NO_2)-N-(CH_2C(propane-1,3-diyl)Ph)]-Gly-
10
          boroLys-OH
     [N-(-C(0)(CH_2)_2Ph-3-NO_2)-N-(CH_2C(butane-1,4-diyl)Ph)]-Gly-
          boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-CO_2H)-N-(CH_2CH_2Ph)]-Gly-boroLys-OH
     [N-(-C(0)(CH_2)_2Ph-3-CO_2H)-N-(CH_2C(CH_3)_2Ph)]-Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-CO_2H)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-boroLys-CH_2C(CH_2CH_3)_2Ph)
     [N-(-C(0)(CH_2)_2Ph-3-CO_2H)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-boroLys-
20
     [N-(-C(O)(CH_2)_2Ph-3-CO_2H)-N-(CH_2C(ethanediyl)Ph)]-Gly-
           boroLys-OH
     [N-(-C(0)(CH_2)_2Ph-3-CO_2H)-N-(CH_2C(propane-1,3-diyl)Ph)]-
           Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-CO_2H)-N-(CH_2C(butane-1,4-diyl)Ph)]-Gly-
25
           boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-CO_2Me)-N-(CH_2CH_2Ph)]-Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-CO_2Me)-N-(CH_2C(CH_3)_2Ph)]-Gly-boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-CO_2Me)-N-(CH_2C(CH_2CH_3)_2Ph)]-Gly-
30
           boroLys-OH
     [N-(-C(O)(CH_2)_2Ph-3-CO_2Me)-N-(CH_2CH(i-C_3H_7)Ph)]-Gly-
           boroLys-OH
      [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-CO<sub>2</sub>Me)-N-(CH<sub>2</sub>C(ethanediyl)Ph)]-Gly-
35
           boroLys-OH
```

```
[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-CO<sub>2</sub>Me)-N-(CH<sub>2</sub>C(propane-1,3-diyl)Ph)]-Gly-boroLys-OH
```

- [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-CO<sub>2</sub>Me)-N-(CH<sub>2</sub>C(butane-1, 4-diyl)Ph)]-Gly-boroLys-OH
- 5 [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-NH<sub>2</sub>)-N-(CH<sub>2</sub>CH<sub>2</sub>Ph)]-Gly-boroLys-OH
  - [N-(-C(0)(CH2)2Ph-3-NH2)-N-(CH2C(CH3)2Ph)]-Gly-boroLys-OH
  - [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-NH<sub>2</sub>)-N-(CH<sub>2</sub>C(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>Ph)]-Gly-boroLys-OH
  - [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-NH<sub>2</sub>)-N-(CH<sub>2</sub>CH(*i*-C<sub>3</sub>H<sub>7</sub>)Ph)]-Gly-boroLys-
- $[N-(-C(0)(CH_2)_2Ph-3-NH_2)-N-(CH_2C(ethanediyl)Ph)]-Gly-boroLys-OH$ 
  - [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-NH<sub>2</sub>)-N-(CH<sub>2</sub>C(propane-1,4-diyl)Ph)]-Gly-boroLys-OH
- 15 [N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph-3-NH<sub>2</sub>)-N-(CH<sub>2</sub>C(butane-1,4-diyl)Ph)]-Gly-boroLys-OH

This invention also provides compositions comprising one or more of the foregoing compounds and methods of using such compositions in the treatment of aberrant proteolysis such as thrombosis in mammals.

## Detail Description of the Invention

- As used throughout the specifications, the following abbreviations for amino acid residues or amino acids apply:
  - Ala = L-alanine
  - Arg = L-arginine
- 30 Asn = L-asparagine
  - Asp = L-aspartic acid
  - Cys = L-cysteine
  - Gln = L-glutamine
  - Glu = L-glutamic acid
- 35 Gly = glycine

His = L-histidine

Ile = L-isoleucine

Leu = L-leucine

Lys = L-lysine

5 Met = L-methionine

Phe = L-phenylalanine

Pro = L-proline

Ser = L-serine

Thr = L-threonine

10 Trp = L-tryptophan

Tyr = L-tyrosine

Val = L-valine

Sar = L-sarcosine

Irg = L-arginine where the guanidine is replaced

with an isothiouronium (-SC(=NH)NH<sub>2</sub>)

The "D" prefix for the foregoing abbreviations indicates the amino acid is in the D-configuration. indicates the amino is present in mixture of the D- and the L-configuration. The prefix "boro" indicates amino acid 20 residues where the carboxyl is replaced by a boronic acid or a boronic acid ester. For example, if R1 is isopropyl and  $Y^1$  and  $Y^2$  are OH, the C-terminal residue is abbreviated "boroVal-OH" where "-OH" indicates the boronic acid is in the form of the free acid. The pinanediol boronic acid 25 ester and the pinacol boronic acid ester are abbreviated "- $C_{10}H_{16}$ " and "- $C_{6}H_{12}$ ", respectively. Examples of other useful diols for esterification with the boronic acids are 1,2-ethanediol, 1,3-propanediol, 1,2-propanediol, 2,3butanediol, 1,2-diisopropylethanediol, 5,6-decanediol, and 1,2-dicyclohexylethanediol. Other abbreviations are: formamidino, HC(=NH)-; N-methylamidino, CH<sub>3</sub>NHC(=NH)-; Z, benzyloxycarbonyl; BSA, benzene sulfonic acid; THF, tetrahydrofuran; Boc-, t-butoxycarbonyl-; Ac-, acetyl; pNA, p-nitro-aniline; DMAP, 4-N, N-dimethylaminopyridine; Tris, 35

Tris(hydroxymethyl)aminomethane; MS, mass spectrometry; FAB/MS, fast atom bombardment mass spectrometry. LRMS(NH3-CI) and HRMS(NH3-CI) are low and high resolution mass spectrometry, respectively, using NH3 as an ion source. Thus, an example of the chemical structure based on the nomenclature used herein is:

[N-(-C(0)(CH<sub>2</sub>)<sub>2</sub>Ph)-N-(CH<sub>3</sub>)]Gly-Orn(formamidino)-CO<sub>2</sub>CH<sub>3</sub>represents

10

and [N-(-C(O)(CH<sub>2</sub>)<sub>2</sub>Ph-4-OCH<sub>3</sub>)-N-(CH<sub>3</sub>)]Gly-boroPhe(mCN)-OHrepresents

15

It is understood that many of the compounds of the present invention contain one or more chiral centers and that these stereoisomers may possess distinct physical and biological properties. The present invention comprises all of the stereoisomers or mixtures thereof. If the pure 20 enantiomers or diasteromers are desired, they may be prepared using starting materials with the appropriate stereochemistry, or may be separated from mixtures of undesired stereoisomers by standard techniques, including

chiral chromatography and recrystalization of diastereomeric salts.

The term "amine-blocking group" or "amine-protecting group" as used herein, refers to various acyl, thioacyl, alkyl, sulfonyl, phosphoryl, and phosphinyl groups comprised of 1 to 20 carbon atoms. Substituents on these groups maybe either alkyl, aryl, alkaryl which may contain the heteroatoms, O, S, and N as a substituent or as inchain component. A number of amine-blocking groups are recognized by those skilled in the art of organic 10 synthesis. Examples of suitable groups include formyl, acetyl, benzoyl, trifluoroacetyl, and methoxysuccinyl; aromatic urethane protecting groups, such as, benzyloxycarbonyl; and aliphatic urethane protecting groups, such as t-butoxycarbonyl or adamantyloxycarbonyl. 15 Gross and Meienhofer, eds., The Peptides, Vol 3; 3-88 (1981), Academic Press, New York, and Greene and Wuts Protective Groups in Organic Synthesis, 315-405 (1991), J. Wiley and Sons, Inc., New York disclose numerous suitable 20 amine protecting groups and they are incorporated herein by reference for that purpose.

"Amino acid residues" as used herein, refers to natural or unnatural amino acids of either D- or L-configuration. Natural amino acids residues are Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val. Roberts and Vellaccio, The Peptides, Vol 5; 341-449 (1983), Academic Press, New York, discloses numerous suitable unnatural amino acids and is incorporated herein by reference for that purpose.

"Amino acids residues" also refers to various amino acids where sidechain functional groups are coupled with appropriate protecting groups known to those skilled in the art. "The Peptides", Vol 3, 3-88 (1981) discloses numerous suitable protecting groups and is incorporated herein by reference for that purpose.

30

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms.

"Alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge.

"Cycloalkyl" is intended to include saturated ring groups, including mono-, bi- or poly-cyclic ring systems, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl and cyclooctyl, and so forth.

"Alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl, propenyl, and the like.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo.

The term "aryl" is defined as phenyl, fluorenyl, biphenyl and naphthyl, which may be unsubstituted or include optional substitution with one to three substituents.

The term "heteroaryl" is meant to include 5-, 6-, 9-, or 10-membered mono- or bicyclic aromatic rings which can optionally contain from 1 to 3 heteroatoms selected from the group consisting of O, N, and S; said ring(s) may be unsubstituted or include optional substitution with one to three substituents. Included in the definition of the group heteroaryl, but not limited to, are the following: 2-, or 3-, or 4-pyridyl; 2-or 3-furyl; 2- or 3-benzofuranyl; 2-, or 3-thiophenyl; 2- or 3-benzo[b]thiophenyl; 2-, or 3-, or 4-quinolinyl; 1-, or 3-, or 4-isoquinolinyl; 2- or 3-pyrrolyl; 1- or 2- or 3- indolyl; 2-, or 4-, or 5-oxazolyl; 2-benzoxazolyl; 2- or 4- or 5-imidazolyl; 1- or 2-benzimidazolyl; 2- or 4- or 5-thiazolyl; 2-benzothiazolyl; 3- or 4- or 5-isothiazolyl; 3- or 4- or 5-is

5

20

25

30

pyrimidinyl; 2-pyrazinyl; 2-triazinyl; 3- or 4- cinnolinyl;
1-phthalazinyl; 2- or 4-quinazolinyl; or 2-quinoxalinyl
ring. Particularly preferred are 2-, 3-, or 4-pyridyl; 2-,
or 3-furyl; 2-, or 3-thiophenyl; 2-, 3-, or 4-quinolinyl;
5 or 1-, 3-, or
4-isoquinolinyl.

The term "heterocycle" is meant to include 5-, 6-, 9-, or 10-membered mono- or bicyclic rings which can optionally contain from 1 to 3 heteroatoms selected from the group consisting of N, O, or S, with the proviso that proline is excluded from this group; said ring(s) may be unsubstituted or include optional substitution with one to three substituents. Included in the definition of the group heterocycle, but not limited to, are tetrahydroisoquinoline, tetrahydroquinoline, tetrahydroquinoline, piperazine, morpholine. Particularly preferred are 1-, 3-, or 4-tetrahdroisoquinolinyl.

Unless otherwise specified, the substituents that may be attached to the aryl, heteroaryl or heterocycle ring(s) may be independently selected at each occurrence from the group consisting of:

halogen,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy, methylenedioxy, -NO2, -CF3, -SH, -S(0)<sub>r</sub>-( $C_1$ - $C_4$  alkyl), CN, -OH, -NH2, -NH( $C_1$ - $C_4$  alkyl), -N( $C_1$ - $C_4$  alkyl)2, -C(=NH)NHR<sup>4</sup>, -NHC(=NR<sup>4</sup>), -NHC(=NH)NHR<sup>4</sup>, -NHC(=O)R<sup>4</sup>, -(CH<sub>2</sub>)<sub>p</sub>-CO<sub>2</sub>R<sup>4</sup>, -NHCO C<sub>1</sub>-C<sub>4</sub> alkoxy), -NH( $C_1$ - $C_4$  alkoxy)2, -N( $C_1$ - $C_4$  alkoxy), phenyl which may be unsubstituted or substituted with  $R^{13}$ .

By "stable compound" or "stable structure" is meant herein a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

10

15

20

25

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound of formula (I) is modified by making acid or base salts of the compound of formula (I). Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

"Prodrugs" are considered to be any covalently bonded 10 carriers which release the active parent drug according to formula (I) in vivo when such prodrug is administered to a mammalian subject. Prodrugs of the compounds of formula (I) are prepared by modifying functional groups present in the compounds in such a way that the modifications are 15 cleaved, either in routine manipulation or in vivo, to the parent compounds. Prodrugs include compounds of formula (I) wherein hydroxy, amine, or sulfhydryl groups are bonded to any group that, when administered to a mammalian subject, cleaves to form a free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formula (I); and the like.

Pharmaceutically acceptable salts of the compounds of the invention can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

35

25

## Synthesis '

The compounds of Formula (I) can be prepared using the reactions and techniques described below, in addition to synthetic procedures described in Applicant's Assignee's commonly assigned patent applications USSN 08/010,731 (filed January 29, 1993), USSN 08/036,378 (filed March 24, 1993), and USSN 08/052,835 (filed April 27, 1993), all of which are hereby incorporated by reference. The reactions are performed in a solvent appropriate to the reagents and 10 materials employed and suitable for the transformations being effected. It will be understood by those skilled in the art of organic synthesis that the functionality present on the molecule should be consistent with the transformations proposed. This will sometimes require a 15 judgment to modify the order of the synthetic steps or to select one particular process scheme over another in order to obtain a desired compound of the invention. It will also be recognized that another major consideration in the planning of any synthetic route in this field is the 20 judicious choice of the protecting group used for protection of the reactive functional groups present in the compounds described in this invention. An authoritative account describing the many alternatives to the trained practitioner is Greene and Wuts, Protective Groups In 25 Organic Synthesis, Wiley and Sons (1991).

## Scheme 1

M is a carboxylic acid protecting group

5 The preparation of the required N, N-disubstituted amino acid subunit can be accomplished by the sequence outlined in Scheme 1. A variety of methods for the asymmetric synthesis of the amino acids required for the amino acid ester substrate (II) is reviewed by Morrison and 10 Mosher (Asymmetric Organic Reactions, American Chemical Society, 297-334 (1976) and references there in). appropriate ester (II), where M is the ester residue, can be conjugated to give the N-substituted intermediate (III) by N-monoalkylation with an alkyl halide related to  $\mathbb{R}^{11}$ . 15 Typical conditions for N-monoalkylation include the admixture of an excess of (II), the required alkyl bromide or iodide and a base in an anhydrous polar aprotic solvent, such as acetone, acetonitrile, N, N-dimethylformamide or methyl sulfoxide. The option exists for stirring this mixture at room temperature or heating at temperatures up 20 to the reflux point of the selected solvent. The base added is chosen so that it will not interfere with the ester functionality of (II); among those recommended are non-nucleophilic bases such as sodium hydride or potassium 25 carbonate. Another general route for the preparation of compounds of this type is the reductive amination of (II) with a selected aldehyde related to R11. In this procedure, a mixture of (II) and the aldehyde are heated in an anhydrous non polar solvent, such as benzene, toluene or xylene, with continuous removal of evolved water by drying 30

agents or azeotropic distillation. This process causes condensation of the aldehyde with amine (II). The condensation product is reduced to monoalkylated (III) by treatment with a selective hydride reducing agent such as sodium cyanoborohydride or sodium borohydride, according to the method of Getson et. al., J. Heterocycl. Chem. 1, 300 (1964), or by catalytic hydrogenation with platinum, palladium, nickel or Raney nickel in an alcohol solvent like propanol, ethanol or methanol, according to the method of Hudlicky, Reductions In Organic Synthesis, John Wiley and Sons, pp. 134 (1984).

While a number of coupling or acylation methods can be contemplated for the preparation of the disubstituted derivative (IV) from (III), (see Bodanszky and Bodanszky, The Practice of Peptide Synthesis, Springer-Verlag, p. 87-15 150 (1984)), three methods are preferred. In the first, a solution of (III) in an anhydrous non polar solvent, such as tetrahydrofuran or dichloromethane, at -78°C or higher is treated sequentially with a selected acid chloride related to  $\mathbb{R}^3$  followed by a trialkylamine base. This 20 mixture is allowed to warm to ambient temperature over several hours if required. The second method is the mixed anhydride procedure of Anderson et al. reported in J. Am. Chem. Soc. 89, 5012 (1967). In this procedure the alkyl mixed anhydride is generated by dissolving a selected 25 carboxylic acid related to R3 in non-polar anhydrous solvent, such as tetrahydrofuran or dichloromethane, and adding one equivalent of a trialkylamine base. solution is stirred at .- 78°C or higher and one equivalent of an alkylchloroformate is added. After formation of the mixed anhydride is complete, a solution of one equivalent each of intermediate (III) and a trialkylamine base is added dropwise. The mixture is stirred with or without cooling until the reaction is complete. The third method preferred for amide formation is the hydroxybenzotriazole / 35

dialkylcarbodiimide method of Koing and Geiger in Chem. Ber. 103, 788 (1970). Thus, to (III) and a selected carboxylic acid related to R<sup>3</sup>, dissolved in an aprotic solvent like N,N-dimethylformamide, dichloromethane or tetrahydrofuran, at -78°C or higher, is added dialkylcarbodiimide, hydroxybenzotriazole and a trialkylamine base. If necessary, the stirred solution is allowed to thaw to ambient temperature over several hours.

An alternative preparation of N, N-disubstituted amino acids uses  $\alpha$ -halo- or  $\alpha$ -sulfonate acylesters such as (V) of Scheme 2. Compound (V) can be treated with a primary amine related to  $R^{11}$  in the presence of a variety of bases like potassium carbonate, triethyl amine or sodium hydride and in solvents such as ethyl ether, acetone or

dimethylformamide at temperatures ranging from -78° C to the reflux point of the solvent selected. From this reaction can be isolated the N-alkyl aminoacid ester (III) of Scheme 1; the N-alkyl-N-acylamino acid ester (IV) can be prepared from compound (III) by any of the methods outlined in Scheme 1 and the related discussion hereafter.

The preparation of intermediates which will lead to compounds where  $\mathbb{R}^3$  and  $\mathbb{R}^{11}$  may be taken together to form a cyclic amide or phthalimide is described in Scheme 2. N,N-Disubstituted  $\alpha$ -amino acid subunits (VI) which lead to compounds of this type are best derived by reaction of an appropriate  $\alpha$ -haloester (V), where J = Cl, Br, I.

10

### Scheme 2

$$R^4$$
  $R^5$   $OM$   $L = \begin{cases} -N - R^{11} \\ R^3 \end{cases}$   $J = \text{Leaving group}$   $M = \text{carboxylic acid protecting group}$ 

with the alkali metal salt of an amide or phthalimide related to the desired cyclized combination of  ${\rm R}^3$  and  ${\rm R}^{11}$ in a polar aprotic solvent according to the method of Daly et. al.in J. Med. Chem. 33, 2818 (1990); Neuberger and In a typical Scott, J. Chem. Soc. p 1820 (1954). preparation the alkali metal salt of the required cyclic amide or phthalimide is generated by adding one equivalent of a strong non-nucleophilic base such as sodium or potassium hydride, a lithium dialkylamine, or lithium trimethyl- or lithium triphenylmethane to a solution of the amide or phthalimide in an anhydrous inert solvent, such as tetrahydrofuran or 1,2-dimethoxyethane, at -78° C or higher. When the salt formation is complete, the solvent is removed by distillation and replaced by the appropriate polar aprotic solvent such as acetonitrile, N, Ndimethylformamide or methylsulfoxide. The appropriate  $\alpha$ chloro- or  $\alpha$ -bromoester is introduced and the mixture stirred at room temperature or with heating until the haloester is consumed.

It will be recognized by those skilled in the art of organic synthesis that the acid derivatives of the N,N-disubstituted  $\alpha$ -amino acid esters (IV) and (VI) are the required precursors for the preparation of the thrombin inhibitors disclosed in this invention. It is recommended that compounds (IV) and (VI) be prepared as either the benzyl, methyl or t-butyl esters because of the ease with which esters of these types may be converted to their

10

15

20

25

corresponding acids. In the case of a benzyl ester (e.g., (IV) or (VI), where  $M = -CH_2C_6H_5$ ), hydrogenolysis of an alcohol solution of the compound may be effected under an atmosphere of hydrogen gas in the presence of platinum or palladium on carbon catalyst according to the reported by Hartney and Simonoff, Org. React. VII, 263 (1953); with a methyl ester (IV) or (VI), where  $M = -CH_3$ , treatment of an ethanol solution of the compound with an aqueous base, such as one equivalent of sodium hydroxide solution, will give 10 the desired acid. The t-butyl ester (IV) or (VI), where M = -C(CH<sub>3</sub>)<sub>3</sub>, is readily cleaved by acid under anhydrous conditions; for example trifluoroacetic acid in dichloromethane solution removes the t-butyl ester of derivatives of (IV) at ambient temperature as reported by 15 Bryan et. al., J. Am. Chem. Soc. 99, 2353 (1977). A number of alternative esters and procedures are detailed in Greene and Wuts (1991).

Scheme 3 illustrates the coupling of the acid derivatives of (IV) or (VI) with boropeptide synthons (VII) or (VIII) to give intermediates (IX) or (X).

### Scheme 3

$$(IV): L = R^{3}R^{11}N$$

$$(VI): L = R^{11} - N$$

$$(VII): y = 3$$

$$(VIII): y = 4$$

$$(X): y = 4$$

25

The preparation of synthons (VII) and (VIII) has been described by Kettner and Shenvi (EP 293 881 A2). It will be recognized by those skilled in the art of organic synthesis that the methodology described by Kettner and Shenvi can be applied to make homologous boropeptide 5 synthons related to (VII) and (VIII). These homologues may be used in the appropriate processes described herein to prepare the corresponding thrombin inhibitors. coupling of the carboxylic acid derivative of (IV) or (VI) to boropeptide synthon (VII) or (VIII) has been described 10 previously by Kettner et al. in J. Biol. Chem. 265, 18289 (1990) and, in general, the standard amino acid coupling protocols detailed by Bodanszky and Bodanszky (1984) are effective for making the compounds of this invention. Preferred methods are the mixed anhydride procedure of 15 Anderson et al. (1967) and the hydroxybenzotriazole/dialkylcarbodiimide method of Koing and Geiger (1970). In the mixed anhydride procedure, the anhydride is generated by dissolving a carboxylic acid related to (IV) or (VI) in a non polar anhydrous solvent, 20 such as tetrahydrofuran or dichloromethane, and adding one equivalent of a trialkylamine base. The solution is stirred at -78° and up to 0°C, then one equivalent of an alkylchloroformate is added. After formation of the mixed anhydride is complete, a solution of boropeptide synthon 25 (VII) or (VIII) and a trialkylamine base is added. The mixture is stirred for one hour with cooling followed by several hours at ambient temperature. By the hydroxybenzotriazole/dialkylcarbodiimide method, (VII) or (VIII) and the acids of (IV) or (VI) are dissolved in an 30 aprotic solvent, such as N, N-dimethylformamide, dichloromethane or tetrahydrofuran, at -78° or higher. To this solution one equivalent each of dialkylcarbodiimide, hydroxybenzotriazole and a trialkylamine base are added.

If necessary, the solution is allowed to stir and thaw to ambient temperature over several hours.

A process for the preparation of the boropeptide thrombin inhibitors of this invention from intermediates (IX) and (X) is disclosed in Scheme 4. Compound (IX) serves as a starting point for isothiouronium thrombin inhibitors (XI) and (XII). The boronic ester (XI) is prepared by stirring a solution of (IX) and thiourea in an inert polar solvent, such as an alcohol or N,N-dimethylformamide, at temperatures ranging from ambient to the reflux temperature of the selected solvent. It is understood that a boronic acid ester like compound (XI) is an effective thrombin inhibitor, however, it may be transformed to the corresponding free boronic acid (XII) without a loss of biological activity. Compound (XII) is derived from the boron ester (XI) by transesterification

under equilibrium conditions.

10

## Scheme 4

$$(CH_{2})_{y} = 3$$

$$(CH_{2})_{y} = 4$$

$$(N) : y = 3$$

$$(XIII) : y = 3$$

$$(XIV) : y = 4$$

$$(XIV) : y = 4$$

$$(XVI)$$

$$(CH_{2})_{y} = 4$$

$$(XVI)$$

$$(XIV) : y = 3$$

$$(XIV) : y = 4$$

$$(XVI)$$

$$(XVI) : y = 3$$

$$(XVIII)$$

 $(XII): A = B(OH)_2$ 

Thus stirring ester (XI) with an excess of an alkyl- or aryl boric acid in a biphasic mixture of neutral or acidic water and an immiscible solvent, such as ethyl ether or toluene, gives (XII) after several hours at ambient temperature. The conditions generally preferred use 5 to 10 equivalents of phenylboric acid in ethyl ether/water at neutral pH. Thrombin inhibitors (XIII) to (XVI) are obtained by reduction of an azide intermediate prepared from (IX) or (X). The azide intermediate is prepared by heating either (IX) or (X) with an inorganic azide, such as sodium or potassium azide, in an anhydrous polar aprotic solvent, such as acetone, dimethylformamide or methyl

10

sulfoxide at temperatures ranging from ambient to 130°C. Alternatively, phase transfer conditions may be employed to prepare the azide intermediate from (IX) or (X). For example, a tetraalkylammonium azide in a non-polar aprotic solvent, such as tetrahydrofuran or toluene, or a crown ether and inorganic azide in biphasic mixtures of water and an immiscible solvent, such as benzene, toluene or xylene, can be stirred at room temperature or heated up to the reflux point of the selected solvent. The primary amines (XIII) and (XIV) are most conveniently obtained from the catalytic hydrogenation of the azide in an inert solvent, such as an alcohol, ethyl acetate or tetrahydrofuran with a transition metal catalyst such as platinum or palladium on carbon under an atmosphere of hydrogen gas. A variety of alternative methods are also useful and can be found in the monograph by Hudlicky (1984, pp. 76). The acid salt of the resulting amines (XIII) and (XIV) may be formed by the addition of one equivalent of the desired acid to the hydrogenation mixture. Phenylboric acid mediated hydrolysis of esters (XIII) and (XIV) gives the free boronic acid thrombin inhibitors (XV) and (XVI), compounds of formula (I) of the invention.

Compounds containing a primary guanidine or N- alkyl guanidine functionality may be prepared by the alternative process outlined in Scheme 4. As illustrated with primary amine (XIII), the transformation to (XVII) is effected with a guanidinylation agent, such as an S-alkyl thiourea, aminoiminomethane sulfonic acid reported by Miller and Bischoff Synthesis 9, 777 (1986), cyanamide reported by Kettner et al. (1990) or their N-alkyl derivatives. This mixture is stirred at room temperature or higher with a base, such as potassium carbonate, triethylamine or N,N-dimethylaminopyridine in an inert solvent like water, alcohol, N,N-dimethylformamide or acetone. The guanidine boronic acid esters (XVII) can be deesterified to give the

10

15

20

25

PCT/US94/11280

WO 95/09634

corresponding boronic acid (XVIII) by the phenylboric acid procedure described above.

According to Scheme 5, the bromide (X) is converted to the corresponding alkylnitrile (XIX) upon exposure to the cyanide anion under a variety of conditions.

5

### Scheme 5

$$(CH_{2})_{4} \qquad NR^{3}R^{11} \qquad (CH_{2})_{4} \qquad NR^{3}R^{11} \qquad (CH_{2})_{4} \qquad NR^{3}R^{11} \qquad (CH_{2})_{4} \qquad NR^{3}R^{11} \qquad (CH_{2})_{5} \qquad NR^{3}R^{11} \qquad (CH_$$

Effective methods include the use of potassium or sodium cyanides in polar aprotic solvents, such as N,N-dimethylformamide, methylsulfoxide, acetone or ethylmethyl ketone, at temperatures ranging from ambient up to the reflux point of the selected solvent. More useful, however, are conditions employing phase transfer agents such as tetrabutylammonium cyanide in a nonpolar aprotic solvent such as tetrahydrofuran or toluene, or a biphasic mixture of a crown ether and an inorganic cyanide in water with an immiscible solvent like benzene, toluene or xylene. These mixtures can be stirred at ambient temperature or heated up

10

to the reflux temperature of the selected solvent. An amidine like (XX) is prepared by first treating nitrile (XIX) with a saturated solution of a mineral acid such as hydrogen chloride in an alcohol solvent at room temperature or lower. The intermediate O-alkylimidate can be exposed to ammonia, or a primary or secondary amine under anhydrous conditions with or without an inert solvent. As illustrated in Scheme 5, compound (XX) is produced by treating the O-alkylimidate formed from (XIX) with neat anhydrous ammonia at reflux. The free boronic acid (XXI) is obtained by transesterification of (XX) with phenylboric acid in a mixture of water and diethyl ether.

The formamidine substituted boronic acid (XXIII) is prepared from alkylamine (XV) as shown in Scheme 6.

15 Compounds of (XV) can be stirred with an O-alkyl or O-aryl formimidate from 0°C or up to the reflux temperature of an inert anhydrous solvent such as tetrahydrofuran or N,N-dimethylformamide to give formamidine (XXII). Free boronic acid (XXIII) is produced from (XXII) by the phenylboric acid transesterification protocol.

5

### Scheme 6

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(XV) \qquad (XXIII)$$

$$(HO)_{2}B \qquad O \qquad NR^{3}R^{11}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(XXIII) \qquad (XXIII)$$

5 As shown in Scheme 7, the N-cyanoguanidine substituted boronic acid (XXVI), can be prepared by the reaction of (XV) with an N-cyanoisourylation agent such as S, S-dimethyl N-cyanoiminodithiocarbonate or O, O-diphenyl N-cyanodiiminocarbonate. In this general process, compounds of Formula 10 (XV) are combined with a selected iminocarbonate in an inert, anhydrous solvent like tetrahydrofuran or N, Ndimethylformamide. The mixture is stirred at 0°C or up to the reflux temperature of the chosen solvent until there is obtained an N-cyano-S-isourea or N-cyano-O-isourea of the 15 Formula (XXIV) similar to that reported by Barpill et al., J. Heterocyclic Chem. 25, 1698 (1988). This intermediate is treated with an amine such as

### Scheme 7

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(XV) \qquad (XXIV) \text{ wherein}$$

$$X \text{ is -SMe, -OPh}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

5 ammonia, or more generally, an alkylamine or an arylamine with or without an inert solvent like water, tetrahydrofuran or an alcohol at temperatures ranging from 0°C to reflux to give the aminolysis product (XXV).

Treatment of (XXV) as described above with phenylboric acid can provide (XXVI).

The N-hydroxyguanidino inhibitors, as shown in Scheme 8, are prepared by treating amine (XV) with cyanogen bromide or cyanogen chloride followed by hydroxylamine in an inert solvent to yield (XXVII) according to Nakahara et. al., Tetrahedron 33, 1591 (1977); and Belzecki et al., J. Chem. Soc. Chem. Commun. p. 806 (1970). Transesterification of (XXVII) by the phenylboric acid method can provide (XXVIII).

#### Scheme 8.

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

$$(XV) \qquad (XXVII)$$

$$(XV) \qquad (XXVII)$$

$$(HO)_{2}B \qquad O \qquad NR^{3}R^{11}$$

$$(CH_{2})_{3} \qquad NR^{3}R^{11}$$

5 A general preparation for the new aromatic boronic acids is illustrated in Scheme 9. Functionalized benzylic anions containing either a halo- or cyano- substituent are obtained with a variety of metalation agents, such as activated zinc metal/CuCN.2LiCl based on the report of Berk 10 et al. Organometallics 9, 3053 (1990); or use of lithium metal according to Michel et al., J. Organometallic Chem. 204, 1 (1981); or lithium naphthalenide in the presence of zinc chloride based on the report of Zhu et al., J. Org. Chem. 56, 1445 (1991) in an inert solvent like 15 tetrahydrofuran or 1,2-dimethoxyethane at temperatures of -78°C or higher. Dichloromethyl boronic acid pinanediol, prepared by the method described by Tsai et al. in Organometallics 2, 1543 (1983), is allowed to react with the transmetallated anion in the selected solvent to give 20 The a-aminoboronic acid, (XXX), can be obtained by treating (XXIX) with the sodium or lithium salt of hexamethyldisilizane in a polar aprotic solvent like

5

acetone, N,N-dimethylformamide or methyl sulfoxide with heating at temperature up to the reflux point of the selected solvent, if necessary. The trimethylsilyl protecting groups are removed by treatment with

Scheme 9

$$\begin{array}{c} CH_2ZnBr \\ CN \\ CN \\ CN \\ CN \\ (XXIX) \\ (XXX) \\ (XXXX) \\ (XXXX) \\ (XXXX) \\ (XXXX) \\ (XXXX) \\ (XXXIa) \\ (XXXIa) \\ (XXXIa) \\ (XXXIb) \\ (XXXIb) \\ (XXXIb) \\ (XXXIb) \\ (XXXIb) \\ (XXXIIb) \\ (XXXIIb$$

anhydrous acid such as gaseous hydrogen chhloride or trifluoroacetic acid in an inert solvent like tetrahydrofuran or dichloromethane at -78°C or higher. Compound (XXX) was coupled to the N,N-disubstituted a-amino acids (IV) or (VI), using the techniques described in Scheme 3, to give the boronic acid ester (XXXIa). Transesterification of (XXXIa) by the phenylboric acid protocol (vide infra) gives inhibitor (XXXIb). In Scheme

9, the aromatic nitrile (XXXIa) is converted to the amidine (XXXIIa) by methods described for the synthesis of aliphatic amidine (XX) in Scheme 5. Removal of the pinanediol protecting group of (XXXIIa) gives the free boronic acid derivative (XXXIIb).

As detailed in Scheme 10, compound (XXXIa) is a versatile intermediate that can be hydrogenated to yield the aminomethyl derivative (XXXIIIa) under a variety of conditions (Hudlicky, (1984), pp 173).

10

# Scheme 10.

15 Catalysts recommended for this transformation include transition metals like rhodium, Raney nickel, nickel

boride, nickel, platinum or palladium; these reductions occur readily under atmospheres of hydrogen or ammonia at pressures ranging from 1 to 300 atmospheres, at room temperature or higher, and in inert solvents such as water, an alcohol, ethyl acetate or tetrahydrofuran. Furthermore, from (XXXIIIa), the formamidino- (XXXIVa), cyanoguanidino- (XXXVa), hydroxyguanidino- (XXXVIa) and guanidino- analogs (XXXVIIa) can prepared by the same procedures described for the aliphatic series in Schemes 4, 6 through 8. The boronic acid esters (XXXIIIa)-(XXXVIIa) can all be trransesterified to the corresponding free boronic acid inhibitors (XXXIIIb)-(XXXVIIb) using the phenyl boronic acid method prviously described.

Aromatic boronic acid inhibitors (XLIa,b), with the 15 guanidine functionality substituted directly on the aromatic nucleus, can be prepared from precursor (XXXVIII) according to Scheme 11. Nitration of the aromatic ring of (XXXVIII) according to the method of Olah and Kuhn, J. Amer. Chem. Soc. 84, 3684 (1962) can occur readily with agents such as acetyl nitrate, nitrosonium 20 tetrafluoroborate (NO2+BF4-) and nitrosonium hexafluorophosphate (NO2+PF6-) in inert solvents like tetrahydrofuran or dichloromethane at -78°C or higher. products of formula (XXXIX) can be reduced to the aniline derivative (LX) by catalytic hydrogenation. The catalysts 25 recommended for this procedure include iron, zinc, platinum oxide, rhodium - platinum oxide, palladium, Raney nickel, copper chromite, and rhenium sulfide. While reduction occurs readily under an atmosphere of hydrogen gas at 30 pressures range from 1 to 350 atmosphere in an inert solvent like water, an alcohol or ethyl acetate, other reagents which may affect this reaction are transfer agents such as hydrazine, formic acid or triethyl formate (Hudlicky, (1984), pp 73).

## Scheme 11.

WO 95/09634

- 5 The aniline (LX) can be converted to the phenylguanidine (LXIa) by the procedure described for (XVII) in Scheme 4.

  The transformation of (LXIa) to the free boronic acid (LXIb) is effected as in Scheme 4.
- The several types of inhibitors disclosed in this

  invention can be broadly classified by their electrophilic functional group A, as defined in Formula (I). The

compounds described below, unlike the boron containing peptides, utilize a highly electrophilic carbon atom at A to interact with the active site serine of thrombin. The precursor for the electrophilic carbon inhibitors is the appropriately protected amino acid (LXII) of Scheme 12.

# Scheme 12

 $Q = -(CH_2)_y$ - or  $-(CH_2)_{q-1}C_6H_4(CH_2)_{p-1}$ -

v = 0-10

M = alkyl or benzyl

PG = protecting group

The preparation of (LXII) can be found in the general chemical literature, one such reference being the review by Morrison and Mosher (1976). According to Scheme 12 various terminal functional groups are available from (XLII): the

formamidino- (XLIII), cyanoguanidino- (XLIV), hydroxyguanidino- (XLV) and guanidino- analogs (XLVI). These compounds are prepared by the same procedures described for the boropeptide series in Schemes 4, 6-8.

The preparation of amidine derivative (XLVIII) and phenylguanidines of formula (L) from amino acids (XLVII) and (XLIX) is shown in Scheme 13. The conditions used to prepare amidines of formula (XLVIII) is discussed for (XX) of Scheme 5 while the method for formamidinylation of (XLIX) to give (L) is the same as that described to prepare (XVII) of Scheme 4.

### Scheme 13.

 $Q = -(CH_2)_y$ - or  $-(CH_2)_{q-1}C_6H_4(CH_2)_{p-1}$ -

$$MO_2C$$
 $NPG$ 
 $NPG$ 
 $NH_2$ 
 $NHC(=NH)NH_2$ 
 $NHC(=NH)NH_2$ 

 $Q = -(CH_2)_{q-1}C_6H_4$ 

v = 0-10

M = alkyl or benzyl

PG = suitable amine protecting group

As shown in Scheme 14, appropriately protected derivatives of formulae (XLII)-(L), wherein M is an alkyl

15

5

or benzyl group can be coupled with N, N-disubstituted acid (IV) or (VI) (wherein M is hydrogen). The methodology to accomplish these transformations is the same as that used to prepare boropeptides (IX) and (X) of Scheme 3. The X group in compounds of formulae (XLII) through (L) and (LI) in Scheme 14, as well as in compounds illustrated in the Schemes to follow, is a protected version of the terminal functional group X, as defined by Formula (I), unless deprotection is indicated to obtain the final compound of the sequence.

### Scheme 14.

$$(VI): L = R^{3}R^{11}N$$

$$(VI): L = R^{1} - N$$

$$R^{3}$$

15

10

It is understood that the protecting group(s) used should compatible with the conditions of the process discussed; a good source for information on protecting group chemistry is Greene and Wuts (1991).

20

25

The preparation of the thrombin inhibitors trihalomethyl ketone (LIII) and a-ketoester (LIV) are shown in Scheme 15. The coupled ester (LI), wherein M is alkyl or benzyl can be converted to the acid (M is hydrogen) by the methodology appropriate for the particular ester functionality as described in Greene and Wuts (1984). The aldehyde (LII) can be prepared by selective reduction of the acid (LI, M is hydrogen) to the primary alcohol followed by oxidation.

## Scheme 15.

To obtain the primary alcohol, the acid can be transformed to the mixed anhydride by condensation of the trialkylammonium salt of the acid with an alkyl- or arylchloroformate in an inert non-polar solvent such as tetrahydrofuran, 1,2-dimethoxyethane or toluene at-78°C to 10 room temperature. The solution of the resulting mixed anhydride is filtered and reduced to the peptidyl alcohol with an excess of a borohydride reducing agent in a compatible solvent like water or an alcohol at -78°C to room temperature according to the method of Rodriguez et. 15 al., Tetrahedron Lett. 32, 923 (1991). The peptidyl alcohol can be oxidized to aldehyde (LII) without over oxidation by a variety of procedures, as detailed by Hudlicky in Oxidations in Organic Chemistry, American Chemical Society, p. 114 (1991); the preferred methods 20 include Swern oxidation described by Omura and Swern, Tetrahedron 34, 1651 (1978); and the Pfitzner-Moffat oxidation described by Fearon et al.in J. Med. Chem. 30, 1617 (1987). A two step protocol reported by Edwards,

PCT/US94/11280 WO 95/09634

the trifluoromethyl ketones (LIII) (J is fluorine) from aldehyde (LII). In this procedure a metallated trifluoromethyl anion is generated from an excess of trifluoromethyliodide or -bromide and an active metal such as zinc, magnesium, lithium or cadmium in inert, anhydrous 5 solvents like tetrahydrofuran or N,N-dimethylformamide at temperatures of -100°C up to the reflux point of the solvent. Alternatively, the metalated trifluoromethyl anion may be generated by the transmetallation of trifluoromethyliodide or -bromide with an organometallic 10 compound such as a Grignard reagent or alkyllithium compound in an inert solvent like tetrahydrofuran, hexane or ether at temperatures ranging from -78°C up to the reflux point of the selected solvent. Aldehyde (LII) can be added to the solution of the metalated trifluoromethyl 15 anion to form the trifluoroethanol derivative at temperatures of -100°C or higher. To obtain the trifluoromethyl ketone (LIII) where J is fluoro, the alcohol is oxidized by the Pfitzner-Moffat or Swern procedure. Removal of the protecting group(s) on terminal 20 group X by the appropriate method will provide the thrombin inhibitors of formulae (LIII).

Trihalomethyl analogs of (LIII), where J is fluoro can also be prepared from aldehyde (LII) by a different method. The trihalomethyl ketones are prepared by treating aldehyde (LII) with either the trimethylsilyl trihaloacetate or the potassium or sodium trihaloacetate in a polar solvent such as an alcohol, N,N-dimethylformamide or methylsulfoxide with or without a base such as a trialkyl amine, potassium 30 carbonate or sodium hydroxide at temperatures of -78°C or higher according to the method of Beaulieu, Tetrahedron Lett. 32, 1031 (1991); Shell Int. Res., European Patent Application EP 16504 ). The resulting a,a,a-trihaloethanol is oxidized and group X can be deprotected as above to give the thrombin inhibitors or formulae (LIII).

35

The a-ketoester thrombin inhibitors, exemplified by (LV), are prepared according to a route disclosed by Iwanowicz et. al. in Bioorgan. Med. Chem. Lett. 12, 1607 (1992). The tris(ethylthio)methyl anion is added to the peptidyl aldehyde (LII) in a solvent such as tetrahydrofuran, 1,2-dimethoxyethane or toluene at -100°C or higher to give the alcohol (LIV). The a-hydroxyl ester is generated from (LIV) by treatment with a mixture of mercuric salts, such as mercuric chloride and mercuric oxide, in an alcohol or water. Swern or Pfitzner-Moffat oxidation of the a-hydroxyl ester followed by the deprotection of substituent X protecting group provides thrombin inhibitors of formula (LV).

Another method for the preparation of compound (LV) 15 substitutes a 1-lithio-1-alkoxyethene or 1-magnesio-1alkoxyethene for the tris(ethylthio)methyl anion of Scheme 15 in an addition reaction with peptidyl aldehyde (LII). There can be obtained an adduct analogus to the tris(ethylthio)hydroxyethyl compound (LIV) when excess 1-20 magnesio- or 1-lithio-1-alkoxyethene anion is stirred at temperatures ranging from -100 °C to ambient temperature with (LII) in anhydrous solvents such as diethyl ether or tetrhydrofuran. This alkoxyolefin product may then be transformed to (LV) by oxidative cleavage with reagents such as ozone or periodate in an inert solvent such as a 25 halohydrocarbon, lower alkyl ketone, an alcohol or water at temperatures ranging from -100 °C to ambient temperature, followed by oxidation of the intervening a-hydroxyester and deprotection as described above.

The preparation of the a,a-dihalomethylketone thrombin inhibitors of this invention is outlined in Scheme 16.

PCT/US94/11280

Scheme 16.

R<sup>11</sup>N

R<sup>2</sup>

R<sup>3</sup>

(LII)

$$Q = (CH_2)_{1-12} \text{ or } (CH_2)_q C_6 H_4 (CH_2)_p$$

R<sup>11</sup>N

 $Q = (CH_2)_{1-12} \text{ or } (CH_2)_q C_6 H_4 (CH_2)_p$ 

(LVII)

The a,a-dihalomethylketone (LVII), where J is fluoro can be 5 prepared from the aldehyde (LII) by selective reaction of the aldehyde with the anion of the corresponding dihalomethane. The metalated dihalomethane anion is generated from one equivalent each of a strong hindered base, such as lithium tetramethylpiperidide or tertbutyllithium, and the selected dihalomethane in an anhydrous, inert solvent like tetrahydrofuran or 1,2dimethoxyethane at -100°C or higher according to the method of Taguchi et. al. Bull. Chem. Soc. Jpn., 50, 1588 (1977). The metalated dihalomethane anion can be added to the 15 aldehyde (LII) at -100°C or higher. Alternatively, the dihalomethane anion is generated from a dihalomethyl(trimethyl)silane and an anhydrous fluoride ion source such as tris(diethylamino)sulfonium difluoromethyl silicate in an inert solvent like benzene, acetonitrile or 20 tetrahydrofuran at -78°C or higher, then (LII) can be added to give dihaloethanol (LVI) according to the method of

WO 95/09634

Fujita and Hiyama, J. Am. Chem. Soc. 107, 4085 (1985). The resulting dihaloethanol can be oxidized to ketone (LVII) by the Swern or Pfitzner-Moffat procedure. Removal of the protecting group(s) on substituent X of (LVII) gives the a,a-dihalomethylketone thrombin inhibitors.

a-Halomethylketone thrombin inhibitors can be prepared by the process illustrated in Scheme 17. The acid chloride (LVIII) can be prepared from acid (LI), wherein M is hydrogen or its trialkylammonium, sodium or 10 potassium salt with a chlorinating agent such as thionyl chloride, oxalyl chloride or dichloromethylmethyl ether in a solvent like tetrahydrofuran or dichloromethane with or without a catalytic amount of N, N-dimethylformamide at -78°C or higher. Alternatively, the mixed anhydride of 15 (LI) may be prepared as described for (LI) in Scheme 15. Compound (LVIII) or the mixed anhydride of (LI) can be treated with an ether solution of diazomethane and either anhydrous hydrogen fluoride or hydrogen chloride gas according to that described by McPhee and Klingsbury, Org. 20 Synth. Coll. III, 119 (1955); or hydrogen bromide according to the method Miescher and Kaji, Helv. Chim. Acta. 24, 1471 (1941).

## Scheme 17.

$$R^{11}N \longrightarrow R^{3} \longrightarrow R^{4} \longrightarrow R^{5} \longrightarrow R^{5} \longrightarrow R^{4} \longrightarrow R^{5} \longrightarrow R^$$

5 Selection of the hydrogen fluoride gas will give the a-fluoromethylketone analog. (LIX) wherein J is fluoro; and hydrogen chloride gas gives the a-chloromethylketone analog (LIX) wherein J is chloro. Deprotection of X gives the corresponding thrombin inhibitors of (LIX).

The general preparative route for the a,b-diketoester, -amide and -ketone thrombin inhibitors of this invention is exemplified in Scheme 18. Compound (LVIII) or the mixed anhydride of (LI) can be reacted with a Wittig reagent such as methyl (triphenyl-phosphoranylidene)acetate in a solvent like tetrahydrofuran or acetonitrile at temperatures ranging from 0°C to the reflux point of the solvent to give (LX). Oxidative cleavage of the phosphoranylidene (LX) with an oxidizing agent like ozone or OXONE<sup>TM</sup> in an inert solvent such as tetrahydrofuran, dichloromethane or water at temperatures of -78°C or higher gives the vicinal

10

15

tricarbonyl compound (LXI), analogous to that described by Wasserman and Vu, *Tetrahedron Lett.* 31, 5205 (1990). Cleavage of the protecting group can provide thrombin inhibitors of formula (LXI).

5

### Scheme 18.

10

15

20

The preparative routes for the synthesis of the amono- and a,a-dihalo-b-ketoester -amide and ketone thrombin inhibitors of this invention are summarized in Scheme 19. The exemplified b-ketoester (LXII) is available from the acid derivative (LI). The acid (LI) can be treated with carbonyl diimidazole in an inert solvent such as tetrahydrofuran or dichloromethane at 0°C or higher to form the acyl imidazole. This acyl imidazole, or the mixed anhydride of (LI), can be further reacted with lithioethylacetate in solvents such as 1,2-dimethoxyethane or tetrahydrofuran/hexane at temperatures ranging from -100°C to ambient temperature, according to the method of Dow, J. Org. Chem. 55, 386 (1990) to give b-ketoester (LXII).

# Scheme 19.

$$R^{11}N$$

$$R^{3}$$

$$Q$$

$$X$$

$$C(O)CH_{2}CO_{2}CH_{3}$$

$$R^{11}N$$

$$Q$$

$$X$$

$$(LXIII)$$

(LXIII): J = H, halogen

 $Q = (CH_2)_{1-12}$  or  $(CH_2)_q C_6 H_4 (CH_2)_p$ 

(LXIV): J = dihalogen

Compound (LXII) serves as a substrate for both mono- and .5 dihalogenation. The a-monochloro analog of (LXIII), where J is each chlorine and hydrogen, can be prepared by controlled halogenation reactions with reagents like Nchlorosuccinimide or thionyl chloride in an inert halogenated solvent and at temperatures ranging from -20°C 10 to the reflux point of the selected solvent according to the methods of Uhle, J. Am. Chem. Soc. 83, 1460 (1961); and DeKimpe et. al., Synthesis 2, 188 (1987). The a,a-dihalo analog (LXIV) where J is chloro is available from halogenation with molecular chlorine in a halogenated 15 solvent at temperatures of -20°C or higher according to the method of Bigelow and Hanslick, Org. Syn. Coll. II, 244 (1943). Reagents such as N-

fluorobis[(trifluoromethyl)sulfonyl]imide are useful for the preparation of mono- and difluoro analogs (LXIII) and (LXIV) by reacting the appropriate stoichiometry of this reagent with (LXII) in a halogenated solvent at temperatures of -78°C or higher according to the method of Resnati and DesMarteau, J. Org. Chem. 56, 4925 (1991). Deprotection of substituent X of the halogenation products (LXIII) and (LXIV) can provide the corresponding thrombin inhibitors.

10 Compounds of formula (LXII) also serves as a substrate for the preparation of tricarbonyl derivatives such as (LXI) (Scheme 18). Condensation of (LXII) with an aldehyde, such as benzaldehyde, gives an b-ene-a,g-dione. This ene-dione can be oxidatively cleaved with reagents 15 like ozone or periodate to give tricarbonyl analog (LXI).

The preparation of the mono- and dihalomethylketone thrombin inhibitors is outlined in Scheme 20. The intermediates formed in the preparation of the a-mono- and a,a-dihalo-b-ketoester thrombin inhibitors of Scheme 19 can be used in these preparations.

#### Scheme 20.

$$R^{11}N$$
 $R^{3}$ 
 $R^{3}$ 
 $R^{3}$ 
 $R^{4}$ 
 $R^{5}$ 
 $R^{5}$ 
 $R^{11}N$ 
 $R^{3}$ 
 $R^{3}$ 
 $R^{3}$ 
 $R^{4}$ 
 $R^{5}$ 
 $R^{5}$ 
 $R^{4}$ 
 $R^{5}$ 
 $R^{5}$ 
 $R^{7}$ 
 $R^{7}$ 

(LXIII): J = halogen, H

(LIX): J = dihalogen

(LXIV): J = dihalogen

(LVII): J = halogen, H

25

20

The decarboxylation of these halogenation products, (LXIII) and (LXIV), can be effected by saponification of the ester with mild aqueous base such as potassium carbonate or sodium hydroxide in water miscible solvents like an

alcohol, tetrahydrofuran or N,N-dimethylformamide, followed by adjusting the pH to a range of 4 to 6. This mixture can be either stirred at ambient temperatures or heated at various temperatures up to the reflux point of the solvent chosen until the formation of (LVII) or (LIX) is complete and is similar to that reported in Matsuda et. al., Tetrahedron Lett. 30, 4259 (1989). Removal of protecting group(s) can provide thrombin inhibitors corresponding to (LVII) or (LIX).

10

15

Compounds of the present invention wherein the electrophilic group A is an a-hydroxy ester are prepared according to Scheme 21. The appropriate amino acid (LXVI) is reduced to the corresponding alcohol (LXVII) via NaBH4 treatment of the mixed anhydride. (LXVII) is then oxidized to the aldehyde (LXVIII) utilizing a Swern oxidation. (LXVIII) is converted to the thiocarbinol (XIX) via a lithiated orthoethylthioformate followed by conversion to the a-hydroxy methyl ester (LXX) upon treatment with mercuric salts. (LXX) is readily converted to the peptides of the invention via coupling with (IV) (M=H) to form (LXXI) under the conditions described in Scheme 3.

The preparation of compounds of the present invention wherein A is an a-keto ester, a-keto acid or a-keto ester hemiacetal is shown in Scheme 22. (LXXI) is converted to the a-keto ester LXXIII via a Swern Oxidation. (LXXIII) is further elaborated to hemiketal (LXXIV) by treating with methanol or to (LXXV) by treatment with hydroxide.

# Scheme 22.

The compounds of the present invention wherein A is an alkyl carbinol (LXXVII) or an alkyl ketone (LXXVIII) are prepared according to Scheme 23. (LXVIII) is treated with an alkyl-CeCl<sub>2</sub> to form the alkyl carbinol (LXXVII) which is then subjected to the Swern oxidation to yield the alkyl ketone (LXXVIII). (LXXVIII) is then further elaborated to the peptide compounds of the present invention (LXXX) by following the conditions outlined in Scheme 3.

Scheme 23.

1)RLi

2)CeCl<sub>3</sub>

3)H<sub>3</sub>O<sup>+</sup>

THF

-78°c

1) Oxalychloride

2)DMS

3)TEA

CH<sub>2</sub>Cl<sub>2</sub>

-10°c

R = alkyl

$$n = 2,3$$

LXXX

 $R^4$ 
 $R^5$ 
 $R^1$ 

LXXX

 $R^4$ 
 $R^5$ 
 $R^1$ 

Compounds of the present invention wherein R<sup>3</sup> is an optionally substituted thiophenylbenzoyl group (LXXXIII) are prepared according to Scheme 24. The desired thiophenol (LXXXII) is coupled to the bromobenzoic acid via a copper promoted coupling reaction to form (LXXXIII). (LXXXIII) is then coupled to (IV) (M=H) under the conditions outlined in Scheme 3 to form (LXXXIV) which is then coupled to (LXXXVI) under the conditions outlined in Scheme 3 to form (LXXXV). It is understood that (LXXXV) need not be in final form and that any or all of the functional groups present may be converted to their desired final form using methods known to the skilled artisan.

Compounds of the present invention wherein R<sup>3</sup> is an optionally substituted benzylbenzoyl group are prepared according to Scheme 25. An optionally substituted bromophenyl compound (LXXXVII) is converted to its phenyl lithium derivative and reacted with a bromobenzaldehyde to form the optionally substituted diphenylmethyl carbinol (LXXXVIII). (LXXXVIII) is further reduced to the substituted diphenylmethane (LXXXIX) with Et<sub>3</sub>SiH. (LXXXIX) is converted to XC. XC is then further elaborated to XCI and XCII by following the appropriate outline in previous schemes.

Ļ

Inhibitors which contain a substituted phenethyl group as R11 are easily prepared according to Scheme 26. The appropriate phenylacetate (XCIV) is readily dialkylated with an excess of a small, unbranched alkyl halide (R18-X) and a suitable base such as potassium tert-butoxide to form an a,a-bisalkylated ester (XCVa). Reduction of XCVa to the

primary alcohol may accomplished with many hydride reducing agents, a preferred agent being lithium aluminum hydride.

Oxidation of the alcohol under Swern conditions or with pyridinium chlorochromate affords the aldehyde (XCV).

(XCV) is best coupled to the appropriate glycine derivative by reductive amination, a preferred procedure being reduction with sodium cyanoborohydride. The resulting amine (XCVI) is then coupled with R3 by any of several standard amide bond forming reactions familiar to those skilled in the art. A preferred method involves treating the amine with the appropriate acid chloride in the presence of a tertiary amine base, such as N-methylmorpholine or triethylamine. Saponification of the ester affords the carboxylic acid (XCVII).

The acid (XCVII) is then coupled to (LXXXVI) and elaborated to the inhibitors of the present invention by following the procedures outlined in Schemes 3, 4, and 5.

Inhibitors which contain a 1,w-alkanediyl substituted

phenethyl group (XCVb) as R<sup>11</sup> are easily prepared according
to Scheme 27. This procedure is similar to that of Scheme
26, except that a 1,w-bifunctional alkylating agent (X-R<sup>18</sup>X) instead of a monofunctional alkylating agent (R<sup>18</sup>-X).
In this manner, inhibitors of the present invention with
the general formula (XCVIIIb) may be prepared.

Inhibitors in which R<sup>3</sup> is an acylalkyl terminated by a carboxylic acid or ester (CI) are prepared by the general route described in Scheme 28. Reaction of a suitably substituted cyclic anhydride (C) with an alkoxide such as sodium benzyl oxide affords a mono-protected diacid (CI).

(CI) is then coupled to an appropriate N-alkylglycine

10 derivative ((IV), (M=H) by any of a number of methods known
to the skilled artisan. The choice of ester groups should
allow for selective deprotection of the glycine
carboxylate. Preferred ester groups are methyl or ethyl on
the glycine carboxylate and benzyl on the acylalkyl chain,

so that saponification gives the acid (CII). (CII) is then converted to the final products (CIII) following methods described in previous schemes.

5

## Experimental Section

#### Intermediate 1

10

15

20

N-Methyl-N-[(3-phenyl)propionyl]glycine

Part A: To hydrocinnamic acid (10.0 g, 66.7 mmol) and 4-methylmorpholine (6.74 g, 66.7 mmol) in tetrahydrofuran (THF, 200 mL) at 0°C was added n-butylchloroformate. The reaction was maintained at 0°C for 15 minutes, and the hydrochloride salt of sarcosine ethyl ester (10.23 g, 66.7 mmol) followed by triethylamine (Et<sub>3</sub>N, 16.84 g, 166.8 mmol) was added. The reaction was allowed to thaw to ambient temperature and stirred for 18 hours. After this time, the solvent was removed and the residue partitioned between aqueous hydrochloric acid (HCl, 1 N, 200 mL) and ethyl acetate (EtOAc, 200 mL). The aqueous acid phase was extracted with additional EtOAc (200 mL), the combined

organic extracts were washed with HCl (1 N, 100 mL), saturated sodium bicarbonate (NaHCO $_3$ , 100 mL) and brine (100 mL). The organic phase was dried over sodium sulfate and evaporated to give ethyl N-methyl-N-[(3-phenyl)propionyl]glycine (11.81 g, 71% yield). This material was used in the next step without further purification.

part B: To a solution of ethyl N-methyl-N-[(310 phenyl)propionyl]glycine (11.81 g, 47.4 mmol) in ethanol
 (300 mL) was added aqueous sodium hydroxide (NaOH, 1 N,
94.8 mL, 94.8 mmol). The reaction was stirred at ambient
 temperature for 18 hour, afterwhich the solvent was removed
 by distillation in vacuo. The residue was dissolved in HCl
15 (1 N, 100 mL) and the solution extracted with methylene
 chloride (CH<sub>2</sub>Cl<sub>2</sub>, 2 x 100 mL). The extracts were dried over
 sodium sulfate (Na<sub>2</sub>SO<sub>4</sub>), evaporated and the resulting solid
 (9.42 g) was recrystallized from EtOAc to give the title
 compound (7.14 g, 68% yield) as a solid (mp: 123 - 126 °C).

20

# Example 21.1.3

Pinanediol  $N-\{N-\text{methyl-}N-\{(3-\text{phenyl})\text{propionyl}\}\text{glycyl}\}-1-$ amido-5-aminopentaneboronate, benzenesulfonic acid salt

Part A: A mixture of Intermediate 1 (0.31 g, 1.5 mmol), pinanediol 1-amino-5-bromopentaneboronate (0.57 g, 1.85 mmol), 1-hydroxybenzotriazole (0.20 g, 1.5 mmol), 4-methylmorpholine (0.17 mL, 1.5 mmol), and 1,3-dicyclohexylcarbodiimide (DCC, 0.33 g,1.5 mmol) were stirred in dry CH<sub>2</sub>Cl<sub>2</sub> at 0°C for 1 hour. The reaction was thawed to ambient temperature and stirred an additional 18 hour. After this time, the reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (25 mL) and filtered. The filtrate was washed with aqueous citric acid (10 %) and saturated NaHCO<sub>3</sub> (25 mL each), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The intermediate

pinanediol  $N-(N-\text{methyl}-N-[(3-\text{phenyl})\text{propionyl}]glycyl}-1-$ amido-5-bromopentaneboronate (0.75 g, 92% yield) was carried on to the next step without further purification.

- 5 Part B: The intermediate from Part A (0.75 g, 1.4 mmol) was heated with sodium azide (NaN3, 0.15 g, 2.3 mmol) in N,N-dimethylformamide (DMF, 10 mL) at 100°C for 2 hours. The reaction mixture was partitioned between water (H2O) and EtOAc (25 mL each), and the EtOAc layer was washed with additional H2O (6 x 15 mL). The organic layer was dried (Na2SO4) and evaporated to give pinanediol N-{N-methyl-N-[(3-phenyl)propionyl]glycyl}-1-amido-5-azidopentaneboronate (0.67 g) in 95% yield.
- 15 Part C: The azide from Part B (0.48 g. 0.9 mmol) was dissolved in methanol (MeOH, 15 mL) with benzenesulfonic acid (0.15 g, 0.9 mmol) and Pearlman's catalyst (palladium hydroxide on carbon, 0.05 g). This mixture was shaken under an atmosphere of 50 psi of hydrogen for 18 hours at ambient 20 temperature. The reaction mixture was purged with nitrogen and the catalyst was removed by filtration through a pad of diatomaceous earth. The clear filtrate was evaporated and pinanediol N-{N-methyl-N-[(3-phenyl)propionyl]glycyl}-1amido-5-aminopentaneboronate was obtained as its 25 benzenesulfonate salt (0.51 g) in 88% yield. High Res Mass Spec: found  $(M+H)^{+}$  484.334758; calculated  $(M+H)^{+}$  = 484.334663.

#### <u>Intermediate 2</u>

N-[(2-Phenyl)ethyl]-N-[(3-phenyl)propionyl]glycine

Part A: A mixture of benzyl glycinate, p-toluenesulfonic acid salt (2.68 g, 7.94 mmol), (2-phenyl)bromoethane (0.98 g, 5.29 mmol), and solid NaHCO<sub>3</sub> (1.56 g, 18.5 mmol) in acetonitrile (25 mL) were heated at reflux for 18 hour.

The reaction was concentrated and diluted with EtOAc (25 mL). The organic solution was washed with  $H_2O$  (25 mL) and brine (25 mL), dried ( $Na_2SO_4$ ) and concentrated in vacuo. The residue was purified by elution through a pad of silica gel with a gradient mixture of hexane:EtOAc. The intermediate benzyl N-[(2-phenyl)ethyl]-glycinate (0.82 g) was obtained in 38% yield. Low Res MS:  $(M+H)^+=270$ .

Part B: A mixture of intermediate from Part A (0.82 g, 3.04 mmol) and 4-methylmorpholine (0.62 g, 6.08 mmol) in THF (15 mL) at 0°C was added hydrocinnamoyl chloride (0.51 g, 3.04 mmol). The reaction was thawed to ambient temperature and stirred for 1 hour. The reaction mixture was diluted with EtOAc (50 mL), washed with HCl (10%, 25 mL) and NaHCO3 (saturated, 25 mL), dried (MgSO4) and evaporated. The intermediate benzyl N-[(2-phenyl)ethyl]-N-[(3-phenyl)-propionyl]glycinate (1.2 g) prepared was used in the next procedure without further purification. LRMS: (M+NH3)+ = 419.0, (M+H)+ = 402.1.

20

Part B A methanol solution (20 mL) of benzyl N-[(2-phenyl)ethyl]-N-[(3-phenyl)propionyl]glycinate (1.3 g, 3.24 mmol) and palladium on carbon (10%, 180 mg) was stirred under 1 atmosphere of hydrogen gas for 18 hours. The reaction was purged with nitrogen and filtered through a pad of diatomaceous earth and evaporated to give N-[(2-phenyl)ethyl]-N-[(3-phenyl)propionyl]glycine (1.0 g) in quantitative yield. LRMS:  $(M+NH_3)^+ = 326$ .

30

25

### Example 23.1.3

Pinanediol N-{N-[(2-phenyl)ethyl]-N-[(3-phenyl)propionyl]glycyl}-1-amido-5-aminopentaneboronate, hydrochloride salt

To a solution of N-[(2-phenyl)ethyl]-N-[(3-phenyl)ethyl]phenyl)propionyl]glycine (1.0 g 3.24 mmol) and 4methylmorpholine (0.66 g, 6.48 mmol) in THF (15 mL) at 0°C was added isobutylchloroformate (0.44 g, 3.24 mmol). The 5 reaction was stirred for 15 min at 0°C, pinanediol 1-amino-5-bromopentaneboronate (1.23 g, 3.24 mmol) was added followed by additional 4-methylmorpholine (0.33 g, 3.24 mmol) and the reaction was stirred at ambient temperature for 18 hours. The reaction was diluted with EtOAc (50 mL), 10 washed sequentially with HCl (10%) and NaHCO3 (saturated) and brine (25 mL each), then dried over magnesium sulfate (MgSO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography (silica gel) using 3:1 EtOAc:hexane to give pinanediol  $N-\{N-\{(2-phenyl)ethyl\}-N-\{(3-phenyl)ethyl\}\}$ 15 phenyl)propionyl]glycyl}-1-amido-5-bromopentaneboronate (0.8 g) in 43% yield. LRMS:  $(M+H)^+ = 637/638$ ,  $(M-HBr)^+ =$ 

- Part B: The intermediate pinanediol N-{N-{(2-20 phenyl)ethyl]-N-{(3-phenyl)propionyl}glycyl}-1-amido-5-bromopentaneboronate (0.8 g, 1.4 mmol) and NaN3 (0.11 g, 1.7 mmol) in DMF (10 mL) was stirred at 100°C for 2 h. The cooled reaction mixture was diluted with EtOAc (50 mL), then it was washed with H2O (6 x 20 mL) and dried (MgSO4).

  The EtOAc solution was evaporated to give pinanediol N-{N-
- The EtOAc solution was evaporated to give pinanediol  $N-\{N-\{(2-phenyl)ethyl\}-N-\{(3-phenyl)propionyl\}glycyl\}-1-amido-5-azidopentaneboronate (0.7 g) in 86% yield. LRMS: <math>(M+H)^+=600$ .
- 30 Part C: A mixture of the product from Part B (0.7 g, 1.2 mmol) and Pearlman's catalyst (0.1 g) in HCl (1.2 N, 1 mL, 1.2 mmol) and MeOH (20 mL) was stirred under an atmosphere of hydrogen (1 atm) for 2 hours. The reaction mixture was purged with nitrogen, filtered through a pad of diatomaceous earth and evaporated. The residue was dried

557.

by azeotropic distillation with benzene and triturated with hexane to give the title compound  $(0.45~\rm g)$  as a yellow powder in 67% yield. LRMS:  $(M+H)^+ = 574.4$ .

# Example 8.1.3

 $N-\{N-\{(2-\text{Pheny1})\text{ ethy1}\}-N-\{(3-\text{pheny1})\text{ propiony1}\}\text{ glycy1}\}-1-$ amido-5-aminopentaneboronic acid, hydrogen chloride salt

A mixture of Example 23.1.3 (0.45 g, 0.8 mmol) in diethyl ether(Et<sub>2</sub>O): H<sub>2</sub>O (15 mL:15 mL) was stirred with phenylboric acid (0.45 g, 3.7 mmol) at ambient temperature for 18 hours. The phases were separated, the Et<sub>2</sub>O layer was discarded, and the H<sub>2</sub>O layer was washed with Et<sub>2</sub>O (15 mL) and 1:1 hexane:EtOAc (15 mL). The H<sub>2</sub>O solution was concentrated by distillation under reduced pressure anddried by azeotropic distillation with toluene. The dried residue was dissolved in a minimal amount of CH<sub>2</sub>Cl<sub>2</sub> and the title compound was precipitated from solution by the addition of hexane. LRMS: (M+H of glycerol ester)+ = 496.

#### Intermediate 3

N-methyl-N-[(3,4-dichlorophenyl)acetyl]glycine

25 The title compound was prepared from commercially available sarcosine ethyl ester and 3,4-dichlorophenylacetic acid according to the procedure of Intermediate 1.

#### Example 50

Pinanediol N-{N-methyl-N-[(3,4-dichlorophenyl) acetyl]glycyl}-1-amido-4-formamidinobutaneboronate, hydrochloride salt

Part A: By substituting pinanediol 1-amino-4-35 bromobutaneboronate for pinanediol 1-amino-5-

bromopentaneboronate and coupling with Intermediate 3 according to the procedure in Example 21.1.3, Parts A-C, the amino intermediate was prepared.

5 Part B: To an ethanol solution (20 mL) of the product from Part A (600 mg, 1.2 mmol) was added ethylformimidate hydrogen chloride (400 mg, 3.62 mmol) and 4-dimethylamino pyridine (DMAP, 442 mg, 3.62 mmol). This mixture was heated at reflux for 5 hours, and the reaction mixture was evaporated. The residue was chromatographed (Sephedex LH 20, MeOH elutant) to give the title compound as a yellow solid. LRMS: (M+H)<sup>+</sup> = 551.

15

#### Example 51

Pinanediol N-{N-methyl-N-[(3,4-dichlorophenyl) acetyl]glycyl}-1-amido-4-guanidinobutaneboronate, hydrochloride salt

Part A: The intermediate from Example 50, Part A hydrochloride salt (1.0 g, 2 mmol), formamidine sulfonic acid (0.496 g, 4 mmol) and DMAP (0.488 g, 4 mmol) in ethanol (50 mL) were heated at reflux for 3 hours. The reaction was cooled to ambient temperature, filtered through a pad of Celite, rinsed with chloroform (CHCl<sub>3</sub>) and evaporated. The residue was dissolved in CHCl<sub>3</sub> and washed with HCl (0.1 N) and brine, dried and evaporated. The title compound was obtained as a white solid. HRMS calcd for C26H39BN5O4Cl<sub>2</sub>:582.247216+; found: 566.247905.

30

#### Intermediate 4

Pinanediol 1-amino-2-(3-cyanophenyl)ethylboronate, hydrochloride salt

The intermediate, C1-CH[CH2-(m-cyanophenyl)]BO2- $C_{10}H_{16}$ , was prepared from m-cyanobenzyl bromide and dichloromethyl boronate pinanediol. Zinc dust (1.0 g) in THF (1 mL) was cooled to  $0-5^{\circ}$ C and a solution of mcyanobenzyl bromide (1.37 g, 7.0 mmol) in THF (7 mL) was 5 added dropwise (5 sec/drop). The reaction mixture was allowed to stir at 5°C for 2 hours. A mixture consisting of lithium bromide (LiBr, 1.22 g, 14 mmol), copper(I) cyanide (CuCN, 0.63 g, 7.0 mmol) and THF (6 mL) was placed in a 50 mL flask and cooled to -40°C; the benzylic 10 organozinc reagent was added by cannulation. The mixture was allowed to warm to -20°C and stir for 5 minutes. Following cooling to -78°C, neat dichloromethyl boronic acid pinanediol (1.47 g, 5.6 mmol) was added dropwise and the resulting mixture was stirred at -78°C for 2 h, and 15 additionally at room temperature for 2 days. ammonium chloride (NH4Cl, saturated, 20 mL) was added to the mixture and the aqueous solution was extracted with  ${\tt Et_2O}$  (3 x 20 mL). The combined organic layers was dried over anhydrous MgSO4 and evaporated in vacuo to give crude 20 compound (1.8 g). Purification was carried out using silica gel chromatography where the column was stepwise eluted with hexane (100 mL) and then 15% ether in hexane (200 mL) to give the desired product (0.53 g) in 27% yield. LRMS(NH<sub>3</sub>-CI) m/e for M+NH<sub>4</sub>+ calcd. for  $C_{19}H_{23}NO_2BCl$ : 361.2; 25 found: 361.1.

Part B: To a solution of hexamethyldisilazane (0.21 mL, 0.98 mmol) in THF (2 mL) at -78°C was added n-butyllithium (1.45 M, 0.67 mL, 0.98 mmol). The solution was allowed to slowly warm to room temperature to ensure the anion generation was complete and recooled to -78°C, upon which a solution of product from Part A (0.33 g, 0.98 mmol) in THF (2 mL) was added. The mixture was allowed to warm to room temperature and to stir overnight. The volatiles were

evaporated and hexane (8 mL) was added to give a suspension. Anhydrous hydrogen chloride in dioxane (4.1 N, 1.5 mL, 6.0 mmol) was added at -78°C and the mixture was slowly warmed to room temperature and stirred for 2 hour.

5 Additional hexane (6 mL) was added and crude product was isolated as a precipitate. This product was dissolved in CHCl<sub>3</sub> and insoluble material was removed by filtration. The filtrate was evaporated at a reduced pressure to give an oil (~0.2 g). Final purification was achieved by chromatography on a column of Sephedex<sup>TM</sup> LH 20 column using MeOH as a solvent. H-boroPhe(m-CN)-CloHl6·HCl was obtained as an oil (0.12 g) in 34% yield. HRMS(NH<sub>3</sub>-CI) m/e (M+H)+ calcd. for Cl9H<sub>26</sub>BN<sub>2</sub>O<sub>2</sub>: 325.2087; found: 325.2094.

15

# Example 52

Pinanediol N-{N-methyl-N-[(3,4-dichlorophenyl)acetyl]glycyl}-1-amido-2-(3-cyanophenyl)ethylboronate

20

25

30

Part A: To Intermediate 3 (0.77 g, 2.8 mmol) and 4-methylmorpholine (0.28 g, 2.8 mmol) in THF (50 mL) at -20°C was added isobutylchloroformate (0.38 g, 2.8 mmol). After 20 minutes at -20°C, the cold solution was added to a -20°C solution of Intermediate 4 (1.0 g, 2.8 mmol) and Et<sub>3</sub>N (0.28 g, 2.8 mmol) in CHCl<sub>3</sub> (50 mL). This mixture was maintained at -20°C for 5 hours and stirred at ambient temperature for 18 hours. The reaction was filtered and concentrated in vacuo. The residue was dissolved in EtOAc, washed with HCl (0.1N), saturated NaHCO<sub>3</sub> and brine. After the solution was dried and evaporated, the resulting yellow solid was applied to a column of Florisil and the desired product eluted with a gradient of CHCl<sub>3</sub>: MeOH (0% MeOH to 7% MeOH). The title compound was obtained as a white solid. HRMS:

PCT/US94/11280 WO 95/09634

calcd. (M+H)+ for C<sub>30</sub>H<sub>35</sub>BN<sub>3</sub>O<sub>4</sub>Cl<sub>2</sub>: 582.209768; found: 582.209631.

5

# Example 53

Pinanediol N-{N-methyl-N-[(3-phenyl)propionyl]glycyl}-1amido-4-(N-methylguanidino)butylboronate, hydrochloride salt

Pinanediol N-{N-methyl-N-[(3phenyl)propionyl]glycyl}-1-amido-4-aminobutylboronate, 10 hydrochloride salt was prepared by the method outlined for Example 21.1.3, wherein pinanediol 1-amino-4bromobutylboronate hydrochloride was used instead of pinanediol 1-amino-5-bromopentane-boronate hydrochloride.

15

20

To a solution of the product from Part A (0.45 g, Part B: 0.89 mmol) in ethanol (10 mL) was added DMAP (0.22 g, 1.78 mmol). After 15 minutes at room temperature, Nmethylaminoiminomethanesulfonic acid (0.25 g, 1.78 mmol) was added and the resulting suspension stirred at reflux for 5 hours. The reaction was cooled to room temperature, filtered, the precipitate washed with CHCl3 and the combined filtrate concentrated under vacuum. The resulting oil was dissolved in CHCl<sub>3</sub> (40 mL) and the organic solution washed with ice cold HCl (0.1 M,  $1 \times 10$  mL), ice cold H<sub>2</sub>O 25 (1x10 mL), brine and dried (MgSO4). The filtered solution was concentrated in vacuo to give of the desired N-methyl guanidino compound (0.35 g) in 70% yield. The material was purified through a florisil column using 10% MeOH/CHCl3 as eluant to give the purified product (0.22 g); LRMS: (M+H)+: 526 .

35

#### Example 54

# Ac-(D) Phe-Sar-boroLys-OH

Boc-(D)Phe-OH (6.9 g, 26 mmoles)was dissolved in THF (50 mL) and 4-methylmorpholine (2.86 mL, 26 mmoles) was added. The solution was cooled to -20°C and isobutylchloroformate (3.38 ml, 26 mmoles) was added. After stirring 5 minutes at -20°C, the mixture was added to a cold solution of H-Sar-Bzl·toluenesulfonic acid dissolved in CHCl $_3$  (50 mL), followed by Et $_3N$  (3.6 mL, 26 mmoles). 10 The mixture was allowed to stir for 1 hour at -20°C and 2 h at room temperature. The solids were removed by filtration and the solvent was removed by evaporation. The residue was dissolved in EtOAc and was washed with HCl (0.20 N), NaHCO3 (5%), and brine. The solution was dried over 15 anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to yield Boc-(D) Phe-Sar-O-Bzl as thick oil (10.4 g).

Part B: The product from Part A (10.4 g) was dissolved in 20 MeOH (100 mL) and the sample was hydrogenated for 2 hour on a Parr apparatus in the presence of palladium on carbon (10%, 0.5 g). The catalysis was removed by filtration and solvent was evaporated to yield Boc-(D)Phe-Sar-OH as a foam (7.8 g).

25

30

Part C: The mixed anhydride of product from Part B (4.42 g, 13.1 mmoles) was prepared as previously described and coupled to NH2-CH[(CH2)4Br]BO2C10H16•HCl (5.0 g, 13.1 mmoles) using the procedure described in Part A. The crude product (7.7 g) was purified by chromatography of a 4.2 g portion on a 2.5 x 100 cm column of Sephedex LH-20 using MeOH as a solvent to give Boc-(D)Phe-Sar-NH-CH[(CH2)4-Br]BO2C10H16.

Part D: The product from Part C (3.5~g,~5.8~mmoles) was dissolved in anhydrous HCl in dioxane (4.1~N,~50~mL) and was stirred for 1 hour at room temperature. Solvent and excess HCl were removed by evaporation. The residue was triturated with hexane to yield H-(D)Phe-Sar-NH-CH[(CH<sub>2</sub>)4-Br]BO2C10H16•HCl (2.9~g).

The product from Part D (2.9 g, 4.8 mmoles) was Part E: dissolved in 30 mL of a 50% (v/v) solution of dioxane: water and acetic anhydride (0.92 ml, 9.7 mmoles) was added. 10 NaHCO3 (0.81 g, 9.7 mmoles) was added and the solution was allowed to stir 45 minutes at room temperature. Acetic acid (3 ml) was added and solution was concentrated It was diluted to 100 mL approximately 50% by evaporation. with EtOAc and was washed with NaHCO3 (5%), HCl (0.2 N), 15 and brine. The organic phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to yield Ac-(D)Phe-Sar-NH- $CH[(CH_2)_4-Br]BO_2C_{10}H_{16}$  as a foam (2.7 g). HRMS calcd for C29H43N3O5BBr (M+H): 604.2557; found: 604.2558.

20

25

Part F: The product from Part E (2.5 g, 4.1 mmoles) and NaN3 (0.54 g, 8.3 mmoles) were dissolved in DMF (5 mL) and heated at  $100^{\circ}$ C for 1 hour. The reaction was allowed to cool and was diluted with EtOAc (100 mL). The organic phase was washed with H2O and saturated brine, dried over Na2SO4, filtered, and evaporated to yield Ac-(D)Phe-Sar-NH-CH[(CH2)4-N3]BO2C10H16 as a white foam (2.2 g).

Part G: The product from Part F (2.0 g, 3.5 mmoles) was dissolved in MeOH (100 mL) and was hydrogenated on a Parr apparatus in the presence of HCl in dioxane (4.1 N, 1.3 ml, 5.3 mmoles) and palladium on carbon (10%, 0.5 g). The catalysis was removed by filtration and solvent was removed by evaporation. The product, Ac-(D)Phe-Pro-boroLys
CloH16·HCl (Ac-(D)Phe-Sar-NH-CH[(CH2)4-NH2]BO2CloH16·HCl),

was purified by chromatography on 2.5 X 100 cm column of LH-20 in MeOH to yield 1.8 g.

The product from Part G (1.5 g, 2.5 mmoles) and Part H: phenyl boronic acid (1.5 g, 12 mmoles) was dissolved in  $H_2O$ 5 and  $Et_2O$  (15 ml each). The mixture was stoppered and allowed to stir for 3 hour at room temperature. The phases were separated and the aqueous phase was washed extensively with Et20. The aqueous phase was evaporated, dried in vacuo., and triturated with Et<sub>2</sub>O to yield the title 10 compound (0.98 g). An analytical sample was prepared as the pinacol ester by treating 4 mg of the boronic acid with 2 equivalents of pinacol in 1.4 ml of MeOH for 5 minutes and evaporating solvent. HRMS calcd for the pinacol ester 15 C<sub>25</sub>H<sub>4</sub>1N<sub>4</sub>O<sub>5</sub>B (M+H): 489.3248. found: 489.3242.

# Example 55

Pinanediol N-{N-2-propyl-N-[(3-phenyl)propionyl]glycyl}-1-20 amido-5-aminopentaneboronate, hydrochloride salt

Part A: A mixture of glycine methyl ester hydrochloride (3.83 g, 30.5 mmol), acetone (1.77 g, 30.5 mmol) and NaOH (1.22 g, 30.5 mmol) in MeOH (200 mL) was stirred under an atmosphere of hydrogen (1 atm) in the presence of palladium on carbon (10%, 0.4 g) for 24 hours. The reaction was flushed with nitrogen and filtered through a Celite pad, acidified with HCl (1 $\underline{N}$ ), dried (MgSO<sub>4</sub>) and evaporated. Trituration of the residue with Et<sub>2</sub>O gave N-2-propylglycine methyl ester hydrochloride (1.0 g) as an off-white solid; LRMS (M+H)+ = 132.1.

Part B: To the hydrochloride salt prepared above (1.0 g, 5.97 mmol) and hydrocinnamic acid (0.9 g, 5.97 mmol) in DMF (20 mL) was added O-benzotriazole-N,N,N',N'-

25

tetramethyluronium hexafluorophosphate (2.26 g, 5.97 mmol) followed by N,N-diisopropylethylamine (1.69 g, 13.1 mmol). The reaction was stirred at ambient temperature for 48 hours. The reaction mixture was diluted with 1:1

5 EtOAc:hexane and washed with H<sub>2</sub>O (2 x), HCl (10%), saturated NaHCO<sub>3</sub> and brine. The solution was dried (MgSO<sub>4</sub>), evaporated, and combined with an additional material obtained by the same acylation procedure. The combined batches were purified by flash chromatography on silica gel with 2:1 hexane:EtOAc as the eluent. There was obtained N-2-propyl-N-[(3-phenyl)propionyl]glycine methyl ester (2.8 g); LRMS (M+H)+ = 264.0.

Part C: A mixture of the ester from Part B (2.8 g, 10.52 mmol) and LiOH monohydrate in THF:H<sub>2</sub>O (20 mL:10 mL) was stirred at ambient temperature for 18 hours. The reaction was diluted with H<sub>2</sub>O and washed with 1:1 hexane:EtOAc and the organic washings were discarded. The aqueous layer was acidified with HCl (10%) and extracted with EtOAc. The EtOAc extract was washed with brine, dried (MgSO<sub>4</sub>) and evaporated to give N-2-propyl-N-[(3-phenyl)propionyl]glycine (2.0 g); LRMS (M+H)+ = 250.1.

Part D: N-2-propyl-N-[(3-phenyl)propionyl]glycine was

reacted with pinanediol 1-amino-5-bromopentaneboronate
according to the procedure of Example 23.1.3 to prepare
pinanediol N-{N-2-isopropyl-N-[(3-phenyl)propionyl]glycyl}1-amido-5-bromopentaneboronate; LRMS (M+H)+ = 575/577.

30 Part E-F: This material was reacted with NaN<sub>3</sub> according to the procedure in Example 23.1.3 to give pinanediol N-{N-2-isopropyl-N-[(3-phenyl)propionyl]glycyl)-1-amido-5-azidopentaneboronate; LRMS (M+H)+ = 538.3. The azide was hydrogenated under the conditions in Example 23.1.3 to give the title compound, pinanediol N-{N-2-propyl-N-[(3-

phenyl)propionyl]-glycyl}-1-amido-5-aminopentaneboronate hydrochloride; LRMS  $(M+H)^+ = 512$ .

5

### Example 56

 $N-\{N-2-\text{propyl}-N-[(3-\text{phenyl})\text{propionyl}]\text{glycyl}\}-1-\text{amidosomiopentaneboronic acid, hydrochloride salt}$ 

Part A:  $N-\{N-2-\text{propyl}-N-[(3-\text{phenyl})\text{propionyl}]\text{glycyl}\}-1-10$  amido-5-aminopentane-boronic acid, hydrochloride salt, was prepared from pinanediol  $N-\{N-2-\text{propyl}-N-\{(3-\text{phenyl})\text{propionyl}]\text{glycyl}\}-1-amido-5-aminopentaneboronate, hydrochloride salt by the procedure of Example 8.1.3; LRMS <math>(M+H-H_2O)^+=360.1$ ,  $(M+H-2H_2O)^+=342.0$ .

15

#### Example 57

Pinanediol  $N-\{N-\text{methyl}-N-\{2-\text{(methylphenyl)benzoyl}\}glycyl}-1-amido-5-aminopentaneboronate, hydrochloride salt$ 

20

A mixture of 2-(methylphenyl)benzoic acid (3.09 Part A: g, 14.55 mmol), sarcosine ethyl ester hydrochloride salt (2.23 g, 14.55 mmol), DCC (3.0 g, 14.55 mmol), HOBT (1.97 g, 14.55 mmol) and Et<sub>3</sub>N (1.47 g, 14.55 mmol) in THF (50 mL) 25 were stirred at ambient temperature for 48 hours. reaction was evaporated and the residue dissolved in EtOAc. The EtOAc solution was washed with HCl (10%), saturated NaHCO3 and brine, and dried (MgSO4). The EtOAc solution was filtered through a plug of silica gel follwoed by a 30 second filtration through a plug of neutral alumina. Evaporation of the solution gave N-methyl-N-[2-(methylphenyl)benzoyl]glycine ethyl ester (4.5 g); LRMS  $(M+H)^+ = 312.2.$ 

Part B: N-methyl-N-[2-(methylphenyl)benzoyl]glycine ethyl ester (4.5 g, 14.45 mmol) and KOH (2.43 g, 43.4 mmol) in MeOH/H<sub>2</sub>O (200 mL/50 mL) were heated at reflux for 45 minutes. The solvent was removed and the residue dissolved in H<sub>2</sub>O. The aqueous solution was washed with Et<sub>2</sub>O and acidified with HCl (10%). The acidified aqueous layer was extracted with EtOAc, the EtOAc extract was washed with brine (2 x), dried over (MgSO<sub>4</sub>) and evaporated. There was obtained N-methyl-N-[2-(methylphenyl)benzoyl]glycine (2.0 g); LRMS (M+H)+ = 284.1, (M+NH<sub>4</sub>)+ = 301.1

Part C-E: Pinanediol  $N-\{N-\text{methyl}-N-[2-\text{methyl}-N]\}$ (methylphenyl)benzoyl]glycyl}-1-amido-5-bromopentaneboronate was prepared by reaction of Nmethyl-N-[2-(methylphenyl)benzoyl]glycine with pinanediol15 1-amino-5-bromopentaneboronate according to the procedure of Example 23.1.3; LRMS (M+H)+=611.2. Pinanediol  $N-\{N-1\}$ methyl-N-[2-(methylphenyl)benzoyl]glycyl}-1-amido-5bromopentaneboronate was reacted withe NaN3 by under the conditions detailed above to give pinanediol N-{N-methyl-N-20 [2-(methylphenyl)benzoyl]glycyl}-1-amido-5-azidopentaneboronate; LRMS (M+H)+=572.4. Hydrogenation of the azide was effected by the conditions previously described to give the title compound, pinanediol N-(Nmethyl-N-[2-(methylphenyl)benzoyl]glycyl}-1-amido-5-amino-25 pentaneboronate, hydrochloride acid salt; LRMS (M+H)+ = 546.3.

30

35

## Example 35.5.3

Part A: Trimethylsilyl cyanide (5.80 mL, 44.0 mmol) was added dropwise to a solution of benzaldehyde (3.10 g, 29.0 mmol) and zinc iodide (280 mg, 8.80 mmol) at 0 °C. The reaction mixture was warmed to room temperature over 18 h

then treated with saturated aqueous NaHCO $_3$  (ca. 100 mL). The layers were separated and the aqueous was extracted with EtOAc (2 x 75 mL). The combined organics were washed with saturated aqueous NaCl (1 x 50 mL), dried (Na $_2$ SO $_4$ ),

and concentrated under reduced pressure to give 3.78 g of 2-[(trimethylsilyl)oxy]-phenylacetonitrile as an oil, which was carried on without purification.

Lithium aluminum hydride (2.10 g, 55.0 mmol) was added in portions over 15 min to a solution of [2 -(trimethylsilyl)oxy]-phenylacetonitrile (3.75 g, 18.3 mmol) 10 in anhydrous THF (75 mL) at 0 °C. The reaction was quenched by the sequential addition of  $\rm H_2O$  (2.10 mL), 10% aqueous NaOH (2.10 mL), and  $H_2O$  (6.30 mL) then dried  $(Na_2SO_4)$  and filtered through a pad of Celite using EtOAc (ca. 75 mL). The filtrate was concentrated under reduced 15 pressure to provide an oil which was purified by flash chromatography, elution with 9:1 CH2Cl2-MeOH containing 2% Et<sub>3</sub>N, to give 2-hydroxy-1-phenethylamine (2.40 g) as an oil in 95% yield. ( ${}^{1}$ H NMR, 300 MHz) d 7.32 (comp, 5H), 4.66 (dd, 1H, J = 7.8, 4.0 Hz), 3.03 (dd, 1H, J = 12.5, 4.0 Hz).20 2.83 (dd, 1H, J = 12.5, 9.1 Hz), 2.46 (br s, 3H). LRMS 155  $(M+NH_4)$ , 138 (M+H).

Phosgene (6.0 mL of a 1.93 M solution in toluene Part B: 25 (PhCH<sub>3</sub>), 12.0 mmol) was added to a solution of 2-hydroxy-1phenethylamine (1.18 g, 8.60 mmol) in PhCH<sub>3</sub> (100 mL) at 0 °C followed by the dropwise addition of Et<sub>3</sub>N (1.80 mL, 13.0 mmol). The reaction mixture was warmed to room temperature over 48 h and poured into EtOAc (ca. 200 mL). The layers were separated and the aqueous was extracted with EtOAc (1 30 The combined organics were washed with saturated aqueous NaCl (1 x 100 mL), dried ( $Na_2SO_4$ ) and concentrated under reduced pressure to give 5-phenyl-2-oxazolidinone (1.05 g) as a solid in 75% yield. (1H NMR, 300 MHz) d 7.39 (comp, 5H), 5.69 (br s, 1H), 5.63 (dd, 1H, J = 8.4, 8.1) 35

Hz), 3.99 (dd, 1H, J = 8.4, 8.1 Hz), 3.55 (dd, 1H, J = 8.4, 8.1 Hz). LRMS 181 (M+NH<sub>4</sub>), 164 (M+H).

A solution of 5-phenyl-2-oxazolidinone (500 mg, Part C: 3.1 mmol) in anhydrous THF was added dropwise to a suspension of NaH (91 mg, 3.7 mmol) in anhydrous THF at 0 °C. The reaction mixture was warmed to room temperature over 30 min then heated at reflux for 15 min. Methyl bromoacetate (0.32 mL, 3.4 mmol) was added and the mixtue was heated at reflux for 2h. The reaction mixture was 10 cooled to room temperature and quenched with  ${\rm H}_2{\rm O}$  (ca. 20 mL). The aqueous was extracted with EtOAc (2  $\times$  75 mL). The combined organics were washed with saturated aqueous NaCl (1  $\times$  50 mL), dried (MgSO<sub>4</sub>), and concentrated under reduced pressure to give an oil which was purified by flash 15 chromatography, elution with 3:1 EtOAc-hexanes, to provide 2-[3-(5-phenyl-2-oxazolidino)]-acetic acid, methyl ester (545 mg) as an oil in 76% yield. ( $^{1}H$  NMR, 300 MHz) d 7.42 $^{\circ}$ (comp, 5H), 5.56 (dd, 1H, J = 8.4, 8.1 Hz), 4.10 (d, 2H, J)= 3.0 Hz), 4.06 (dd, 1H, J = 8.4, 8.1 Hz), 3.78 (s, 3H), 20 3.64 (dd, 1H, J = 8.4, 8.1 Hz). LRMS 253 (M+NH<sub>4</sub>, base), 236 (M+H).

Part D: A solution of 2-[3-(5-phenyl-2-oxazolidino)] acetic acid, methyl ester (540 mg, 2.30 mmol) in MeOH (10 mL) and H<sub>2</sub>O (10 mL) was treated with NaOH (138 mg, 3.40 mmol) and heated at reflux for 15 min. The reaction mixture was cooled to room temperature, acidified to pH 2 with 2M aqueous HCl, and extracted with EtOAc (3 x 50 mL). The combined organics were washed with saturated aqueous NaCl (1 x 25 mL), dried (MgSO<sub>4</sub>), and concentrated to give 2-[3-(5-Phenyl-2-oxazolidino)] acetic Acid (505 mg) as an oil in 99% yield. (1H NMR, 300 MHz) d 7.41 (comp, 5H), 5.57 (dd, 1H, J = 8.0, 8.0 Hz), 4.15 (s, 2H), 4.05

(dd, 1H, J = 8.0, 8.0 Hz), 3.65 (dd, 1H, J = 8.0, 8.0 Hz). LRMS 239 (M+NH<sub>4</sub>, base), 222 (M+H).

- Part E: Triethylamine (0.23 mL, 1.60 mmol) was added to a mixture of (2R)-4-bromo-1-aminobutane-1-boronic acid, (+)-pinanediol ester hydrochloride (540 mg, 1.47 mmol), HOBT (200 mg, 1.47 mmol), DCC (300 mg, 1.47 mmol), and 2-[3-(5-phenyl-2-oxazolidino)]acetic acid (325 mg, 1.47 mmol) in anhydrous THF (12 mL) and anhydrous DMF (3 mL) at 0 °C.
- The reaction mixture was warmed to room temperature over 18 h, diluted with  $Et_2O$  (ca. 30 mL), and filtered through Celite with additional  $Et_2O$  (ca. 50 mL). The filtrate was washed with  $H_2O$  (3 x 25 mL), saturated aqueous NaCl (1 x 25 mL), dried ( $Na_2SO_4$ ), and concentrated under reduced
- pressure to give (2R)-2-[[3-(5-phenyl-2-oxazolidino)]-acetamido]-4-bromo-1-aminobutane-1-boronic acid, (+)-pinanediol ester (770 mg) as a foam in 98% yield. LRMS 535, 533 (M+H) 453 (base).
- 20 Part F: A solution of (2R)-2-[[3-(5-phenyl-2-oxazolidino)]-acetamido]-4-bromo-1-aminobutane-1-boronic acid, (+)-pinanediol ester (770 mg, 1.45 mmol) and thiourea (220 mg, 2.90 mmol) in EtOH (15 mL) was heated at reflux for 36 hours then cooled to room temperature and diluted with Et<sub>2</sub>O (ca. 100 mL) which was decanted. The residue was purified by size exclusion chromatography on Sephadex LH -
  - 20, elution with MeOH, to give a foam. The foam was dissolved in 5 mL of anhydrous THF and treated with  $\rm Et_2O$  (ca. 30 mL) to give a solid that was washed with  $\rm Et_2O$  (ca.
- 30 10 mL) and dried to afford the title compound (180 mg) as a white powder in 21% yield, mp 93 96 °C. LRMS 529 (M+1, base); HRMS Calcd for  $C_{26}H_{38}BN_4O_5S$ : 529.2656. Found: 529.2644.

# Example 35.6.2

5

15

A solution of (2R)-2-[[3-(5-phenethyl-2-oxazolidino)]acetamido]-4-bromo-1-aminobutane-1-boronic acid, (+)pinanediol ester (710 mg, 1.27 mmol) was prepared by an analogous method to that reported for Example 35.5.3, however substituting hydrocinnamaldehyde for benzaldehyde. 10 Further reaction with thiourea (190 mg, 2.50 mmol) in EtOH (13 mL) was heated at reflux for 36 h then cooled to room temperature and diluted with  $Et_2O$  (ca. 200 mL) which was decanted. The residue was purified by size exclusion chromatography on Sephadex LH - 20, elution with MeOH, to give a foam. The foam was dissolved in 5 mL of anhydrous THF and treated with  $Et_2O$  (ca. 40 mL) to give a solid that was washed with  $\mathrm{Et_2O}$  (ca. 10 mL) and dried to afford the title compound (150 mg) as a white powder in 19% yield, mp 93 - 96 °C. LRMS 557 (M+H, base); HRMS Calcd for 20  $C_{28}H_{42}BN_4O_5S$ : 557.2969. Found: 557.2965.

# Example 59

A mixture of 1,8-napthalic anhydride (1.0 g, 5.1 25 Part A: mmol), glycine (0.42g, 5.6 mmol) and camphorsulfonic acid (ca. 100 mg) was suspended in absolute EtOH (60 mL) and DMF (20 mL). The reaction mixture was heated at reflux for 72 h, then the solvent removed by distillation in vacuo. The residue was diluted with  ${\rm H}_2{\rm O}$  (10 mL), acidified with HCl (1N) to pH = 3 and the resulting solid was isolated by filtration and air dried. There was obtained N, N-(1, 8napthyldiimido)-glycine (1.22 g) in 95% yield. LRMS:  $(M+H)^+ = 256$ ; mp 278-279 °C.

An alternative to the above prepartion of N,N-(1,8-napthyldiimido)glycine would be to react the sodium salt of 1,8-napthalic phthalimide with ethyl bromoacetate in dimethylformamide at 60°C. The resulting ester can then be hydrolyzed with 1N sodium hydroxide in ethanol solution to give the title compound.

N, N-(1, 8-Napthyldiimido) glycine (0.56 g, 2.1) mmol) and N-methylmorpholine (0.53 mL, 4.82 mmol) were 10 dissolved in THF (10 mL) and DMF (1 mL) then cooled to -20 OC. Isobutylchloroformate (0.31 mL, 2.32 mmol) was added to the cold solution and the reaction was stirred at -20 °C for 20 min. After this time a THF suspension (5 mL) of pinanediol 1-amino-4-bromobutane-boronate hydrochloride 15 salt (0.80 g, 2.19 mmol) was added to the mixture and the reaction was allowed to warm to room temperature over 3 h. The reaction was partitioned between H2O (10 mL) and EtOAc (15 mL). The organic layer was washed with  $H_2O$  (3 x 15 mL), then with saturated NaHCO<sub>3</sub> (15 mL), and brine (15 mL). After the solution was dried (MgSO<sub>4</sub>) and evaporated under reduced pressure, there was obtained pinanediol N-[N,N-(1,8-napthyl-diimido)glycyl]-1-amido-4-bromobutaneboronate (1.2 g) in 98% yield. LRMS:  $(M+H)^+ = 568$ .

25 Part C: To a solution of pinanediol N-[N,N-(1,8napthyldiimido)glycyl]-1-amido-4-bromobutaneboronate (1.0 g, 1.76 mmol) in MeOH (30 mL) was added thiourea (0.27 g, 3.53 mmol). The reaction mixture was heated at reflux for 4h, then was allowed to cool to ambient temperature and the 30 solvent was removed by distillation. The resulting viscous liquid was dissolved in a minimal amount of MeOH and passed through a short column (35 g, LH-20 Sephadex) by elution with MeOH. Product containing fractions were combined and concentrated in vacuo, then the resulting foam 35 was dissolved in a minimal amount of MeOH and triturated

with Et<sub>2</sub>O. After solvent was decanted, the residue was rinsed with additional Et<sub>2</sub>O and placed under vacuum. There was obtained pinanediol  $N-[N,N-(1,8-napthyldiimido)glycyl]-1-amido-4-S-thiourylbutane-boronate hydrogen bromide (1.0 g) as an amorphous foam in 100% yield. LRMS: <math>(M+H)^+ = 563$ .

- Part D: To a solution of pinanediol N-[N,N-(1,8-napthyldimido)glycyl]-1-amido-4-bromobutaneboronate (1.21g, 2.13 mmol) in DMF (10 mL) was added NaN3 (0.28 g, 4.27 mmol). The reaction mixture was heated at 65 °C for 8 h, then it was allowed to cool to room temperature and partitioned between H2O (15 mL) and EtOAc (20 mL). The layers were separated and the organic phase was washed with H2O (3x20 mL) and brine (20 mL). This solution was dried (MgSO4) and concentrated in vacuo to give pinanediol N-[N,N-(1,8-napthyldimido)glycyl]-1-amido-4-azidobutaneboronate (0.92 g) in 82% yield. LRMS: (M+H)+ = 530.
- 20 Part D: A suspension of the azide prepared above (1.19g, 2.25 mmol) and 10% Pd/C (100 mg) in MeOH (15 mL) was placed under an atmosphere of H<sub>2</sub> (1 atm). The reaction mixture was stirred at room temperature for 5 h, then purged with a stream of N<sub>2</sub>. The catalyst was removed by filtration
  25 through a pad of diatomaceous earth and the filtrate was concentrated under reduced pressure to give pinanediol N-[N,N-(1,8-napthyldiimido)glycyl]-1-amido-4-amino-butaneboronate (1.1 g) in 85% yield. LRMS: (M+H) + = 504.
- 30 Part E: To a solution of pinanediol N-[N,N-(1,8napthyldiimido)glycyl]-1-amido-4-aminobutaneboronate (0.51
  g, 1.01 mmol) in pyridine (10 mL) was added
  aminoiminomethanesulfonic acid (0.13 g, 1.01 mmol). The
  reaction mixture was heated at reflux for 4 h, then was
  35 concentrated in vacuo to give pinanediol N-[N,N-(1,8-

napthyldiimido)glycyl]-1-amido-4-guanidinobutane-boronate sulfonic acid salt (0.21 g) in 35% yield.

Part F: Pinanediol N-(N, N-(1, 8-napthyldiimido)glycyl]-1-5 amido-4-guanidinobutane-boronate sulfonic acid salt (0.21 g, 0.35 mmol) was dissolved in MeOH (2 mL) and Et<sub>2</sub>O (10 mL). A single portion of phenylboric acid (0.21g, 1.76 mmol) was added to the solution followed by H2O (10 mL) and this mixture was stirred for 15 h. The phases were 10 separated and the aqueous layer was washed with  $Et_2O$  (6 x 10 mL). The aqueous layer was concentrated in vacuo and the resulting residue was placed under vacuum to give N-[N, N-(1,8-napthyldiimido)-glycyl]-1-amido-4guanidinobutane-boronic acid as the sulfonic acid salt 15 (0.16 g) in quantitative yield.

The following compounds were prepared according to the methods outlined in the Synthesis and Experimental sections. Appropriate physical data to characterize the compounds are provided:

#### Example 52.1.2

25

20

11

Part A. Preparation of 3-methylphenethyl bromide.

To a solution of 3-methylphenethyl alcohol (5.0 g, 36.7 mmol) in methylene chloride at 0° C was added triphenylphosphine (10.6 g, 40.4 mmol) and carbon tetrabromide (13.4 g, 40.4 mmol). The mixture was allowed to stir with warming to 25° C for 16 h. The solvent was removed in vacuo and the residue was taken up in ether and filtered through a pad of silica gel. The solvent was removed in vacuo to afford 7.0 g (95%) of the title bromide.

Part B. Preparation of N-(3-methylphenethyl)-Gly-OMe.

To a solution of 3-methylphenethyl bromide (7.0 g, 35.0 mmol) in acetonitrile was added glycine methyl ester hydrochloride (6.6 g, 52.5 mmol) and sodium bicarbonate (10.3 g, 122.5 mmol). The resulting mixture was allowed to stir at 80°C for 16 h. The reaction mixture was allowed to cool to 25°C and then was diluted with ethyl acetate. The mixture was washed with water and brine, dried (MgSO4), and concentrated to afford the title compound. MS (CI): m/z 208 (M+H)+. Part C. Preparation of hydrocinnamoyl-[N-(3-methylphenethyl)-Gly]-OMe.

To a solution of N-(3-methylphenethyl)-Gly-OMe (3.15 g, 15.2 mmol) in THF at 0°C was added N-methylmorpholine (3.34 mL, 30.4 mmol) and hydrocinnamoyl chloride (2.26 mL, 15.2 mmol). The mixture was allowed to stir with warming to 25°C for 6 h. The solvent was removed in vacuo and the residue was taken up in ethyl acetate and washed with 10% aq HCl, sat'd aq NaHCO3 and brine. The organic layer was dried (MgSO4) and concentrated to afford 4.4 g (86%) of the title compound. MS (CI): m/z 340 (M+H)+.

Part D. Preparation of Hydrocinnamoyl-[N-25 (3-methylphenethyl)-Gly]-OH.

To a solution of hydrocinnamoyl-[N-(3-methylphenethyl)-Gly]-OMe (4.4 g, 13.0 mmol) in 50 mL of 2:1 THF/H<sub>2</sub>O was

added lithium hydroxide monohydrate (0.65 g, 15.6 mmol). The reaction mixture was allowed to stir at 25° C for 4 h and then the THF was removed in vacuo. The basic solution was extracted with 1:1 hexanes/ethyl acetate and the organic layer was discarded. The aqueous layer was

- 5

acidified with concentrated HCl and then was extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried (MgSO4) and concentrated to afford the title compound as a white solid. MS (CI): m/z 326 (M+H)+.

5

Part E. Preparation of Hydrocinnamoyl-[N-(3-methyl)henethyl)-Gly]-boroLys-OH hydrochloride salt.

10

The carboxylic acid hydrocinnamoyl-[N-(3-methylphenethyl)-Gly]-OH was elaborated to the title compound according to the procedures described in Example ???. MS (ES): m/z 454.4 (M+H)+.

15

25

30

#### Example 53.1.1

Hydrocinnamoyl-[N-(2,2-dimethyl)-phenethyl-Gly]boroLys-OH hydrochloride salt

20 Part A. Preparation of Methyl 2,2-(dimethyl)-2phenylacetate.

To a cooled (-78° C) solution of methyl phenylacetate (1.0 g, 6.7 mmol) in THF was added methyl iodide (0.91 mL, 14.7 mmol) followed by potassium tert-butoxide (14.7 mL of a 1.0 M solution in THF, 14.7 mmol). The reaction mixture was allowed to stir while slowly warming to 25° C. After 1 h the reaction was quenched by addition of saturated aq NH4Cl, diluted with ethyl acetate and washed with brine. The organics were dried (MgSO4) and concentrated to afford 1.1 g (92%) of the title compound.

Part B. Preparation of 2,2-(Dimethyl)-2-phenylethyl alcohol.

To a cooled  $(0^{\circ} C)$  solution of 1M lithium aluminum

hydride in ether (31.1 mL, 31.1 mmol) was added methyl 2,2-(dimethyl)-2-phenylacetate (5.54 g, 31.1 mmol) as a solution in ether. The reaction mixture was allowed to warm to 25° C and was stirred for 3 h. The mixture was cooled to 0° C and was quenched by slow sequential addition of 1.2 mL of water, 1.2 mL of 15% aq NaOH and 3.6 mL of water. The resulting slurry was stirred vigorously with warming to 25° C for 1 h, and then was dried (MgSO4), filtered and concentrated to afford 3.8 g (81%) of the title compound.

Part C. Preparation of 2,2-(Dimethyl)-2-phenylacetaldehyde.

To a solution of 2,2-(dimethyl)-2-phenylethyl alcohol

(3.8 g, 25.1 mmol) in methylene chloride was added
pyridinium chlorochromate (16.2 g, 75.3 mmol) and the
resulting mixture was stirred vigorously at 25° C for 4h.
The mixture was filtered through a pad of layered silica
gel (bottom)/Celite/Florasil (top) and concentrated to the
give the title aldehyde.

Part D. Preparation of N-(2,2-dimethyl)-phenethyl-Gly-OEt.

To a solution of glycine ethyl ester hydrochloride
(1.7 g, 12.3 mmol) in methanol was added sodium

25 cyanoborohydride (0.77 g, 12.3 mmol) and 2,2-(dimethyl)-2phenylacetaldehyde (2.0 g, 13.5 mmol). Glacial acetic acid
was added if necessary to maintain the pH at 5-6. The
mixture was allowed to stir at 25° C for 16 h. The
reaction was quenched by addition of excess satd. aq. K2CO3

30 and then was diluted with ethyl acetate. The layers were
separated and the organic layer was washed with brine (2x),
dried (MgSO4) and concentrated to afford 2.5 g (86%) of the
title compound. MS (CI): m/z 236 (M+H)+.

5

Part E. Preparation of Hydrocinnamoyl-N-(2,2-dimethyl)-phenethyl-Gly-OEt.

To a cooled (0° C) solution of N-(2,2-dimethyl)phenethyl-Gly-OEt (2.4 g, 10.1 mmol) in THF was added Nmethylmorpholine (2.22 mL, 20.2 mmol) followed by
hydrocinnamoyl chloride (1.50 mL, 10.1 mmol). The
resulting solution was allowed to warm to 25° C and was
stirred for 3h. The THF was removed in vacuo and the
residue was taken up in ethyl acetate and washed with 10%
aq HCl, satd. aq. NaHCO3 and brine. The organics were
dried (MgSO4) and concentrated. The residue was purified
by silica gel flash chromatography (solvent gradient 7:1
hexanes/ethyl acetate to 3:1 hexanes/ethyl acetate) to
afford 2.0 g (54%) of the title compound. MS (CI): m/z

368 (M+H)+.

Part F. Preparation of Hydrocinnamoyl-N-(2,2-dimethyl)-phenethyl-Gly-OH.

phenethyl-Gly-OEt (1.5 g, 4.1 mmol) in 25 mL of 1:1 methanol/water was added potassium hydroxide (0.34 g, 6.1 mmol). The reaction mixture was allowed to stir at reflux for 1 h and then was allowed to cool to 25° C and the methanol was removed in vacuo. The basic aqueous solution was extracted with 1:1 hexanes/ethyl acetate and the organic layer was discarded. The aqueous layer was acidified with concentrated HCl and then was extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried (MgSO4) and concentrated to afford the title compound as a white solid. MS (CI): m/z 340 (M+H)+.

Part G. Preparation of Hydrocinnamoyl-[N-(2,2-dimethyl)-phenethyl-Gly]-boroLys-OH hydrochloride salt.

The carboxylic acid hydrocinnamoyl-N-(2,2-dimethyl)35 phenethyl-Gly-OH was elaborated to the title compound

according to the procedures described in Example ???. MS (ES): m/z 468.4 (M+H)+.

#### Example 1.1.3

 $N-(N-methyl-N-[(3-phenyl)propionyl]glycyl}-1-amido-4-(guanidino)butylboronate, hydrochloride salt; LRMS (M+H, ethylene glycol ester)+ = 404.$ 

#### Example 6.1.1

N-{N-methyl-N-benzoylglycyl}-1-amido-5aminopentaneboronate, hydrochloride salt; HRMS calcd: (M+H, ethylene glycol ester)+ = 348.209462, obs.: (M+H, ethylene glycol ester)+ = 348.208418.

# Example 6.1.2

 $N-(N-methyl-N-\{phenylacetyl\}glycyl\}-1-amido-5-$  aminopentaneboronic acid, hydrochloride salt; LRMS (M+H, ethylene glycol ester) = 362.

#### 20 <u>Example 6.1.3</u>

 $N-\{N-\text{methyl-}N-[(3-\text{phenyl})\text{propionyl}]\text{glycyl}\}-1-\text{amido-}5-$ aminopentaneboronate; HRMS (M+H)+ calcd: 376.240762, found: 376.240727.

### 25 Example 9.1.2

 $N-\{N-\text{phenyl-}N-\{\text{phenylacetyl}\}\text{glycyl}\}-1-\text{amido-}5$ aminopentaneboronate, hydrochloride salt; HRMS calcd: (M+H, ethylene glycol ester)+ = 424.240762, obs.: (M+H, ethylene glycol ester)+ = 424.242097.

### Example 9.1.3

 $N-\{N-\text{phenyl}-N-[(3-\text{phenyl})\text{propionyl}\}\text{glycyl}\}-1-\text{amido}-5-\text{aminopentaneboronate}$ , hydrochloride salt; HRMS calcd: (M+H, ethylene glycol ester)+ = 438.256412, obs.: (M+H, ethylene glycol ester)+ = 438.256557.

30

35

5

#### Example 15.1.3

Pinanediol N-{N-methyl-N-[(3-phenyl)propionyl]glycyl}1-amido-4-(guanidino)butylboronate, hydrochloride salt;
LRMS (M+H)+ = 512.3.

#### Example 21.1.1

Pinanediol  $N-\{N-\text{methyl}-N-\text{benzoylglycyl}\}-1-\text{amido}-5$ aminopentaneboronate, hydrochloride salt; HRMS calcd: 10  $(M+H)^+ = 456.303362$ , obs.:  $(M+H)^+ = 456.302964$ .

#### Example 21.1.2

Pinanediol  $N-\{N-\text{methyl-}N-\{\text{phenylacetyl}\}\text{glycyl}\}-1$ amido-5-aminopentaneboronate, hydrochloride salt; LRMS 15  $(M+H)^+ = 470$ .

#### Example 21.9.1

Pinanediol N-{N-methyl-N-[2-(phenyl)benzoyl]glycyl}-1amido-5-amino-pentaneboronate, hydrochloride acid salt;
20 LRMS (M+H)+ = 532.3.

#### Example 24.1.2

Pinanediol  $N-\{N-\text{phenyl}-N-\{\text{phenylacetyl}\}\text{glycyl}\}-1$ amido-5-aminopentaneboronate, hydrochloride salt; HRMS 25 calcd:  $(M+H)^+ = 532.334663$ , obs.:  $(M+H)^+ = 532.334090$ .

### Example 24.1.3

Pinanediol  $N-\{N-\text{phenyl}-N-\{(3-\text{phenyl})\text{propionyl}\}\text{glycyl}\}-1-\text{amido}-5-\text{aminopentaneboronate}$ , hydrochloride salt; HRMS calcd:  $(M+H)^+=546.350313$ , obs:  $(M+H)^+=546.352069$ .

### Example 24.59.1

Pinanediol  $N-\{N-\text{phenyl}-N-\{N'-\text{methyl}-N'-\text{methyl}\}$  aminocarbonyl]glycyl}-1-amido-4-

isothiouroniumbutylboronate; HRMS (M+H) + calcd: 606.338533, found: 606.329421.

#### Example 26.1.3

Pinanediol  $N-\{N-\text{methyl}-N-\{(3-\text{phenyl})\text{propionyl}\}\text{glycyl}\}-1-\text{amido}-4-\text{isothiouroniumbutylboronate}; HRMS (M+H)+ calcd: 529.301983, found: 529.302078.$ 

# Example 26.9.1

Pinanediol N-{N-methyl-N-[(2-phenyl)benzoyl]glycyl}-1-amido-4-isothiouroniumbutylboronate; HRMS (M+H)+calcd: 577.301983, found: 577.302704.

### Example 26.9.3

pinanediol N-{N-methyl-N-[(3-(2phenyl)phenyl)propionyl]glycyl}-1-amido-4isothiouroniumbutylboronate; HRMS (M+H)+ calcd: 605.3333,
found: 605.3325.

20 <u>Example 26.12.1</u>

Pinanediol  $N-\{N-\text{methyl}-N-[(3-(2-\text{phenyl})\text{phenyl})\text{propionyl}]\text{glycyl}\}-1-\text{amido}-4-$  isothiouroniumbutylboronate; HRMS  $(M+H)^+$  calcd: 606.3285, found: 606.3294.

25

30

# Example 27.1.3

Pinanediol  $N-\{N-[(4-hydroxyphenyl)methyl]-N-[(3-phenyl)propionyl]-glycyl\}-1-amido-4-isothiouroniumbutylboronate; HRMS <math>(M+H)^+$  calcd: 621.328198, found: 621.329437.

# Example 28.1.3

Pinanediol  $N-\{N-\{(2-phenyl)ethy1\}-N-\{(3-phenyl)propionyl\}glycyl\}-1-amido-4-$ 

isothiouroniumbutylboronate; HRMS (M+H) + calcd: 619.348934, found: 619.348587.

#### Example 29.1.3

Pinanediol  $N-\{N-\text{phenyl}-N-\{(3-\text{phenyl})\text{propionyl}\}glycyl}-1-\text{amido}-4-\text{isothiouroniumbutylboronate}; HRMS (M+H)+ calcd: 591.317633, found: 591.316620.$ 

#### Example 30.1.3

Pinanediol N-{N-[(naphth-2-yl)methyl]-N-[(3-phenyl)propionyl]glycyl)-1-amido-4-isothiouroniumbutylboronate; HRMS (M+H)+ calcd: 655.348934, found: 655.347870.

# 15 <u>Example 36.6.1</u>

Pinanediol  $N-\{2-[(2-\infty x_0-4-\text{methylphenyl})-4,5-(H) \text{ oxazol-}3-yl]$  acetyl $\}-1-\text{amido-}4-\text{isothiouroniumbutylboronate}$ ; mp. 120-130 °C; Anal calcd. for C27H39BN4O5S HBr %C: 52.02; %H: 6.47; %N: 8.99; %B: 1.73; found: %C: 52.01; %H: 6.43,; %N: 8.84; %B: 1.75.

# Example 49.1.1

Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn-C10H16 HCl; MS (ESI) (M+H)+ 560.4.

25

20

#### Example 49.1.2

 $\label{eq:hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn(CH=NH)-C10H16} $$HCl; MS (NH_3-CI) (M+H) + 587.7.$ 

30

## Example 49.1.3

Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn(CH=NH)-OH HCl;
MS (ESI) (M+H)+ 453.1.

#### Example 49.2.1

Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroArg(CH3)-C10H16 HC1;
MS (ESI) (M+H)+ 616.4.

#### Example 50.1.1

5 Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroOrn-C10H16 HCl; MS (ESI) (M+H) + 499.2

#### Example 50.1.2

Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroOrn(CH=NH)-C10H16 HCl; 10 MS (ESI)  $(M+H)^+$  526.1

# Example 50.1.3

Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroLys-C10H16 HCl; MS  $(NH_3-CI)$  (M+H)+ 513.5.

15

### Example 51.1.1

 $\label{eq:hydrocinnamoy1-[N-(3-(Trifluoromethyl)-Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H)^+ 642.5}$ 

20

# Example 51.1.2

Hydrocinnamoyl-[N-(3-(Methyl)-Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H)+ 588.4

# Example 52.1.1

25 Hydrocinnamoyl-[N-(3-(Trifluoromethyl)-Phenethyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H)+ 534 for ethylene glycol ester.

# **EXAMPLE 61.1.1**

30 Hydrocinnamoyl-Sar-Lys[C(O)CO<sub>2</sub>H]

Part A: Preparation of Na-t-Boc-Ne-Cbz-lysine[N(OMe)Me]
A flask was charged 150 ml of anhydrous CH<sub>2</sub>Cl<sub>2</sub>
followed by the addition of Na-t-boc-NE-Cbz-lysine (15.00 grams, 39.43 mmol), N-methylmorpholine (13.0 ml, 118.29

mmol) and was cooled to -78°c followed by the addition of isobutylchloroformate (5.11 ml, 39.43 mmol). The mixture was stirred for 1 hr at which time the N,O-Dimethylhydroxylamine hydrochloride was added, stirred for 1 hr. and allowed to warm to room temperature. The solvent was removed in vacuo, the residue diluted with EtOAc and washed with 10% aq HCl, sat'd aq NaHCO3 and brine. After drying (MgSO4), the solution was filtered through a pad of silica gel and the solvent removed in vacuo to give the product (16.32 g). MS: ESI, m/z 424.2 (M+H)+.

Part B: Preparation of Part A: Na-t-boc-Ne-Cbz-lysinal A flask was charged with 200 ml of anhydrous THF followed by the addition of Na-t-boc, NE-Cbz 15 lysine[N(OMe)Me] (4.00 gr, 9.44 mmol), cooled to 0°c and followed by the addition of Lithium aluminum hydride (450 mg, 11.80 mmol). The solution was stirred for 30 min. and was slowly quenched with a sat'd  $KHSO_3$  solution (2.25 gr, 16.53 mmol). The volatiles were removed in vacuo and the residue dissolved in EtOAc and washed with 10% aq HCl, 20 sat'd aq  $NaHCO_3$  and brine. After drying  $(MgSO_4)$ , the solution was filtered through a pad of silica gel and the solvent removed in vacuo to give the product (3.29 grams) MS: ESI, m/z 365.2 (M+H)+.

25

10

Part C: Preparation of Na-t-boc, Ne-Cbz lysine[C(OH)CO<sub>2</sub>CH<sub>3</sub>]

A flask was charged with 150 ml of anhydrous THF followed by the addition of orthoethylthioformate (6.82 gr, 34.77 mmol), cooled to -78°c and lithiated with n-butyllithium (2.5 M, 14.0 ml, 34.7 mmol). After stirring for 20 min., Na-t-boc-NE-Cbz-lysinal was added as a THF solution via cannula and continued to stir at -78°c for and additional 4 hrs; the solution was quenched using sat'd NH4Cl and the volatiles removed in vacuo. The residue was

dissolved in EtOAc and washed twice with brine, dried over MgSO<sub>4</sub>, filtered and dried in vacuo. The residue was dissolved in 95% MeOH followed by the addition of HgO (12.57 gr, 58.06 mmol) and HgCl<sub>2</sub> (40.8 gr, 150.2 mmol) and stirred at rt. for 3 hrs. The solution was filtered through a pad of celite and the volatiles removed in vacuo followed by the addition of chloroform, the solution filtered and the volatiles removed in vacuo. The residue was dissolved in EtOAc and washed with 10% aq HCl, sat'd aq NaHCO<sub>3</sub> and brine. After drying (MgSO<sub>4</sub>), the solution was filtered through a pad of silica gel and the solvent removed in vacuo. The residue was purified by flash chromatography to give the product (1.49 grams) MS: CI m/z 425.2 (M+H)+

15

10

Part D: Preparation of NE-Cbz-lysine[C(OH)CO<sub>2</sub>CH<sub>3</sub>] TFA salt Na-t-boc-Ne-Cbz-lysine[C(OH)CO<sub>2</sub>CH<sub>3</sub>] (1.49 gr, 3.51 mmol) was dissolved in 50 ml of neat TFA and the reaction monitored by TLC. The volatiles were removed in vacuo and the residue dissolved in a minimum amount of CH<sub>2</sub>Cl<sub>2</sub> followed by the addition of Et<sub>2</sub>O, cooled to -78°c and the product ppt. out with hexane (1.36 gr). MS: CI m/z 325.0 (M+H)+.

Part E: Preparation of Hydrocinnamoyl-Sar-ethyl ester
Hydrocinnamic acid was added to 100 ml of anhydrous
CH<sub>2</sub>Cl<sub>2</sub> followed by the addition of N-methylmorpholine (44.0
ml, 400.0 ml), and cooled to -78°c. To the resulting
solution was added isobutylchloroformate (17.3 ml, 133.17
mmol) and stirred for 1 hr. The sarcosine ethyl ester
hydrochloride (20.00 gr, 133.17 mmol) was added and the
solution stirred for and additional hr at -78°c and allowed
to warm to rt. The volatiles were removed in vacuo and the
residue was dissolved in EtOAc and washed with 10% aq HCl,
sat'd aq NaHCO<sub>3</sub> and brine. After drying (MgSO<sub>4</sub>), the

solution was filtered through a pad of silica gel and the solvent removed in vacuo (31.56 gr) MS: CI m/z 250.0 (M+H)+.

Part F: Preparation of Hydrocinnamoyl-Sar-OH
Hydrocinnamoyl-Sar-ethyl ester (31.57 gr, 126.63 mmol)
and KOH (21.32 gr, 380 mmol) were combined in a 50/50
methanol-water solution and stirred at rt, the reaction was
monitored by TLC. The methanol was removed in vacuo and
the organics were extracted using EtOAc. The aq extract
was acidified with a 10% HCl solution and the organics
extracted with EtOAc and washed with brine. After drying
(MgSO<sub>4</sub>), the solution was filtered and the solvent removed
in vacuo to a white solid. MS: CI m/z 222.0 (M+H)+.

15

30

35

Part G: Preparation of Hydrocinnamoyl-Sar-NE-Cbz-lysine[C(OH)CO<sub>2</sub>CH<sub>3</sub>]

Hydrocinnamoyl-Sar-OH (690 mg, 3.10 mmol), NE-Cbz lysine[C(OH)CO<sub>2</sub>CH<sub>3</sub>] TFA salt (1.36 gr, 3.10 mmol), N-Methyl morpholine (1.0 ml, 9.3 mmol), HOBT (420 mg, 3.10 mmol) and 1-(3-Dimethyl amino propyl)-3 ethyl carbodiimide (600 mg, 3.10 mmol) were dissolved in 50 ml of anhydrous DMF and stirred overnight at rt. The resulting solution was diluted with 300 ml of EtOAc and washed repeatedly with brine. The organic were dried over MgSO<sub>4</sub>, filtered through a pad of silica gel, and the volatiles dried in vacuo to an oil (1.08 gr). MS: CI m/z 528.4 (M+H)+.

Part H: Preparation of Hydrocinnamoyl-Sar-NE-Cbz-lysine[C(0)CO<sub>2</sub>CH<sub>3</sub>]

Anhydrous  $CH_2Cl_2$  (100 ml) was charged with oxalylchloride (20 ml 2.25 mmol) and cooled to -40°c; this was followed by the addition of anhydrous DMSO (.35 ml, 4.91 mmol) and stirred for 20 min. Hydrocinnamoyl-Sar-NE-Cbz-lysine[C(OH)CO<sub>2</sub>CH<sub>3</sub>] (1.08 gr, 2.04 mmol) was added as a

 ${\rm CH_2Cl_2}$  solution and stirred for an additional 20 minutes. Triethylamine (1.42 ml, 10.23 mmol) was added to the resulting solution and stirred for an additional 20 min. The volatiles were removed in vacuo and the resulting residue subject to flash chromatography yielding the product as an oil.(73 gr) MS: CI m/z 526.4 (M+H)+.

Part I: Preparation of Hydrocinnamoyl-Sar-NE-Cbz-lysine[C(0)CO<sub>2</sub>H]

Hydrocinnamoyl-Sar-NE-Cbz lysine[C(O)CO<sub>2</sub>CH<sub>3</sub>] (.73 gr, 1.39 mmol) and LiOH (150 mg, 3.48 mmol) were combined in a 50/50 mixture of methanol/water, stirred at rt and the reaction monitored by TLC. The volatiles were removed in vacuo, the residue dissolved in EtOAc and acidified with 10% HCl. The organics were dried (MgSO<sub>4</sub>) and the volatiles removed in vacuo to yield .57 gr of product MS: CI m/z 468.2 (M+H-CO<sub>2</sub>)+.

# Part J: Hydrocinnamoyl-Sar-Lys[C(0)CO2H]

Hydrocinnamoyl-Sar-Ne-Cbz-lysine[C(O)CO<sub>2</sub>H] (570 mg, 1.11 mmol) was dissolved in 100 ml of methanol followed by the addition of 20% Pd/C catalyst (60 mg) and stirred under 1 atm. of H<sub>2</sub> at rt for 3 hrs. The solution was filtered through a pad of celite and the volatiles removed in vacuo to give the product MS: CI m/z 380.3 (M+H)+

## Example 53.1.2

Hydrocinnamoyl-[N-(2-(Cyclopropyl)-Phenethyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H)+ 466.3

30

#### Example 53.2.1

Hydrocinnamoyl-[N-(N-(Methyl)-Phenyl)-Gly]-boroLys-OH HC1; MS (ESI)  $(M+H)^+$  441.3

35

Example 53.2.2

 $\label{eq:hydrocinnamoyl-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-OH HCl;} $$MS (ESI) (M+H)^+ 455.4$ 

## Example 53.4.3

5 Hydrocinnamoyl-[N-(Cyclohexyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H) + 418.3

## Example 54.1.1

Hydrocinnamoyl- $[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroLys-10 C10H16 HCl; MS (ESI) <math>(M+H)^+$  616.3

# Example 54.1.2

 $\label{eq:hydrocinnamoyl-[N-(2-(Cyclopropyl)-Phenethyl)-Gly]-boroLys-Cl0Hl6 HCl; MS (NH_3Cl) (M+H) + 600.5$ 

15

# Example 54.1.3

Hydrocinnamoyl-[N-(2,2-(Diethyl)-Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H) + 630

20

## Example 54.2.1

 $\label{eq:hydrocinnamoyl-[N-(N-(Methyl)-Phenyl)-Gly]-boroLys-C10H16} $$HCl; MS (NH_3CI) (M+H) + 575.4$ 

#### Example 54.2.2

25 Hydrocinnamoyl-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-C10H16 HCl; MS  $(NH_3CI)$  (M+H)+ 589.4

## Example 54.2.3

Hydrocinnamoyl-[N-(Succiny1)-Gly]-boroLys-C10H16; MS (ESI)
30 (M+H) + 542.5

## Example 54.3.1

Hydrocinnamoyl-[N-(Methyl Succinyl)-Gly]-boroLys-C10H16
HCl; MS (ESI) (M+H) + 556.5

#### Example 54.3.2

Hydrocinnamoyl-[N-(2-(Cyclopentyl)-Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H)+ 628.3

# 5 <u>Example 54.3.3</u>

Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H)+ 630.4

#### Example 54.4.1

10 Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-boroLys-C10H16 HCl; MS (ESI) (M+H)+ 602.4

#### Example 54.4.2

Hydrocinnamoy1-[N-(Cyclopropyl)-Gly}-boroLys-C10H16 HC1;
15 MS (NH<sub>3</sub>CI) (M+H) + 510.3

## Example 54.4.3

 $\label{eq:hydrocinnamoyl-[N-(Cyclohexyl)-Gly]-boroLys-C10H16 HCl; MS $$ (NH_3CI) (M+H)^+ 552.4 $$$ 

20

# Example 55.1.1

Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroArg-OH HCl; MS (ESI) (M+H)+ 496.4

25 Example 56.1.1

 $\label{eq:hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroArg-C10H16 HCl; MS (ESI) (M+H)^+ 630.4$ 

# Example 56.1.2

30 Hydrocinnamoyl-[N-(2-(Cyclopropyl)-Phenethyl)-Gly]-boroArg-C10H16 HCl; MS (ESI) (M+H)+ 628.3

# Example 56.3.3

 $\label{eq:hydrocinnamoyl-N-[2,2-(Dimethyl)-2-(3,5-dimethylphenyl)-35} \\ \text{ethyl]-Gly}-\text{boroArg-C10H16 HCl;} \\ \text{MS (ESI) } \\ (\text{M+H})^+ \\ \text{658.4} \\$ 

## Example 56.4.1

 $\label{eq:hydrocinnamoyl-N-[2-(3,5-dimethylphenyl)-ethyl]-Gly} + boroArg-C10H16 HCl; MS (ESI) (M+H) + 630,4$ 

5

#### Example 57.1.1

Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroOrn(CH=NH)-OH HCl; MS (ESI) (M+H)+ 481.2

10

25

35

#### Example 57.1.2

Hydrocinnamoyl-[N-(2-(Cyclopropyl)-Phenethyl)-Gly]-boroOrn(CH=NH)-OH HCl; MS (ESI) (M+H)+ 479

#### Example 57.4.2

Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn(CH=NH)-OH HCl;
MS (ESI) (M+H)+ 389.3

#### Example 58

N-{N-methyl-N-[2-(methylphenyl)benzoyl]glycyl}-1-amido-5-20 aminopentaneboronic acid, hydrochloride salt

Using the procedure of Example 8.1.3, however using pinanediol N-(N-methyl-N-[2-(methylphenyl)benzoyl] glycyl}-1-amido-5-aminopentaneboronate, hydrochloride salt, the title compound was prepared; LRMS  $(M-H_2O)^+=394.1$ ,  $(M-2H_2O)^+=376.1$ .

## Example 58.1.1

Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-30 boroOrn(CH=NH)-Cl0Hl6 HCl; MS (ESI) (M+H)+ 615.4

## Example 58.3.3

Hydrocinnamoyl-{N-[2,2-(Dimethyl)-2-(3,5-dimethylphenyl)-ethyl]-Gly}-boroOrn(CH=NH)-C10H16 HCl; MS (ESI) (M+H)+643.4

## Example 58.4.1

 $\label{eq:hydrocinnamoyl-N-[2-(3,5-dimethylphenyl)-ethyl]-Gly} - boroOrn(CH=NH)-Cl0H16 HCl; MS (ESI) (M+H)+ 615.4$ 

5

#### Example 58.4.2

 $\label{eq:hydrocinnamoy1-[N-(Cyclopropy1)-Gly]-boroOrn(CH=NH)-C10H16} $$ HCl; MS (NH_3CI) (M+H)^+ 524.3$ 

10

#### Example 59.1.1

Hydrocinnamoy1-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H)+ 482.2

## Example 59.4.2

Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl; MS
(ESI) (M+H) + 376.2

#### Example 60

Pinanediol N-{N-((naphth-1-yl)methyl)-N-((3-20 phenyl)propionyl)glycyl}-1-amido-4-isothiouroniumbutylboronate; HRMS (M+H)+calcd: 655.348934, found: 655.349243.

# Example 60.1.1

25 Hydrocinnamoyl- $[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroOrn-C10H16 HCl; MS <math>(NH_3DCI)$   $(M+H)^+$  602.5

# Example 60.3.3

Hydrocinnamoyl- $[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-boroOrn-30 C10H16 HCl; MS (ESI) <math>(M+H)^+$  616.4

# Example 60.4.1

 $\label{eq:hydrocinnamoyl-N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-boroOrn-C10H16 HCl; MS (ESI) (M+H)+ 588.3$ 

## Example 60.4.2

Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn-C10H16 HCl;
MS (NH<sub>3</sub>CI) (M+H) + 496.3

5

#### Example 61

Pinanediol N-{N-methyl-N-[(3-phenyl-2-(phenyl)methyl)propionyl}-glycyl}-1-amido-5-aminopentylboronate; HRMS (M+H)+ calcd: 560.365963, found: 560.364426.

10

#### Example 61.1.1

Hydrocinnamoy1-Sar-Lys[C(=0)-C(=0)-OH]; MS: ESI m/z 380.3 (M+H)+

15

#### EXAMPLE 62.3.3

2-(2-cyanothiophenyl)-benzoyl-Sar-BoroArg C<sub>10</sub>H<sub>16</sub>·HCl

- Part A: Preparation of 2-(2-cyanothiophenyl)benzoyl-Sarethyl ester
- 20 2-(2-cyanothiophenyl)benzoic acid (5.00 gr, 19.58 mmol), sarcosine ethyl ester hydrochloride (3.00 gr, 19.58 mmol), triethylamine (8.2 ml, 58.75 mmol), HOBT (2.64 gr, 19.58 mmol) and 1,3-diisopropylcarbodiimide (3.0 ml, 19.58 mmol) were combined in 100 ml of anhydrous CH<sub>2</sub>Cl<sub>2</sub> and 25 stirred at rt overnight. The velatiles were revered in
- stirred at rt overnight. The volatiles were removed in vacuo, the organics dried over MgSO<sub>4</sub>, filtered and concentrated. The product was purified by flash chromatography (4.93 gr) MS: CI m/z 355.1 (M+H)+.
- Part B: Preparation of 2-(2-cyanothiophenyl)benzoyl-Sar-OH
  2-(2-cyanothiophenyl)benzoyl-Sar-ethyl ester (4.93 gr,
  14.03 mmol) and KOH (790 mg, 14.03 mmol) were combined in
  75 ml of a 50/50 methanol-water solution and stirred at rt,
  the reaction was monitored by TLC. The methanol was
  removed in vacuo and the organics were diluted with EtOAc

and extracted with  $H_2O$ . The aq extract was acidified with a 10% HCl solution and the organics extracted with EtOAc, washed with brine. After drying (MgSO<sub>4</sub>), the solution was filtered and the solvent removed in vacuo to a white solid.

5 MS: CI m/z 327.0 (M+H) +

Part C: Preparation of 2-(2-cyanothiophenyl)benzoyl-Sar-NH-CH[( $CH_2$ )3Br]BO2- $C_{10}H_{16}$ 

2-(2-cyanothiophenyl)benzoyl-Sar-OH (4.23 gr, 12.96

10 mmol) and NMM (4.3 ml, 38.9 mmol) were dissolved in 150 ml of anhydrous CH<sub>2</sub>Cl<sub>2</sub> and stirred at -78°c. This was followed by the addition of isobutylchloroformate(1.70 ml, 12.96 mmol) and stirred for 1 hr, at which time NH<sub>2</sub>CH[(CH<sub>2</sub>)<sub>3</sub>Br]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub>· HCl (4.75 gr, 12.96 mmol) was added and stirred for an additional hr. and allowed to warm to rt. The volatiles were removed in vacuo, the residue was dissolved in EtOAc and the organics were washed with sat'd NaHCO<sub>3</sub>, 10% HCl and brine. The residue was dried (MgSO<sub>4</sub>) and filtered through a pad of florisil and concentrated in vacuo to give the product (6.01 gr) MS: CI m/z 558.2 (M-HBr)+

Part D: Preparation of 2-(2-cyanothiophenyl)benzoyl-Sar-NH-CH[(CH<sub>2</sub>)<sub>3</sub>N<sub>3</sub>]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub>

2-(2-cyanothiophenyl)benzoyl-Sar-NH-CH[(CH<sub>2</sub>)<sub>3</sub>Br]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub> (6.01 gr, 9.21 mmol) and NaN<sub>3</sub> (1.80 gr, 27.64 mmol) were combined in 75 ml of DMF and stirred for 3 hrs. at 110°c. The solution was diluted with EtOAc and washed repeatedly with brine. The organics were dried over MgSO<sub>4</sub> and filtered through a pad of florisil and concentrated in vacuo to give the product (5.38 gr).MS: ESI m/z 601.4 (M+H)+

Part E: Preparation of 2-(2-cyanothiophenyl)benzyloyl-Sar-35 boroOrn-C<sub>10</sub>H<sub>16</sub>·HCl To a solution of 2-(2-cyanothiophenyl)benzyloyl-Sar-NH-CH[(CH<sub>2</sub>)<sub>3</sub>N<sub>3</sub>]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub> (5.38 gr. 8.96 mmol) in MeOH (75 ml) was added 20% Pd/C catalyst (600 mg). The mixture was stirred under 1 atm of H<sub>2</sub> for 2 hrs and then filtered through a pad of celite and concentrated to give the product MS: CI m/z 575.2 (M+H)+.

Part F: Preparation of 2-(2-cyanothiophenyl)benzyloyl-SarboroArg-C<sub>10</sub>H<sub>16</sub>·HCl

#### Example 61,2.1

Hydrocinnamoyl-Sar-Lys-C(=0)-OCH3 HCl; MS: CI m/z 364.2 20  $(M+H)^+$ 

#### Example 61.2.3

Hydrocinnamoyl-Sar-Lys-C(=O)-CH3 HCl; MS: CI m/z 348.2 (M+H)+

25

## Example 61.4.1

Hydrocinnamoyl-Sar-Lys[CH(OH)(OCH3)-C(=0)-OCH3] HCl; MS: CI m/z 392.3(M+H)+

30

#### Example 62

Pinanediol N-{N-methyl-N-[(3,4-dichlorophenyl)acetyl]glycyl}-1-amido-5-aminopentaneboronate, hydrochloride salt; LRMS (M+H)+ = 538.

35

#### Example 62.1.3

DM-6666-C

(2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroIrg-C10H16 HBr; MS: DCI m/z 648.(M+H)+

## Example 62.2.2

5 (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-borohomolrg-C10H16 HBr: MS: ESI m/z 634.4 (M+H)+

# Example 62.3.2

(2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-borohomoArg-C10H16

10 HCl; MS: DCI m/z 631. (M+H)+

## EXAMPLE 64.2.3

2-(3-chlorobenzyl)-benzoyl-Sar-Borolys C10H16·HCl

- Part A: Preparation of 2-bromo-3'-chloro diphenyl methanol Anhydrous THF (400 ml) was charged with 3-(chloro) bromo benzene (19.63 ml, 167.13 mmol), cooled to -78°c, lithiated with n-BuLi (2.5 M, 66.85 ml, 167.13 mmol) and stirred for 20 minutes. 2-bromobenzaldehyde (19.51 ml, 167.13 mmol) was added and stirred for an additional 30 minutes and allowed to warm to rt. The solution was quenched with sat'd NH<sub>4</sub>Cl and the volatiles removed in vacuo. The residue was dissolved in EtOAc and washed with brine; the organics were dried over MgSO<sub>4</sub>, filtered through a pad of silica and the volatiles dried in vacuo. The product was purified by flash chromatography (38.8 gr) MS: CI m/z 281.0 (M+H-H<sub>2</sub>O)+
- Part B: Preparation of 2-bromo-3'-chloro diphenyl methane

  2-bromo-3'-chloro diphenyl methanol (38.8 gr, 130.35

  mmol) and triethylsilane (31.23 ml, 195.53 mmol) were

  combined in 200 ml of TFA and stirred overnight at rt. The

  volatiles were removed in vacuo and the residue purified by

  flash chromatography (33.32 grams) MS: CI m/z 283.0

  (M+H)+.

Part C: Preparation of 3-chlorobenzylbenzoic acid
2-bromo-3'-chloro diphenyl methane was dissolved in
250 ml of anhydrous THF, cooled to -78°c and lithiated with
n-BuLi (2.5 M, 47.5 ml, 118.34 mmol). After stirring for
30 minutes, CO<sub>2</sub> was slowly bubbled in for 15 minutes and
the solution warmed to rt. The volatiles were removed in
vacuo and the residue dissolved in H<sub>2</sub>O and the organics
removed with EtOAc, the aq. layer was acidified with 10%
HCl, the organics extracted with EtOAc, washed with brine,
dried (MgSO<sub>4)</sub> and the volatiles removed in vacuo to a white
solid MS: CI m/z 264.0 (M+H)+.

Part D: Preparation of 2-(3-chlorobenzyl)benzoyl-Sar ethyl
15 ester

3-chlorobenzylbenzoic acid (10.28 gr, 41.67 mmol), sarcosine ethyl ester hydrochloride (6.40 gr, 41.67 mmol), 1-(3-Dimethyl amino propyl)-3 ethyl carbodiimide (8.0 gr, 41.67 mmol), NMM (13.75 ml, 125.0 mmol) and DMAP (1.27 gr, 10.42 mmol) were combined in anhydrous CH<sub>2</sub>Cl<sub>2</sub> and stirred at rt overnight. The volatiles were removed in vacuo, the residue dissolved in EtOAc, washed with 10% aq HCl, sat'd aq NaHCO<sub>3</sub> and brine. After drying (MgSO<sub>4</sub>), the solution was filtered through a pad of silica gel and the solvent removed in vacuo to give the product (3.19 grams) MS: CI, m/z 346.0 (M+H)+.

Part E: Preparation of 2-(3-chlorobenzyl)benzoyl-Sar-OH 3-chlorobenzylbenzoyl-Sar ethyl ester (3.19 gr, 9.22 mmol) and KOH (2.0 gr, 36.90 mmol) were combined in 75 ml of a 50/50 methanol-water solution and stirred at rt, the reaction was monitored by TLC. The MeOH was removed in vacuo and the organics were diluted with EtOAc and extracted with  $\rm H_{2}O$ . The aq extract was acidified with a 35 10% HCl solution and the organics extracted with EtOAc and

washed with brine. After drying  $(MgSO_4)$ , the solution was filtered and the solvent removed in vacuo to a white solid. MS: CI m/z 318.0 (M+H)+

5 Part F: Preparation of 2-(3-chlorobenzyl)benzyloyl-Sar-NH-CH[(CH<sub>2</sub>)<sub>4</sub>Br]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub>

3-chlorobenzylbenzoyl-Sar-OH (2.66 gr, 8.37 mmol) and NMM (2.76 ml, 25.11 mmol) were dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> and stirred at -78°c. This was followed by the addition of isobutylchloroformate (1.1 ml, 8.37 ml) and stirred for 1 hr, at which time NH<sub>2</sub>CH[(CH<sub>2</sub>)<sub>4</sub>Br]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub>. HCl (3.19 gr, 8.37 mmol) was added and stirred for an additional hr. and allowed to warm to rt. The volatiles were removed in vacuo, the residue was dissolved in EtOAc and the organics were washed with sat'd NaHCO<sub>3</sub>, 10% HCl and brine. The residue was dried (MgSO<sub>4</sub>), filtered through a pad of florisil and concentrated in vacuo to give the product (.4.45 gr) MS: CI m/z 645.5 (M+H)+

20 Part G: Preparation of 2-(3-chlorobenzyl)benzyloyl-Sar-NH-CH[( $CH_2$ ) $_4N_3$ ]BO2- $C_{10}H_{16}$ 

3-chlorobenzylbenzyloyl-Sar-NH-CH[(CH<sub>2</sub>)<sub>4</sub>Br]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub> (4.45 gr, 6.91 mmol) and NaN<sub>3</sub> (1.34 gr, 20.73 mmol) were combined in 50 ml of DMF and stirred for 3 hrs. at  $110^{\circ}$ c. The solution was diluted with EtOAc and washed repeatedly with brine. The organics were dried over MgSO<sub>4</sub> and filtered through a pad of florisil and concentrated in vacuo to give the product (3.33 gr) MS: DCI m/z 623.0 (M+NH<sub>4</sub>)+

30

35

25

Part H: Preparation of 2-(3-chlorobenzyl)benzyloyl-Sarborolys-C10H16·HCl

To a solution of 3-chlorobenzylbenzyloyl-Sar-NH-CH[(CH<sub>2</sub>)<sub>4</sub>N<sub>3</sub>]BO<sub>2</sub>-C<sub>10</sub>H<sub>16</sub> (3.33 gr, 5.49 mmol) in MeOH (75 ml) was added 20% Pd/C catalyst (300 mg). The mixture was

stirred under 1 atm of  $H_2$  for 2 hrs and then filtered through a pad of celite and concentrated to give the product MS: CI m/z 580.5 (M+H)+.

5

#### Example 62.4.1

(2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroOrn(CH=NH)-C10H16 HCl; MS: ESI m/z 602.3 (M+H)+

#### Example 62.4.2

10 (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroLys(CH=NH)-C10H16 HCl; MS: ESI m/z 616.2 (M+H)+

#### Example 62.4.3

(2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroLys-C10H16 HCl;

MS: CI m/z 589.3 (M+H)+

#### Example 63

Pinanediol N-{N-methylphenyl-N-[(3phenyl)propionyl]glycyl}-1-amido-5-aminopentaneboronate,

20 hydrochloride salt; HRMS calcd: (M+H)+ = 560.365963, obs:
(M+H)+ = 560.366107.

## Example 63.1.3

2-(Thiophenyl)-Benzoyl-Sar-boroIrg-C10H16 HBr; MS: CI m/z 567.2 (M-H2NCN)+

# Example 63.3.1

2-(Thiophenyl)-Benzoyl-Sar-boroOrn-C10H16 HCl; MS: CI m/z 550.3 (M+H)+

30

#### Example 63.4.1

2-(Thiophenyl)-Benzoyl-Sar-boroOrn(CH=NH)-C10H16 HCl; MS: CI m/z 577.3 (M+H)+

35

#### Example 63.4.2

2-Thiophenyl-Benzoyl-Sar-boroLys(CH=NH)-C10H16; MS: CI m/z 564.2 (M-HCN)+

# Example 63.5.1

5 Pinanediol N-{N-methyl-N-[2-(Thiophenyl)-Benzoyl]Sar}-1-amido-5-thiocyanatobutane boronate; MS: CI m/z 592.2 (M+H)+

## Example 64

- N-{N-methylphenyl-N-[(3-phenyl)propionyl]glycyl}-1amido-5-aminopentaneboronate, hydrochloride salt; HRMS calcd (M+H, ethylene glycol ester)+ = 452.272062, obs.: (M+H, ethylene glycol ester)+ = 452.270496
- Example 64.1.1

  2-Benzyl-(N-Benzyl)-Sar-boroLys-C10H16 HCl; MS: CI m/z

  532.5 (M+H)+

# Example 64.1.2

20 Acetyl-Gly[N-(2-(Benzyl)-Benzyl)]-boroLys-C10H16 HCl; MS: CI m/z 560.4 (M+H)+

# Example 64.1.3

Pinanediol N-{N-methyl-N-[2-(pyrrol-1-ylmethyl)-25 Benzyl]glycyl}-1-amido-5-aminopentaneboronate, hydrochloride salt; MS: CI m/z 535.3 (M+H)+

# Example 64.2.2

[3-(Trifluoromethyl)-Benzyl]-Benzoyl-Sar-boroLys-C10H16 30 HCl; MS: CI m/z 614.3 (M+H)+

## Example 65

Piananediol  $N-\{N-\text{methyl}-N-\{(4-\text{phenyl})\text{butanoyl}\}glycyl}-1-\text{amido}-5-\text{aminopentaneboronate, hydrochloride salt; HRMS}$ 35 calcd:  $(M+H)^+=498.350313$ , obs.:  $(M+H)^+=498.350585$ .

#### Example 65,1.3

N-{N-methyl-N-{2-(pyrrol-1-ylmethyl)-Benzyl]glycyl}-1-amido-5-aminopentaneboronic acid, hydrochloride salt; MS: ESI m/z 601.3 (M+H)+

#### Example 66

N-(N-methyl-N-[(4-phenyl)butanoyl]glycyl}-1-amido-5aminopentaneboronate, hydrochloride salt; HRMS calcd: (M+H,
10 ethylene glycol ester)+ = 390.256412, obs.: (M+H, ethylene
glycol ester)+ = 390.257428.

# Example 66.1.1

Glutaryl-[N-(Phenethyl)-Gly]-boroLys-OH HCl; MS (ESI)  $(M+H)^+$  422.3

#### Example 66.1.2

Glutary1(3,3-Dimethy1)-[N-(Phenethy1)-Gly]-boroLys-OH; MS (ESI) (M+H)+ 450.5

20

5

#### Example 66.1.3

Methyl Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-OH HCl; MS (CDI)  $(M+H)^+$  490

25

#### Example 66.3.2

Methanesulfonyl-Gly-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H) + 458.3

#### Example 67

Pinanediol N-{N-methyl-N-{N-methanesulphonyl-p-phenylalanyl}glycyl}-1-amido-5-aminopentaneboronate, hydrochloride salt; HRMS calcd: (M+H)+ = 577.323113, obs.: (M+H)+ = 577.322891.

35

#### <u>Example 67.1.2</u>

Glutary1(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-C10H16; MS (ESI) (M+H)+ 584.6

# Example 67.1.3

Methyl Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H)+ 598.6

# Example 67.3.2

Methanesylfonyl-Gly-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys10 C10H16 HCl; MS (ESI) (M+H) + 592.3

# Example 68

N-{N-methyl-N-[N-methanesulphonyl-Dphenylalanyl]glycyl}-1-amido-5-aminopentaneboronate,

hydrochloride salt; HRMS calcd: (M+H, ethylene glycol
ester)+ = 469.229212, obs.: (M+H, ethylene glycol ester)+ =
469.228962.

## Example 68.2.1

20 Boc-Glu-[N-(Phenethyl)-Gly]-boroLys-OH

## Example 68.3.1

Succiny1-[N-(Phenethy1)-Gly]-boroLys-OH; MS (ESI) (M+H)+408.3

25

## Example 68.3.3

Methyl Succinyl-[N-(Phenethyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H)+ 422.3

30

#### Example 68.4.1

Methyl Glutaryl-[N-(Phenethyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H)+ 336.3

# Example 68.4.2

Methanesulfonyl-Sar-[N-(Phenethyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H)+ 457.0

## Example 69

5 Pinanediol N-{N-methyl-N-[3-(4methylphenyl)propionyl]glycyl}-1-amido-5aminopentaneboronate, hydrochloride salt; HRMS calcd:
(M+H)+ = 498.350313, obs.: (M+H)+ = 498.349676.

10 Example 69.1.1

Glutaryl-[N-(Phenethyl)-Gly]-boroLys-C10H16; MS (ESI)

(M+H) + 556.4

# Example 69.2.1

15 Boc-Glu-[N-(Phenethyl)-Gly]-boroLys-C10H16; MS (ESI) (M+H) + 671.6

# Example 69.2.2

Boc-Asp-[N-(Phenethyl)-Gly]-boroLys-C10H16; MS (ESI)  $(M+H)^+$  657.6

#### Example 69.2.3

Boc-Glu(OCH3)-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H)+ 685.6

25

#### Example 69.3.2

Methanesulfonyl-Gly-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (NH<sub>3</sub>CI) (M+H) + 577

30 Example 69.4.1

Methyl Glutaryl-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl; MS

(ESI) (M+H)+ 570.5

## Example 69.4.2

-158-

Methanesulfonyl-Sar-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (NH<sub>3</sub>CI) (M+H)+ 591.4

# Example 70

N-{N-methyl-N-[3-(4-methylphenyl)propionyl]glycyl}-1amido-5-aminopentaneboronate, hydrochloride salt; HRMS calcd: (M+H, ethylene glycol ester)+ = 390.256412, obs.: (M+H, ethylene glycol ester)+ = 390.256960.

10 Example 71

Pinanediol N-{N-methyl-N-[2-(methyl(4-methoxyphenyl))benzoyl]glycyl}-1-amido-5-amino-pentaneboronate, hydrochloride acid salt; LRMS (M+H)+ = 576.3.

15

## Example 72

Pinanediol N-{N-methyl-N-[2-(methyl(4-methylphenyl))benzoyl]glycyl}-1-amido-5-amino-pentaneboronate, hydrochloride acid salt; LRMS (M+H)+ = 560.5.

# Example 72.1.3

Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroLys-OH HC1; MS (ESI) (M+H)+ 379.0

25

20

# Example 73

Pinanediol N-(N-((O-tert-butyl))) methylenecarboxylate)-N-[(3-phenyl)] propionyl]-glycyl}-1-amido-5-aminopentaneboronate, hydrochloride salt; LRMS  $(M+H)^+$  = 584.

# Example 73.1.2

Succinyl-[N-(3-(Methyl)-Phenethyl)-Gly]-boroLys-C10H16 HCl; MS (ESI) (M+H) + 556

35

PCT/US94/11280

#### Example 74

Pinanediol  $N-\{N-\text{methyl}-N-\{(3-\text{phenyl})\text{propionyl}\}\text{glycyl}\}-1-\text{amido}-4-(\text{formamidino})\text{butylboronate}, hydrochloride salt; LRMS <math>(M+H)^+=497$ .

5

# Example 75

N-(N-methyl-N-[(3-phenyl)propionyl]glycyl-1-amido-4-(N-methylguanidino)butylboronate, hydrochloride salt; LRMS (M+H, ethylene glycol ester)+ = 418.3.

10

## Example 75.3.1

(2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroArg-OH HCl; MS (ESI) (M+H) + 483.1

15

# Example 76

 $N-\{N-\text{methyl}-N-[(3-\text{phenyl})\text{propionyl}]\text{glycyl}\}-1-\text{amido}-4-(formamidino})$ butylboronate, hydrochloride salt; LRMS (M+H, ethylene glycol ester)+ = 389.2.

20

#### Example 76.1.1

Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys(CH=NH)-OH HCl; MS (ESI) (M+H) + 403.0

# Example 77.1.1

25 Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys(CH=NH)-C10H16 HCl; MS (NH<sub>3</sub>CI) (M+H) + 537.3

# Example 78.1.2

Phenoxyacetyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H)+ 378.3

# Example 78.1.3

Thiophenacetyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl; MS (ESI) (M+H) + 394.2

35

# Example 79.1.2

Phenoxyacetyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl; MS (NH<sub>3</sub>CI)  $(M+H)^+$  512.3

5 <u>Example 79.1.3</u>

Thiophenacetyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl; MS (NH<sub>3</sub>CI) (M+H) + 528.3

# Tables 1-79

# 5 <u>Table 1</u>

Formula I : A = -B(OH)2 ; X = guanidinyl ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = CH3

	.1	.2	.3
1.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
1.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
1.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
1.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
1.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
1.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
1.7	-C(O)naphth-1-yl	-C(O)CH <sub>2</sub> (naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
1.8	-C(O)naphth-2-yl	-C(O)CH <sub>2</sub> (naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
1.9	-C(O)o-biphenyl	-C(O)CH <sub>2</sub> (o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
1.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
1.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
1.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
1.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhOPh)
1.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
1.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
1.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhNHPh)
1.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
1.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-PhSPh)
1.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
1.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
1.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
1.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (m-PhCH <sub>2</sub> SPh)	-C(O)CH2CH2(m- PhCH2SPh)
	O(O) DI OU ODI	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
1.24	-C(O)p-PhCH <sub>2</sub> SPh		-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
1.25	-C(O)adamantyl	-C(O)CH <sub>2</sub> (adamantyl)	-C(O)CH2CH2((cyclopentyl)
1.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclohexyl)
1.27	-C(O)cyclohexyl	-C(O)CH <sub>2</sub> (cyclohexyl)	
1.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
1.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
1.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
1.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
1.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
1.33	-C(O)furan-2-yl	-C(O)CH <sub>2</sub> (furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
1.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)

		C(C)C(Ia(thisphon 2 vI)	-C(O)CH2CH2(thiophen-2-yl)
	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-vl)	-C(O)CH2CH2(thiophen-2-yl)
	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
1.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (minda20-2-yi)
1.38	-C(O)oxazo-2-yl	-C(O)CH <sub>2</sub> (oxazo-2-yl)	
1.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
1.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2-
			vi)
1.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH2CH2(benzofuran-3-
l [			yl)
1.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-yl)	C(O)CH2CH2(benzothiophen-
			2-yl)
		Corporation (15 - han 2 of)	-C(O)CH2CH2(thiophen-2-yl)
1.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	
1.44	-C(O)benzimidazo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo-
			2-vI)
1.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-
			yl) -C(O)CH2CH2(benzothiazo-2-
1.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	
			vl) -C(O)p-Ph(P(O)Ph3)
1.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)Ph-4-(fluoren-9-yl)
1.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)indol-2-vl
1.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)cyclohexyl-2-(Ph)
1.50	-	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclonexyi-2-(Fii)
	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth		
	-2-vl)	2(0)	-C(O)tetrahydrothiophen-3-yl-
1.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-(Ph)	4-(Ph)
1.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
1.53	-C(O)tetrahydroisoquinolin-1-	-C(O)tetrahydroisoquinolin-3-yl	-C(O)CH2((2-oxo)indolin-3-
1	yl	<u> </u>	yl)
1.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
	one)		one)
1.55	-C(O)CH2(N-dihydroimidazol-	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2-
1	2-one)	one)	one)
1.56	<b>C</b> O-	,co-	50-
1	- 10	~ N 0	<u>\(\frac{1}{4}\) \(\frac{1}{4}\)</u>
	(A)" P	<i>US</i> ''4	
			<u> </u>
1.57			
1.57	l : 유	1	Ĭ ŭ
	-oc N N H	-00° N°O	-oc N S
	l <sup>™</sup>		
1.58	00.0	-oc.	-oc o
1	-00 0	1 33	-0C 0
1	'N' >	\ \( \( \bullet \) \\ \( \)	١ "ــــــــــــــــــــــــــــــــــــ
Ī		'N'	
1	\=/	1 英	
			1
1			
1.59	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph

1.60	-C(O)pvridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH2Ph)
1.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
1.62	-C(O)CH2O(o-PhCH2OII)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
1.63	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
1.64	-C(O)CH <sub>2</sub> O(o-PhCOOCH <sub>3</sub> )	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
1.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)		-C(O)CH2O(p-PhCH2COOH)
1.66			

Table 2

Formula I:  $A = -B(OH)_2$ ; X = guanidinyl;  $R^3 = table below$ ;  $R^{11} = -CH_2(p-PhOH)$ .

	.1	.2	.3
2.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
2.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPb
2.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
2.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
2.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
2.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
2.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
2.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
2.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
2.10	-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
2.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
2.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
2.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
2.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhOPh)
2.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
2.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
2.18	-C(O)p-PhNHPh	-C(O)CH <sub>2</sub> (p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
2.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
2.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
2.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
2.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
2.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
			PhCH <sub>2</sub> SPh)
2.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
2.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
2.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
2.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
2.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
2.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
2.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
2.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)

			5.60.00 (0.17.4.2)
2.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
2.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
2.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
2.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
2.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
2.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
2.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
2.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
2.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2- vl)
2.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3- vl)
2.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH2CH2(benzothiophen- 2-yl)
2.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
2.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-vl)
2.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-yl)
2.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo-2-yl)
2.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
2.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
2.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-vl
2.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
2.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
2.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
2.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
2.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
2.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)
2.56	CO-	CVN CO.	CO.
2.57	-oc N NH	-oc-NHO	-oc N s

2.58	°C N	00 z z ~	oc N
2.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph
2.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
2.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
2.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
2.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
2.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
2.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
2.66	-0C 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

5

Formula I :  $A = -B(OH)_2$ ; X = guanidinyl;  $R^3 = table below$ ;  $R^{11} = -CH_2CH_2Ph$ .

	.1	.2	.3
3.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
3.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH2SPh
3.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
3.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH <sub>2</sub> OH
3.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
3.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
3.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)Cl12CH2(napth-1-yl)
3.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
3.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
3.10	-C(O)m-biphenyl	-C(O)CII <sub>2</sub> (m-hiphenyl)	-C(O)CH2CH2(m-biphenyl)
3.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-hiphenyl)	-C(O)CH2CH2(p-biphenyl)
3.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
3.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
3.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
3.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
3.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
3.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
3.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
3.20	-C(O)m-PhSPh	-C(O)CII <sub>2</sub> (m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
3.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
3.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)

3.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(in-PhCH2SPh)	-C(O)CH2CH2(m- PhCH2SPh)
		-C(O)Cl12(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
3.24	-0(0)// 1::01:201::		-C(O)CH2CH2(adamantyl)
3.25	-C(O/acestratify:	-C(O)CH2(adamantyl)	-C(O)CH2CH2((cyclopentyl)
3.26	C(C)0) Citi (Citi)	-C(O)Cli2(cyclopentyl)	-C(O)CH2CH2(cyclohexyl)
3.27	-C(O)cyclohexyl	-C(O)CH <sub>2</sub> (cyclohexyl)	
3.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
3.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
3.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
3.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
3.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
3.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
3.34	-C(O)furan-3-yl	-C(O)CH <sub>2</sub> (furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
3.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
3.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
3.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
3.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
3.39	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH2CH2(benzoturan-2-
3.40	-C(O)benzoiwan-2-yi	-C(O)(:112(12:112:112:112:12:12:12:12:12:12:12:12:1	vi)
3.41	-C(O)benzofuran-3-yl	-C(O)Cl12(benzofuran-3-yl)	-C(O)CH2CH2(benzofuran-3-
5.41	-C(O)telizolulai-5-yi		vl)
3.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	-
J.72	-2(0)2011201110 111011 2 }	yl)	C(O)CH2CH2(benzothiophen-
			2-yl)
3.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
3.44	-C(O)benzimidazo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo-
			2-yl)
3.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-
			vl)
3.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo-2-
•		2.6.000	vl) -C(O)p-Ph(P(O)Ph3)
3.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	
3.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl) -C(O)indol-2-yl
3.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)cyclohexyl-2-(Ph)
3.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth-	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclonexyr-2-(rii)
	2-yl)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-yl-
3.51	-C(O)pyrrolidin-3-yl-4-(Ph)	(Ph)	4-(Ph)
2.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )
1 4 7/		-C(O)tetrahydroisoquinolin-3-	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-
3.52			
3.53	-C(O)tetrahydroisoquinolin-1-yl	h .	<u> </u>
3.53		yl	yl)
	-C(O)CH <sub>2</sub> (N-benzimidazol-2-	yl -C(O)CH <sub>2</sub> (N-benzoxazol-2-	<u> </u>
3.53 3.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	yl -C(O)CH <sub>2</sub> (N-benzoxazol-2- one)	yl) -C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
3.53	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one) -C(O)CH <sub>2</sub> (N-dihydroimidazol-	yl -C(O)CH <sub>2</sub> (N-benzoxazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrooxazol-2-	yl) -C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
3.53 3.54 3.55	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one) -C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	yl  -C(O)CH <sub>2</sub> (N-benzoxazol-2-one)  -C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	yl) -C(O)CH <sub>2</sub> (N-benzothiazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrothiazol-2
3.53	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one) -C(O)CH <sub>2</sub> (N-dihydroimidazol-	yl -C(O)CH <sub>2</sub> (N-benzoxazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrooxazol-2-	yl) -C(O)CH <sub>2</sub> (N-benzothiazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)
3.53 3.54 3.55	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one) -C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	yl  -C(O)CH <sub>2</sub> (N-benzoxazol-2-one)  -C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	yl) -C(O)CH <sub>2</sub> (N-benzothiazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)
3.53 3.54 3.55	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one) -C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	yl  -C(O)CH <sub>2</sub> (N-benzoxazol-2-one)  -C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	yl) -C(O)CH <sub>2</sub> (N-benzothiazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)

3.57	-oc NHNH	-oc N	-00 N S
3.58	OC OC	-0 C Z Z - (1)	(2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
3.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
3.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
3.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
3.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
3.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
3.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
3.65	-C(U)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
3.66	0, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,		

5

Formula I : A = -B(OH)<sub>2</sub> ; X = guanidinyl ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -Ph.

	11	.2	.3
4.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
4.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH2SPh
4.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
4.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
4.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
4.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
4.7	-C(O)naphth-1-yl	-C(O)Cli2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
4.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
4.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
4.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
4.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
4.13	-C(O)o-PhOPh	-C(O)Cl12(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
4.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
4.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
4.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
4.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
4.18	-C(O)p-PhNHPh	-C(O)Cllp(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
4.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)

4.20	-C(O)ın-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
4.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
4.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
4.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m- PhCH <sub>2</sub> SPh)
4.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	-C(O)CH2CH2(p-PhCH2SPh)
4.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
4.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
4.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
4.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
4.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
4.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
4.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
4.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
4.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
4.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
4.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
4.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
4.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
4.38	-C(O)nxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
4.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
4.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2-yl)
4.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
4.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen- 2-yl)
4.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
4.44	-C(O)benzimidazo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (henzimidazo- 2-yl)
4.45	-C(O)benzoxazu-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
4.46	-C(O)benzothiazo-2-yl	-C(O)C112(benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo-2 yl)
4.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
4.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
4.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
4.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
4.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
4.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
4.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
4.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
4.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-diliydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2 one)

	<del> </del>		
4.56		QN CO.	CO.
4.57	-oc N NH	-oc N 0	-oc N
4.58	-0ç o	°°, z, z, (1)	0,000
4.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph
4.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-vl))	<sup>L</sup> C(O)Ph-3-(CH <sub>2</sub> Ph)
4.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
4.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH <sub>2</sub> O(m-PhCH <sub>2</sub> OH)	-C(O)CH2O(p-PhCH2OH)
4.63	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
4.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
4.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
4.66			

 $\frac{Table\ 5}{\text{Formula}\ I: A = -B(OH)_2: X = guanidinyl: } R^3 = table\ below: R^{11} = -CH_2(naphth-2-yl).$ 

Т	.1	.2	.3
5.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
5.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
5.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
5.4	-C(O)o-PhCH <sub>2</sub> OH	-C(O)m-PhCH <sub>2</sub> OH	-C(O)p-PhCH <sub>2</sub> OH
5.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
5.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
5.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
5.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
5.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
5.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
5.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
5.12	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
5.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
5.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhOPh)
5.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
5.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
5.18	-C(O)n-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhNHPh)
5.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
5.20	-C(O)m-PhSPh	-C(O)CH2(in-PhSPh)	-C(O)CH2CH2(m-PhSPh)
5.21	-C(O)n-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
5.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
	(0,012012		PhCH <sub>2</sub> SPh)
5.23	-C(O)m-PhCl·l <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-
			PhCH <sub>2</sub> SPh)
5.24	-C(O)p-PhCH2SPh	-C(O)Cl12(p-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-
1			PhCH <sub>2</sub> SPh)
5.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
5.26	-C(O)cyclopentyl	-C(O)CH <sub>2</sub> (cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
5.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
5.28	-C(0)CH2O(cyclopentyl)	-C(O)CH <sub>2</sub> NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
5.29	-C(O)CH2O(cyclohexyl)	-C(O)Cl12NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
5.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
5.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
5.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
5.33	-C(O)furan-2-yl	-C(O)CH <sub>2</sub> (furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
5.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
5.35	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
5.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
5.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
5.38	-C(O)oxazo-2-vl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
			-C(O)CH2CH2(thioazo-2-yl)
5.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)

5.40	C(O)1	Lower	
L	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 2-yl)
5.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 3-vI)
5.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-vl)
5.43	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
5.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
5.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-yl)
5.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-yl)
5.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
5.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-vl)	-C(O)Ph-4-(fluoren-9-yl)
5.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indol-2-vl
5.50	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
5.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3- vl-4-(Ph)
5.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
5.53	-C(O)tetrahydroisoquinolin- 1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- vl)
5.54	-C(O)CH <sub>2</sub> (N-benzimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
5.55	-C(O)CH <sub>2</sub> (N- dihydroimidazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
5.56	CO.		COO
5.57	ос — м м н	00° 00° 00° 00° 00° 00° 00° 00° 00° 00°	-oc N S
5.58	-0Ç 0	oc, z	-oc 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
5.59	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C3H7)CH2Ph
5.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
5.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph

172

5.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
5.63	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
5.64		-C(O)CH2O(m-	-C(O)CH2O(p-
5.65	-C(O)CH <sub>2</sub> O(o- PhCH <sub>2</sub> COOH)	PhCH2COOH)	PhCH2COOH)
5.66			

Formula I : A = -B(OH)<sub>2</sub> ; X = -CH<sub>2</sub>NH<sub>2</sub> ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = CH<sub>3</sub>

	.1	.2	.3
6.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
6.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH2SPh
6.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
6.4	-C(O)o-PhCH <sub>2</sub> OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
6.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
6.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
6.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
6.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
6.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
6.10	-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
6.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-hiphenyl)	-C(O)CH2CH2(p-biphenyl)
6.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
6.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
6.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
6.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
6.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
6.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
6.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
6.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
6.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
6.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
"	6(6),012012211		PhCH <sub>2</sub> SPh)
6.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-
0.23	S(0) 1 2 2 3 2 3		PhCH <sub>2</sub> SPh)
6.24	-C(O)p-PhCH2SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	-C(O)CH2CH2(p-
0.2.	0(0)/1 1 1 1 2 2		PhCH <sub>2</sub> SPh)
6.25	-C(O)adamantyl	-C(O)Cl12(adamantyl)	-C(O)CH2CH2(adamantyl)
6.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
6.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
6.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
6.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
6.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)

		<del>,</del>	
6.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
6.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
6.33	-C(O)furan-2-yl	-C(O)CH <sub>2</sub> (furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
6.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-vl)
6.35	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-vl)
6.36	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
6.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
6.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
6.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thioazo-2-yl)
6.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-
	(0,000000000000000000000000000000000000	0(0,012(10111111111111111111111111111111	2-vI)
6.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 3-yl)
6.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	-
	1	yl)	C(O)CH2CH2(benzothiophen
<u>L</u>	<u>                                     </u>		-2-vI)
6.43	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
6.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-vl)
6.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzox <sub>2</sub> zo-2-yl)
6.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-vl)
6.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
6.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-vl)	-C(O)Ph-4-(fluoren-9-yl)
6.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indol-2-vl
6.50	- C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
6.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3- vl-4-(Ph)
6.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
6.53	-C(O)tetrahydroisoquinolin- 1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- vl)
6.54	-C(O)CH <sub>2</sub> (N-benzimidazol-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
	2-one)	one)	one)
6.55	-C(O)CH <sub>2</sub> (N- dihydroimidazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
6.56	CO.	COO	C C C C C C C C C C C C C C C C C C C
6.57	-oc N NH	-oc_N <sub>H</sub> o	-oc^Nus

174

6.58	oc N	٥ کر چی کے حرگ	OC O
6.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph
6.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(Cll <sub>2</sub> (thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
6.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
6.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
6.63	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
6.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
6.65	-C(O)CH <sub>2</sub> O(o- PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p- PhCH <sub>2</sub> COOH)
6.66	-oc <sup>k</sup> o		

5

Formula I : A = -B(OH)2 ; X = -CH2NH2 ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -CH2(p-PhOH).

·	1	.2	.3
7.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
7.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
7.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
7.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
7.5	-C(O)n-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
7.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
7.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
7.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
7.9	-C(O)o-hiphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
7.10	-C(O)m-hiphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
7.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
7.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
7.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
7.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
7.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
7.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
7.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
7.19	-C(O)o-PhSPh	-C(O)Cl12(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
7.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
7.21	-C(O)p-PhSPh	-C(O)Cl12(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
7.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (o-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o- PhCH <sub>2</sub> SPh)

7.00	Toron Brown		
7.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
7.24	C(C) - PLOU-CPL	0/0/0//	PhCH <sub>2</sub> SPh)
7.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-
1205	0(0)	0.00	PhCH <sub>2</sub> SPh)
7.25	-C(O)adamantyl	-C(O)CH <sub>2</sub> (adamantyl)	-C(O)CH2CH2(adamantyl)
7.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
7.27	-C(O)cyclohexyl	-C(O)CH <sub>2</sub> (cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
7.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NII(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
7.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
7.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
7.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
7.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
7.33	-C(O)furan-2-yl	-C(O)CH <sub>2</sub> (furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
7.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
7.35	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2- vl)
7.36	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2- vl)
7.37	-C(O)imidazo-2-yl	-C(O)CII <sub>2</sub> (imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
7.38	-C(O)oxazo-2-yl	-C(O)Cl12(oxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (oxazo-2-yl)
7.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thioazo-2-yl)
7.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-
			2-yl)
7.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 3-yl)
7.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	<u> -</u>
		yl)	C(O)CH2CH2(benzothiophen
7.43	CONTRACTOR		-2-vl)
7.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
7.44	-C(O)benzimidazo-2-yl	C(())CHe/har-i-i-i- Q ()	yl)
	-C(C)(kelizilindiiz(F2-y)	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo-
7.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	2-yl)
	CONSTRUCTION OF THE PROPERTY O	-C(O/C.112(tielizoxazo-z-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2- yl)
7.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo-
	1	Storenzelinazorz-yl)	2-yl)
7.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
7.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-vl)
7.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
7.50	-	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth		
	-2-yl)		
7.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-
7.52	-C(O)tetrahydronaphth-1-vl	(Ph) -C(O)tetrahydronaphth-2-yl	yl-4-(Ph)
7.53	-C(O)tetrahydroisoguinolin-	-C(O)tetrahydroisoquinolin-3-	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )
	1-yl	vl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- vl)
			Y1/
7.54		-C(O)CH2(N-henzoxazol-2-	
	-C(O)CH <sub>2</sub> (N-benzimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
7.55	-C(O)CH <sub>2</sub> (N-benzimidazol-		

7.56	CO.	CO.	
7.57	-oc NH NH	0000	oc N s
7.58		0 2 z z - (_)	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
7.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C3H7)CH2Ph
7.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-C(O)Ph-3-(CH2Ph)
7.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
7.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
7.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOII)	-C(O)CH2O(p-PhCOOH)
7.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
7.65	-C(O)CH <sub>2</sub> O(o- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(p- PhCH <sub>2</sub> COOH)
7.66	-oc 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

 $\label{eq:Formula I:A = -B(OH)_2: X = CH_2NH_2; R^3 = table below: R^{11} = -CH_2CH_2Ph.}$ 

	.1	.2	.3
8.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
8.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
8.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
8.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
8.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
8.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
8.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
8.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
8.9	-C(O)o-biphenyl	-C(O)CH <sub>2</sub> (o-biphenyl)	-C(O)CH2CH2(o-hiphenyl)
8.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
8.12	-C(O)p-hiphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
8.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
8.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)Cl12CH2(m-PhOPh)

0.5	Taga sasa		
8.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
8.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
8.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
8.18	-C(O)p-PhNHPh	-C(O)CH <sub>2</sub> (p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
8.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
8.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
8.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
8.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
			PhCH <sub>2</sub> SPh)
8.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (m-PhCH <sub>2</sub> SPh)	-C(O)CH2CH2(m-
			PhCH2SPh)
8.24	-C(O)p-PhCH2SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-
1005			PhCH <sub>2</sub> SPh)
8.25	-C(O)adamantyl	-C(O)CH <sub>2</sub> (adamantyl)	-C(O)CH2CH2(adamantyl)
8.26	-C(O)cyclopentyl	-C(O)CH <sub>2</sub> (cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
8.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
8.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
8.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
8.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
8.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
8.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
8.33	-C(O)furan-2-yl	-C(O)CII <sub>2</sub> (furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
8.34	-C(O)furan-3-yl	-C(O)CH <sub>2</sub> (furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
8.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
8.36	-C(O)thiophen-2-yl	2400	yl)
0.50	-C(O)unopnen-2-yi	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
8.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	yl)
8.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (imidazo-2-yl)
8.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
8.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
		-C(O)C.112(benzoruran-2-yi)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 2-vl)
8.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(U)CH2CH2(benzofuran-
	L	(0,0112(total2(101al1-1,-y1)	3-vl)
8.42	-C(O)benzothiophen-2-yl	-C(O)Cl12(benzothiophen-2-	-
		yl)	C(O)CH2CH2(benzothiophen
9.42	0(0)		-2-yl)
8.43	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
8.44	C(O)barrinida - 2 )	0.00	yl)
0.44	-C(O)benzimidazo-2-yl	-C(O)CII <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo-
8.45	-C(O)benzoxazo-2-yl	C(O)CHe/bessesses 2 "	2-yl)
	O COMMISSIONAL (P-Z-y)	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-
8.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	(C)CH-CH-(I
		-C.C.C.112(CC112O(thaz(r-2-yl))	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo-
8.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	2-yl) -C(())p-Ph(P(O)Ph3)
8.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-vl)	-C(O)Ph-4-(fluoren-9-yl)
8.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(())indol-2-vl
		1 2007/////////// 2-11	-CAC/MIGOI-Z-VI

- 8. A method of treating a physiological disorder in a warm blooded animal catalyzed by trypsin-like enzymes comprising administering to an animal in need such treatment an effective amount of a compound of claim 2.
- 9. A method of treating a physiological disorder in a warm blooded animal catalyzed by trypsin10 like enzymes comprising administering to an animal in need such treatment an effective amount of a compound of claim 3.

# INTERNATIONAL SEARCH REPORT

Int tional application No.

2000	1/4021	100
PK 1/1	INGA/I	1 /AL

1	SSIFICATION OF SUBJECT MATTER				
	:A61K 31/69; C07F 5/02, 5/04	•			
	US CL: 514/64; 546/13; 548/110; 558/5,288,298; 562/7 According to International Patent Classification (IPC) or to both national classification and IPC				
	LDS SEARCHED	matorial Cassification and If C			
		d by alonifostics and also			
	ocumentation searched (classification system followe	d by classification symbols)			
U.S. :	514/64; 546/13; 548/110; 558/5,288,298; 562/7	·	·		
Dogumento	tion searched other than minimum documentation to th	e extent that such documents are incl	uded in the fields seconded		
Documenta	Hou residues office men immunium cocamenescion e di	e exient diat such documents are nici			
Flectronic	lata base consulted during the international search (n	ome of data base and, where practic	shie search terms used)		
		and or one one and, where present	more, senten terms used, .,		
CAS ON	LINE 2 Searched: REGISTRY (Structure-based Search	n) -			
	, , , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,			
C. DOC	CUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where a	poropriate, of the relevant passages	Relevant to claim No.		
		· · · · · · · · · · · · · · · · · · ·	The state of the s		
X	US, A, 5,187,157 (KETTNER ET A	AL.) 16 FEBRUARY 199	3, 1-9		
	see columns 3, line 13-column 4,	line 21.			
Α	US, A, 4,537,773 (SHENVI) 27 /	AUGUST 1985, see ent	ire 1-9		
	document.	7			
Α	US, A, 5,169,841 (KLEEMAN ET /	AL.) 08 DECEMBER 199	2, 1-9		
	see entire document.	i			
	·				
	•				
Furth	er documents are listed in the continuation of Box C	See patent family anne	x.		
• Spe	ecial categories of cited documents:	"T" later document published after the	be international filing date or priority		
	cument defining the general state of the art which is not considered be of particular relevance	principle or theory underlying t	application but cited to understand the ne invention		
	tier document published on or after the international filing date	"X" document of particular relevan	ce; the claimed invention cannot be		
"L" doe	current which may throw doubts on priority claim(s) or which is	when the document is taken alor	pasidered to involve an inventive step ac		
cited to establish the publication date of another citation or other special reason (as specified)  "Y"  document of particular relevance; the claimed invention cannot be					
"O" document referring to an oral disclosure, use, exhibition or other combined with one or more other such documents, such combination being obvious to a nerso stilled in the est					
*P* document published prior to the international filing date but later than *& document member of the same patent family					
the	priority date claimed				
Date of the	actual completion of the international search	Date of mailing of the internationa			
13 DECE	MBER 1994	10 JANY	<del>ชชวี ,                                  </del>		
Name and -	pailing address of the ISA/IIS	Authorized officer	<i>^////</i>		
Commissio	nailing address of the ISA/US ner of Patems and Trademarks	Variation Trees	211/ami		
Box PCT Washington	n, D.C. 20231	MICHAEL G. AMBROSE	- IN house I		
•	0. (703) 305-3230	Telephone No. (703) 308-1235	h 1		

Form PCT/ISA/210 (second sheet)(July 1992)\*

		-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
8.50		2C(O)Cyclopentyi-2-(t ii)	
	C(O)C(CH3)2NHSO2(naphth		
	-2-vl)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-
8.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)(eiranydioidiai-5-yi (Ph)	vl-4-(Ph)
		-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
8.52	-C(O)tetrahydronaphth-1-yl	-(.(O)(et/ai/yd/(nia)nid-2 )1	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-
8.53	-C(O)tetrahydroisoquinolin-	-C(O)tetrahydroisoquinolin-3-	vi)
	1-yl	yl 22	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
8.54	-C(O)CH <sub>2</sub> (N-benzimidazol-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-	one)
	2-one)	one)	-C(O)CH2(N-dihydrothiazol-
8.55	-C(O)CH2(N-	-C(O)CH <sub>2</sub> (N-dihydrooxazol-	2-one)
	dihydroimidazol-2-one)	2-one)	
8.56	,co-	co-	
	1 . 10	N.O	N.Y.
	1 AY Y	('_`Y'``Y	
	ال ا		
			. 0
8.57	0	l C	l: ŭ
	NH NH	OCCN O	-OC N'S
	-OC N NH	1-00 "}-(	
	( )		
			00.0
8.58	-oc Q	-OC <sub>3</sub>	-oc 0
	I IN INCOME	N.	N O
		[]	
1	\ \( \( \)	Į Ņ.	\ <u>\_</u> \
1	l		
0.50	C(C)N(CHa)CHaPh	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
8.59	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph	-C(O)Ph-3-(CH2(thiophen-2-	-C(O)Ph-3-(CH2Ph)
8.60	-C(O)pyridin-3-yl-5-(Ph)	yl))	
		-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
8.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OFH -C(O)CH2(O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
8.62	-C(O)CH2O(o-PhCH2OH)		-C(O)CH <sub>2</sub> O(p-PhCOOH)
8.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2()(m-PhCOOH)	
8.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-
8.65	-C(O)CH2O(o-	-C(O)CH <sub>2</sub> O(m-	
1	PhCH <sub>2</sub> COOH)	PhCH2COOH)	PhCH <sub>2</sub> COOH)
8.66	-00 0	•	}
1	I TUR		
1			
1	\ <b>\</b>	]	
1		ł	
1	\ <b>\_</b> \		1

5

Formula I : A = -B(OH)2 ; X = CH2NH2:  ${\bf R}^3$  = table below ;  ${\bf R}^{11}$  = -Ph.

r	1	.2	.3
9.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph

9.2	-C(O)Cl12OPh	-C(O)CH2NHPh	-C(O)CH2SPh
9.3	-C(O)o-PhOH	-C(O)m-PhOH	
9.3	-C(O)o-PhCH2OH		-C(O)p-PhOH
9.5		-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
9.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
9.7	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
9.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
	-C(O)naphth-2-yl	-C(O)CH <sub>2</sub> (naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
9.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
9.10	-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
9.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
9.13	-C(O)o-PhOPh	-C(O)CII <sub>2</sub> (o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
9.14	-C(O)m-PhOPh	-C(O)CH <sub>2</sub> (m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
9.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
9.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
9.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
9.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNIIPh)	-C(O)CH2CH2(p-PhNHPh)
9.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
9.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
9.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
9.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CII2CH2(o-
	<u>                                     </u>		PhCH <sub>2</sub> SPh)
9.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
			PhCH2SPh)
9.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-
			PhCH2SPh)
9.25	-C(O)adamantyl	-C(O)Cl12(adamantyl)	-C(O)CH2CH2(adamantyl)
9.26	-C(O)cyclopentyl	-C(O)CH <sub>2</sub> (cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
9.27	-C(O)cyclohexyl	-C(O)Cll <sub>2</sub> (cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
9.28	-C(O)CH2O(cyclopentyl)	-C(O)CII2NII(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
9.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
9.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
9.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
9.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-vI)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
9.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
9.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
9.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
w		, , , , , , , , , , , , , , , , , , ,	vl)
9.36	-C(O)thiophen-2-yl	-C(U)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
			vl)
9.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
9.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (oxazo-2-yl)
9.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-vl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thioazo-2-yl)
9.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzoturan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-
	<u> </u>		2-vi)
9.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yI)	-C(O)CH2CH2(benzofuran-
			3-yl)
9.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	-
		yl)	C(O)CH2CH2(benzothiophen
			-2-yl)

9.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
		-C(O)CH2(benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo-
9.44	-C(O)benzimidazo-2-yl		2-vl)
9.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-
			vI) -C(O)CH2CH2(benzothiazo-
9.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	2-yl)
9.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
9.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
9.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indol-2-yl
9.50		-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
9.50	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-yl)		
0.61	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-
9.51	-C(O)pyrrolldin-3-y1-4-(1 ii)	(Ph)	yl-4-(Ph)
9.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )
9.53	-C(O)tetrahydroisoquinolin-	-C(O)tetrahydroisoquinolin-3-	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- vl)
	1-yl	yl	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
9.54	-C(O)CH <sub>2</sub> (N-benzimidazol-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	one)
0.65	2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-	-C(O)CH2(N-dihydrothiazol-
9.55	-C(O)CH <sub>2</sub> (N- dihydroimidazol-2-one)	2-one)	2-one)
9.56	co-	.co-	CO-
		CNS	
9.57	-oc NHNH	-oc N 0	-oc N s
9.58	-oc o	00 z z	-oc 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
9.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
9.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
9.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C <sub>2</sub> H <sub>5</sub> )OPh	-C(O)CH <sub>2</sub> OCH <sub>2</sub> Ph
9.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH <sub>2</sub> O(m-PhCH <sub>2</sub> OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
9.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
9.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
9.65	-C(O)CH2O(o-	-C(O)CH2O(m-	-C(O)CH2O(p-
1	PhCH2COOH)	PhCH2COOH)	PhCH2COOH)

9.66	-0C 0		

Table 10 Formula I :  $A = -B(OH)_2$  ;  $X = -CH_2NH_2$ ;  $R^3 = table below$  ;  $R^{11} = -CH_2(naphth-2-yl)$ .

	.1	.2	.3
10.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
10.1	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
10.2	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
10.3	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
10.4	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
10.5	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
10.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
10.7	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
10.8	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
10.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
10.10	-C(O)p-biphenyl	-C(O)Cl12(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
10.12	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
10.13	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhOPh)
10.15	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CII2CH2(o-PhNHPh)
10.16	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
10.17	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
10.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
10.19	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
10.20	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
10.21	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
10.22	-C(O)0-1 IIC11201 II	(6,6.12,1.1.1.2.1.1.1.1.1.1.1.1.1.1.1.1.1.1.	PhCH <sub>2</sub> SPh)
10.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
10.23	-C(0) : 110.1201 =		PhCH <sub>2</sub> SPh)
10.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-
10.5			PhCH <sub>2</sub> SPh)
10.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
10.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
10.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
10.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
10.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyi)	-C(O)CH <sub>2</sub> S(cyclohexyl)
10.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
10.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
10.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
10.33	-C(O)furan-2-yl	-C(O)Cl12(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
10.34	-C(O)furan-3-yl	C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
10.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
10.36	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
10.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
10.37	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
10.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
10.40	-C(O)benzofuran-2-yl	-C(O)Cl12(benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2-
10.70			vl)

10.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
10.42	-C(O)benzothiophen-2-yl	-C(O)Cl12(benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-vl)
10.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
10.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
10.45	-C(O)benzoxazo-2-yl	-C(O)Cl12(benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2- vl)
10.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-vl)
10.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
10.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(Пиоren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
10.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
10.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-vl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
10.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(Ö)tetrahydrothiophen-3-yl- 4-(Ph)
10.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )
10.53	-C(O)tetrahydroisoquinolin-1- yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
10.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
10.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
10.56		ON CO.	
10.57	-oc NHNH	-oc N	oc N s
10.58	o Z	٥ ٥ عي- حي	-oc 200
10.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph
10.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
10.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
10.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
10.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH <sub>2</sub> O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
10.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )

10.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH2O(p-PhCH2COOH)
10.66	OCN CONTO		

Table 11

Formula I : A = -B(OH)<sub>2</sub> ; X = -SC(=NH)NH2 ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = CH<sub>3</sub>

	1	.2	.3
11.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH <sub>2</sub> CH <sub>2</sub> Ph
11.2	-C(O)CH2OPb	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
11.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
11.5	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
11.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
11.5	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
11.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
11.8	-C(O)naphth-2-yi	-C(O)CH2(naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
11.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
11.10	-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
11.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-hiphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
11.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
11.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
11.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
11.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
11.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
11.17	-C(O)n-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
11.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
11.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
11.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
11.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(U)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
11.22			PhCH <sub>2</sub> SPh)
11.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
			PhCH <sub>2</sub> SPh)
11.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCl12SPh)	-C(O)CH2CH2(p-
			PhCH <sub>2</sub> SPh)
11.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
11.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
11.27	-C(O)cyclohexyl	-C(O)CH <sub>2</sub> (cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
11.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH <sub>2</sub> NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
11.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
11.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
11.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
11.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
11.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)

11.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
11.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
11.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
11.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
11.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
11.39	-C(O)thioazo-2-yl	-C(O)CH <sub>2</sub> (thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
11.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2-yl)
11.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzoturan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
11.42	-C(O)benzothiophen-2-yl	-C(O)Cl12(benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-yl)
11.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
11.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
11.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2- yl)
11.46	-C(O)benzothiazo-2-yl	-C(U)CH2(benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (henzothiazo- 2-yl)
11.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
11.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
11.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
11.50	- C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-vl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
11.51	-C(O)рупоlidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
11.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
11.53	-C(O)tetrahydroisoquinolin-1- yl	-C(O)terrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- vl)
11.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2- one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
11.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol- 2-one)
11.56	C O		
11.57	-oc NH	O O O	OC N S

11.58	-0¢ N	ه در چی جی گے۔ انگ	OC O
11.59	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph
11.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
11.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
11.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
11.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
11.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
11.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH2O(p-PhCH2COOH)
11.66	-0C 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

Table 12

Formula I : A = -B(OH)<sub>2</sub> ; X = -SC(=NH)NH<sub>2</sub> ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -CH<sub>2</sub>(p-PhOH).

	.1	.2	
12.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
12.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH2SPh
12.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
12.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
12.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
12.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
12.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
12.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
12.9	-C(O)o-hiphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
12.10	-C(O)m-hiphenyl	-C(O)CH2(m-hiphenyl)	-C(O)CH2CH2(m-biphenyl)
12.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
12.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
12.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
12.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
12.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
12.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
12.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
12.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
12.20	-C(O)m-PhSPh	-C(O)Cl12(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
12.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
12.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh

12.22	T C(C) Pt CH - SPI	Torrier	
12.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
12.24	C(O) - PLOU- OP	0.00011 ( p) 011 051	PhCH <sub>2</sub> SPh)
12.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	-C(O)CH2CH2(p-PhCH2SPh)
12.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
12.26	-C(O)cyclopentyl	-C(O)CH <sub>2</sub> (cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
12.27	-C(O)cyclohexyl	-C(O)CH <sub>2</sub> (cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
12.28	-C(0)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH <sub>2</sub> NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
12.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
12.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
12.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
12.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
12.33	-C(O)furan-2-yl	-C(O)CH <sub>2</sub> (furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
12.34	-C(O)furan-3-yl	-C(O)CH <sub>2</sub> (furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
12.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
12.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
12.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
12.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
12.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-vl)
12.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzoturan-2-yl)	-C(O)CH2CH2(benzofuran-2-
<u> </u>			yl)
12.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH2CH2(benzofuran-3-
10.10			yl)
12.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	•
l	ĺ	yl)	C(O)CH2CH2(benzothiophen-
12.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	2-yl)
12.44	-C(O)benzimidazo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
	-C(0/001121111102120-2-y)	-C(O)C.H2(helizimidi26-2-yi)	-C(O)CH2CH2(benzimidazo-
12.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	2-yl) -C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-
		C.(O/C412(1/C112(1/212(0-2-y1)	vl)
12.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo-2-
·			vl)
12.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
12.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-vl)	-C(O)Ph-4-(fluoren-9-vl)
12.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indôl-2-yl
12.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth-	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
10.51	2-vl)	·	, , , , , , , , , , , , , , , , , , , ,
12.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-yl-
12.52	C(())totaslandaria - tat.	(Ph)	4-(Ph)
12.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyi-2,2-(Ph <sub>2</sub> )
14.33	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoguinolin-3-	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-
12.54	-C(0)CH <sub>2</sub> (N-benzimidazol-2-	-C(U)CH <sub>2</sub> (N-benzoxazol-2-	yl)
.4.57		i -( (( ))( 'H2(/\/-henzavaza\-2-	-C(O)CH2(N-benzothiazol-2-
	-		
12.55	one)	one)	one)
12.55	-		

12.56	Co.	COO	
12.57	OC NH NH	0000	oc N s
12.58	-oc N	00 2 2 (2)	• • • • • • • • • • • • • • • • • • •
12.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph
12.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
12.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
12.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
12.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
12.64	-C(O)CH <sub>2</sub> O(o-PhCOOCH <sub>3</sub> )	-C(O)Cli2O(m-PhCOOCH3)	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
12.65	-C(O)CH <sub>2</sub> O(o-PhC11 <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
12.66	och		

Formula I : A = -B(OH)<sub>2</sub> : X = -SC(=NII)NII<sub>2</sub> :  $\mathbb{R}^3$  = table below :  $\mathbb{R}^{11}$  = -CH<sub>2</sub>CH<sub>2</sub>Ph.

	1	.2	.3
13.1	-C(O)Ph	-C(O)CII2Ph	-C(O)CH2CH2Ph
13.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH2SPh
13.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
13.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
13.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
13.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
13.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
13.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
13.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
13.10	-C(O)m-biphenyl	-C(O)CH2(m-hiphenyl)	-C(O)CH2CH2(m-biphenyl)
13.12	-C(O)p-hiphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
13.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)

13.14   -C(O)m-PhOPh   -C(O)CH2(n-PhOPh)   -C(O)CH2(H2(n-PhOPh)   13.15   -C(O)p-PhOPh   -C(O)CH2(n-PhOPh)   -C(O)CH2(H2(n-PhOPh)   -C(O)CH2(H2(n-PhOPh)   -C(O)CH2(H2(n-PhOPh)   -C(O)CH2(H2(n-PhOPh)   -C(O)CH2(H2(n-PhNHPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhSPh)   -C(O)CH2(H2(n-PhCH2SPh)   -C(O)CH2(H2(n-PhCH2S				
13.16	13.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
13.17	13.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
13.18	13.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
13.19	13.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
13.20	13.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
13.21   -C(O)p-PhSPh	13.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
13.22	13.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
13.23	13.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
13.24   C(O)p-PhCH <sub>2</sub> SPh   -C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)   -C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)   13.25   -C(O)adamantyl   -C(O)CH <sub>2</sub> (chalgenantyl)   -C(O)CH <sub>2</sub> (CH <sub>2</sub> (chepentyl)   -C(O)CH <sub>2</sub> (CH <sub>2</sub> (cyclohexyl)   -C(O)CH <sub>2</sub> (CH <sub>2</sub> (cyclohexyl)   -C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)   -C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)   -C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)   -C(O)CH <sub>2</sub> S(cyclohexyl)   -C(O)CH <sub>2</sub> S(cy	13.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
13.24   C(O)p-PhCH <sub>2</sub> SPh	13.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
13.25				
13.26	13.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	-C(O)CH2CH2(p-PhCH2SPh)
13.27   C(O)cyclohexyl	13.25	-C(O)adamantyl	-C(O)CH <sub>2</sub> (adamantyl)	-C(O)CH2CH2(adamantyl)
13.28	13.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
13.29	13.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
13.29	13.28		-C(O)CII2NII(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
13.31   -C(O)pyridin-3-yl   -C(O)CH2(pyridin-3-yl)   -C(O)CH2(pyridin-3-yl)   13.32   -C(O)pyridin-4-yl   -C(O)CH2(pyridin-4-yl)   -C(O)CH2(pyridin-3-yl)   -C(O)(pyridin-3-yl)   -C(O)(pyrid	13.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2N11(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
13.32	13.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
13.33   -C(O)Itran-2-yl   -C(O)CH2(Itran-2-yl)   -C(O)CH2(CH2(Itran-2-yl)   13.34   -C(O)Itran-3-yl   -C(O)CH2(Itran-3-yl)   -C(O)CH2(I	13.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
13.34   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2CH2(furan-3-yl)     13.35   -C(O)thiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2CH2(thiophen-2-yl)     13.36   -C(O)thiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2CH2(thiophen-2-yl)     13.37   -C(O)imidazo-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2CH2(thiophen-2-yl)     13.38   -C(O)oxazo-2-yl   -C(O)CH2(oxazo-2-yl)   -C(O)CH2CH2(imidazo-2-yl)     13.39   -C(O)thioazo-2-yl   -C(O)CH2(ch2(imidazo-2-yl)   -C(O)CH2CH2(thiozo-2-yl)     13.40   -C(O)benzofuran-2-yl   -C(O)CH2(benzofuran-2-yl)   -C(O)CH2CH2(benzofuran-2-yl)     13.41   -C(O)benzofuran-3-yl   -C(O)CH2(benzofuran-3-yl)   -C(O)CH2CH2(benzofuran-3-yl)     13.42   -C(O)benzothiophen-2-yl   -C(O)CH2(benzothiophen-2-yl)   -C(O)CH2(CH2(benzothiophen-2-yl)     13.43   -C(O)benzothiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2(CH2(thiophen-2-yl)     13.44   -C(O)benzothiazo-2-yl   -C(O)CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)     13.45   -C(O)benzothiazo-2-yl   -C(O)CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)     13.46   -C(O)benzothiazo-2-yl   -C(O)CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)     13.47   -C(O)benzothiazo-2-yl   -C(O)CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)     13.48   -C(O)Ph-2-(fluoren-9-yl)   -C(O)Ph-3-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-yl)     13.49   -C(O)W-indolin-2-one   -C(O)m-Ph(P(O)Ph3)   -C(O)P-Ph(P(O)Ph3)     13.49   -C(O)W-indolin-2-one   -C(O)indolin-2-yl   -C(O)indol-2-yl     13.50   -C(O)CCH3)2NIISO2(naphth-2-yl)   -C(O)cerabydrofuran-3-yl-4-(Ph)   -C(O)berabydrofuran-3-yl-4-(Ph)	13.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
13.35   -C(O)thiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2(CH2(thiophen-2-yl)   13.36   -C(O)thiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2(CH2(thiophen-2-yl)   -C(O)CH2(CH2(thiophen-2-yl)   -C(O)CH2(CH2(thiophen-2-yl)   -C(O)CH2(CH2(thiophen-2-yl)   -C(O)CH2(CH2(thiodazo-2-yl)   -C(O)CH2(CH2(t	13.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
13.36   -C(O)thiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2(thiophen-2-yl)   13.37   -C(O)imidazo-2-yl   -C(O)CH2(imidazo-2-yl)   -C(O)CH2(i	13.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
13.37   -C(O)imidazo-2-yl   -C(O)CH2(imidazo-2-yl)   -C(O)CH2(CH2(imidazo-2-yl)   13.38   -C(O)oxazo-2-yl   -C(O)CH2(oxazo-2-yl)   -C(O)CH2(CH2(cXazo-2-yl)   -C(O)CH2(CH2(CH2(cXazo-2-yl)   13.39   -C(O)thioazo-2-yl   -C(O)CH2(thioazo-2-yl)   -C(O)CH2(CH2(thioazo-2-yl)   -C(O)CH2(thioazo-2-yl)   -C(O)CH2(thioazo-2-yl	13.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
13.38   -C(O)oxazo-2-yl   -C(O)CH2(oxazo-2-yl)   -C(O)CH2(CH2(cxazo-2-yl)   13.39   -C(O)thioazo-2-yl   -C(O)CH2(thioazo-2-yl)   -C(O)CH2(CH2(thioazo-2-yl)   -C(O)CH2(CH2(thioazo-2-yl)   -C(O)CH2(CH2(thioazo-2-yl)   -C(O)CH2(CH2(thioazo-2-yl)   -C(O)CH2(CH2(thioazo-2-yl)   -C(O)CH2(CH2(thioazo-2-yl)   -C(O)CH2(CH2(thiopten-2-yl)   -C(O)CH2(thiopten-2-yl)   -C(O)CH2(thio	13.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
13.49   -C(O)thioazo-2-yl   -C(O)CH2(thioazo-2-yl)   -C(O)CH2CH2(thioazo-2-yl)   13.40   -C(O)benzofuran-2-yl   -C(O)CH2(benzofuran-2-yl)   -C(O)CH2(CH2(benzofuran-2-yl)   -C(O)CH2(CH2(benzofuran-2-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-2-yl)   -C(O)CH2(benzofuran-2-yl)   -C(O	13.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)Cl12CH2(imidazo-2-yl)
13.40   -C(O)benzofuran-2-yl   -C(O)CH2(benzofuran-2-yl)   -C(O)CH2(benzofuran-2-yl)   -C(O)CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-3-yl)   -C(O)CH2(CH2(benzofuran-2-yl)   -C(O)CH2(CH2(benzofuran-2-yl)   -C(O)CH2(CH2(benzofuran-2-yl)   -C(O)CH2(CH2(benzimidazo-2-yl)   -C(O)CH2(CH2(benzimidazo-2-yl)   -C(O)CH2(CH2(benzofuran-2-yl)   -C(O)CH	13.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
13.41	13.39	-C(O)thioazo-2-yl	-C(O)C112(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
13.41	13.40	-C(O)benzofuran-2-yl	-C(O)CI12(benzoturan-2-yl)	-C(O)CH2CH2(benzofuran-2-
13.42				
13.43   -C(O)thiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2CH2(thiophen-2-yl)   13.44   -C(O)benzimidazo-2-yl   -C(O)CH2(benzimidazo-2-yl)   -C(O)CH2CH2(benzimidazo-2-yl)   -C(O)Ph-4-(fluoren-9-yl)   -C(O)Ph-			-C(O)CH <sub>2</sub> (benzofuran-3-yl)	
13.43	13.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	-
13.43   -C(O)thiophen-2-yl   -C(O)CH2(thiophen-2-yl)   -C(O)CH2CH2(thiophen-2-yl)   13.44   -C(O)benzimidazo-2-yl   -C(O)CH2(benzimidazo-2-yl)   -C(O)CH2CH2(benzimidazo-2-yl)   -C(O)Ph-2-(fluoren-9-yl)   -C(O)Ph-3-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-yl)   -C(O)Ph-3-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-yl)   -C(O)CH2CH2(benzimidazo-2-yl)   -C(O)Ph-4-(fluoren-9-yl)   -C(O)Ph-3-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-			yl)	
13.44   -C(O)benzimidazo-2-yl   -C(O)CH2(benzimidazo-2-yl)   -C(O)CH2CH2(benzimidazo-2-yl)   13.45   -C(O)benzoxazo-2-yl   -C(O)CH2(benzoxazo-2-yl)   -C(O)CH2(benzoxazo-2-yl)   -C(O)CH2(benzoxazo-2-yl)   13.46   -C(O)benzothiazo-2-yl   -C(O)CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)   -C(O)CH2(CH2(benzothiazo-2-yl)   -C(O)Ph-2(fluoren-9-yl)   -C(O)Ph-3(fluoren-9-yl)   -C(O)Ph-3(fluoren-9-yl)   -C(O)Ph-3(fluoren-9-yl)   -C(O)Ph-4(fluoren-9-yl)   -C(O)indol-2-yl   -C(O)indol-2-yl   -C(O)cyclohexyl-2-(Ph)   -C(O)cyclohexyl-2-(Ph)   -C(O)cyclohexyl-2-(Ph)   -C(O)cyclohexyl-3-yl-4-(Ph)   -C(O)cyclohexyl-3	12.42			
13.45		<u> </u>		
13.46				2-yl)
13.47	13.45	-C(O)benzoxazı-2-yl	-C(O)CH2(benzoxazo-2-yl)	
13.47   -C(O)o-Ph(P(O)Ph3)   -C(O)m-Ph(P(O)Ph3)   -C(O)p-Ph(P(O)Ph3)     13.48   -C(O)Ph-2-(fluoren-9-yl)   -C(O)Ph-3-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-yl)     13.49   -C(O)N-indolin-2-one   -C(O)indolin-2-yl   -C(O)indol-2-yl     13.50   -C(O)C(CH3)2NIISO2(naphth-2-yl)   -C(O)cyclopentyl-2-(Ph)     2-yl)   -C(O)pyrrolidin-3-yl-4-(Ph)   -C(O)tetrahydrofuran-3-yl-4-(Ph)     13.51   -C(O)pyrrolidin-3-yl-4-(Ph)   -C(O)tetrahydrofuran-3-yl-4-(Ph)	13.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	
13.48   -C(O)Ph-2-(fluoren-9-yl)   -C(O)Ph-3-(fluoren-9-yl)   -C(O)Ph-4-(fluoren-9-yl)     13.49   -C(O)N-indolin-2-one   -C(O)indolin-2-yl   -C(O)indol-2-yl     13.50   -C(O)C(CH <sub>3</sub> )2NIISO <sub>2</sub> (naphth-2-yl)   -C(O)cyclopentyl-2-(Ph)   -C(O)cyclohexyl-2-(Ph)     13.51   -C(O)pyrrolidin-3-yl-4-(Ph)   -C(O)tetrahydrofuran-3-yl-4-(Ph)   -C(O)tetrahydrothiophen-3-yl-4-(Ph)	13.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	
13.49 -C(O) <i>N</i> -indolin-2-one -C(O)indolin-2-yl -C(O)indol-2-yl 13.50 -C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth-2-yl) -C(O)cyclopentyl-2-(Ph) -C(O)cyclohexyl-2-(Ph) 2-yl) -C(O)pyrrolidin-3-yl-4-(Ph) -C(O)tetrahydrofuran-3-yl-4-(Ph) -C(O)tetrahydrofuran-3-yl-4-(Ph)	13.48	-C(O)Ph-2-(fluoren-9-yl)		
2-ył)  13.51 -C(O)pyrrolidin-3-yl-4-(Ph) -C(O)tetrahydrofuran-3-yl-4 -C(O)tetrahydrothiophen-3-yl-4-(Ph)	13.49			-C(O)indol-2-yl
13.51 -C(O)pyrrolidin-3-yl-4-(Ph) -C(O)tetrahydrofuran-3-yl-4-(Ph) -C(O)tetrahydrothiophen-3-yl-4-(Ph)	13.50		-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
(Ph) 4-(Ph)				
13.52 -C(O)tetrahydronaphth-1-yl -C(O)tetrahydronaphth-2-yl -C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )			(Ph)	4-(Ph)
	13.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )

			C(O)OU.((2)i-dalia 2
13.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3-	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-
		yl12	yl) -C(O)CH <sub>2</sub> (N-benzothiazol-2-
13.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-	one)
	one)	one) -C(O)CH <sub>2</sub> (N-dihydrooxazol-2-	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2-
13.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol-		one)
	2-one)	one)	CO-
13.56	<b>CO-</b>	<b>℃</b> 00-	· (°)
, i	~N*O	N XV	N-V
	<b>  [</b> ] 人人		
	~ (()		
	~		
13.57	0	Ç	
	~N <sup>H</sup> NH	-00°N*0	-OC N S
1	-oc N NH		
13.58	00.0	-OC	-oc 0
13.50	-00 0		]; [[NN_/
		l	ا "ــر
		'N'	
			\ <b>_</b> /
1			
10.50	C(C)N(CN -) CH-Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
13.59	-C(O)N(CH3)CH2Ph	-C(O)Ph-3-(CH2(thiophen-2-	-C(O)Ph-3-(CH <sub>2</sub> Ph)
13.60	-C(O)pyridin-3-yl-5-(Ph)	yl))	(6), 2,5 (61,2,1,5)
13.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
13.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH <sub>2</sub> O(m-PhCH <sub>2</sub> OH)	-C(O)CH2O(p-PhCH2OH)
13.63	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
13.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
13.65	-C(O)CH2O(o-PhCH2COOH)	-C(O)C112O(m-	-C(O)CH2O(p-PhCH2COOH)
13.55		PhCH2COOH)	
13.66	-00 0		
	Till		1
1	- V 0		
	(	,	
1			
ł			<u> </u>

Table 14

Formula I :  $A = -B(OH)_2$ ;  $X = -SC(=NH)NH_2$ ;  $R^3 = table below$ ;  $R^{11} = -Ph$ .

	.1	.2	.3
14.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
14.2	-C(O)CII2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
14.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
14.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
14.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
14.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
14.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
14.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
14.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
14.10	-C(O)m-hiphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
14.12	-C(O)p-biphenyl	-C(O)Cl·l <sub>2</sub> (p-hiphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
14.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
14.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
14.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
14.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
14.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhNHPh)
14.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
14.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
14.20	-C(O)m-PhSPh	-C(O)CH <sub>2</sub> (m-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhSPh)
14.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
14.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(0- PhCH2SPh)
14.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m- PhCH <sub>2</sub> SPh)
14.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p- PhCH <sub>2</sub> SPh)
14.25	-C(O)adamantyl	-C(O)CI-i2(adamantyl)	-C(O)CH2CH2(adamantyl)
14.26	-C(O)cyclopentyl	-C(O)CH <sub>2</sub> (cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
14.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
14.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
14.29	-C(O)CH2O(cyclohexyl)	-C(O)CII2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
14.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
14.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
14.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
14.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
14.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
14.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
14.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
14.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
14.38	-C(O)oxazo-2-yl	-C(O)CH <sub>2</sub> (oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
14.39	-C(O)thioazo-2-yl	-C(O)Cll2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
14.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzoturan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2-yl)

14.42			·	
14.43	14.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
14.43	14.42	-C(O)benzothiophen-2-yl	-C(O)Cl12(benzothiophen-2-yl)	-2-yl)
14.44	14.43	-C(O)thiophen-2-vl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
14.46	14.44		-C(O)CH2(benzimidazo-2-yl)	2-v1)
14.47	14.45	-C(O)benzoxazo-2-yl		yl)
14.57	14.46	-C(O)benzothiazo-2-yl	-C(O)Cl12(benzothiazo-2-yl)	2-vl)
14.48	14.47	-C(O)o-Ph(P(O)Ph3)		
14.49   -C(O)N-indolin-2-one   -C(O)indolin-2-yl   -C(O)indolin-2-yl   -C(O)indolin-2-yl   -C(O)indolin-2-yl   -C(O)indolin-2-yl   -C(O)indolin-2-yl   -C(O)indolin-3-yl-4-(Ph)   -C(O)indolin-3-yl-3-(O)indolin				
14.50		-C(O)N-indolin-2-one		
14.51	14.50	-C(O)C(CH3)2NHSO2(naphth-	•	
14.53   -C(O)CH2(W-benzimidazol-2-one)   -C(O)CH2(W-benzimidazol-2-one)   -C(O)CH2(W-benzimidazol-2-one)   -C(O)CH2(W-benzimidazol-2-one)   -C(O)CH2(W-dihydroinidazol-2-one)   -C(O)CH2(W-dihydroinidazol-2-one	14.51			4-(Ph)
14.53	14.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphti-2-yl	
14.55	14.53	-C(O)tetrahydroisoquinolin-1-yl		yl)
14.55	14.54	1		one)
14.57 -OC N N H -OC N N O	14.55	-C(O)CH2(N-dihydroimidazol-		_
-OC N NH -OC N S  -OC	14.56	C N C O	C N CO.	
14.59 -C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph 14.60 -C(O)pyridin-3-yl-5-(Ph) -C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl)) -C(O)Ph-3-(CH <sub>2</sub> Ph) 14.61 -C(O)C(CH <sub>3</sub> )2OPh -C(O)CH(C <sub>2</sub> H <sub>5</sub> )OPh -C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH) -C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH) 14.62 -C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> OH) -C(O)CH <sub>2</sub> O(m-PhCH <sub>2</sub> OH) -C(O)CH <sub>2</sub> O(p-PhCOOH) 14.63 -C(O)CH <sub>2</sub> O(o-PhCOOH) -C(O)CH <sub>2</sub> O(m-PhCOOH) -C(O)CH <sub>2</sub> O(p-PhCOOH) 14.64 -C(O)CH <sub>2</sub> O(o-PhCOOCH <sub>3</sub> ) -C(O)CH <sub>2</sub> O(m-PhCOOCH <sub>3</sub> ) -C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )	14.57	-0C N N H	-00°N	-oc N S
14.60         -C(O)pyridin-3-yl-5-(Ph)         -C(O)Ph-3-(CH2(thiophen-2-yl))         -C(O)Ph-3-(CH2Ph)           14.61         -C(O)C(CH3)2OPh         -C(O)CH(C2H5)OPh         -C(O)CH2OCH2Ph           14.62         -C(O)CH2O(o-PhCH2OH)         -C(O)CH2O(m-PhCH2OH)         -C(O)CH2O(m-PhCOOH)           14.63         -C(O)CH2O(o-PhCOOH)         -C(O)CH2O(m-PhCOOH)         -C(O)CH2O(p-PhCOOCH3)           14.64         -C(O)CH2O(o-PhCOOCH3)         -C(O)CH2O(m-PhCOOCH3)         -C(O)CH2O(p-PhCOOCH3)	14.58	N CO	-00 Z Z -	N
14.60         -C(O)pyridin-3-yl-5-(Ph)         -C(O)Ph-3-(CH2(thiophen-2-yl))         -C(O)Ph-3-(CH2Ph)           14.61         -C(O)C(CH3)2OPh         -C(O)CH2(C2H5)OPh         -C(O)CH2OCH2Ph           14.62         -C(O)CH2O(o-PhCH2OH)         -C(O)CH2(Cm-PhCH2OH)         -C(O)CH2O(p-PhCH2OH)           14.63         -C(O)CH2O(o-PhCOOH)         -C(O)CH2O(m-PhCOOH)         -C(O)CH2O(p-PhCOOCH3)           14.64         -C(O)CH2O(o-PhCOOCH3)         -C(O)CH2O(m-PhCOOCH3)         -C(O)CH2O(p-PhCOOCH3)	14.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	
14.61         -C(O)C(CH3)2OPh         -C(O)CH(C2H5)OPh         -C(O)CH2OCH2Ph           14.62         -C(O)CH2O(o-PhCH2OH)         -C(O)CH2O(m-PhCH2OH)         -C(O)CH2O(m-PhCOOH)           14.63         -C(O)CH2O(o-PhCOOH)         -C(O)CH2O(m-PhCOOH)         -C(O)CH2O(p-PhCOOCH3)           14.64         -C(O)CH2O(o-PhCOOCH3)         -C(O)CH2O(m-PhCOOCH3)         -C(O)CH2O(p-PhCOOCH3)				
14.62         -C(O)CH2O(o-PhCH2OH)         -C(O)CH2O(m-PhCH2OH)         -C(O)CH2O(p-PhCH2OH)           14.63         -C(O)CH2O(o-PhCOOH)         -C(O)CH2O(m-PhCOOH)         -C(O)CH2O(p-PhCOOH)           14.64         -C(O)CH2O(o-PhCOOCH3)         -C(O)CH2O(m-PhCOOCH3)         -C(O)CH2O(p-PhCOOCH3)				-C(O)CH2OCH2Ph
14.63         -C(O)CH2O(o-PhCOOH)         -C(O)CH2O(m-PhCOOH)         -C(O)CH2O(p-PhCOOH)           14.64         -C(O)CH2O(o-PhCOOCH3)         -C(O)CH2O(m-PhCOOCH3)         -C(O)CH2O(p-PhCOOCH3)				
14.64 -C(O)CII2O(n-PhCOOCH3) -C(O)CII2O(m-PhCOOCH3) -C(O)CH2O(p-PhCOOCH3)				
14.04   -C(O)C1170(0-110:000:117)   C(O)C1120(110:000:117)			-C(O)CII2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
	14.65	-C(O)CH2O(o-PhCH2COOH)	-C(O)CH2O(m-PhCH2COOH)	-C(O)CH2O(p-PhCH2COOH

14.66	-00 0	` .	
	r <sup>N</sup> NO		
	آسر		

Table 15

Formula I : A = -B(OH)<sub>2</sub> ; X = -SC(=NH)NH<sub>2</sub> ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -CH<sub>2</sub>(naphth-2-yl).

15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-2-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)			<u> </u>	· · · · · · · · · · · · · · · · · · ·
15.2	•			. I
15.3	15.1	-C(O)Ph		
15.4         -C(O)o-PhCH2OH         -C(O)m-PhCOOH         -C(O)m-PhCOOH         -C(O)m-PhCOOH         -C(O)m-PhCOOH         -C(O)m-PhCH2COOH         -C(O)m-PhCH2CDOH         -C(O)m-PhCH2CDOH         -C(O)m-PhCH2CDOH         -C(O)m-PhCH2CDOH         -C(O)m-PhCH2CDOH         -C(O)CH2(m-PhCH2)         -C(O)CH2CH2(maphh-1-yl)         -C(O)CH2CH2(maphh-2-yl)         -C(O)CH2CH2(maphh-2-yl)         -C(O)CH2CH2(maphh-2-yl)         -C(O)CH2CH2(m-biphenyl)				<u> </u>
15.5				
15.6		-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	
15.7				
15.8		-C(O)o-PhCH2COOH		
15.9		-C(O)naphth-1-yl	-C(O)Cl12(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
15.10		-C(O)naphth-2-yl		-C(O)CH2CH2(napth-2-yl)
15.12		-C(O)o-hiphenyl	-C(O)CH <sub>2</sub> (o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
15.13		-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
15.14		-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
15.15         -C(O)p-PhOPh         -C(O)CH2(p-PhOPh)         -C(O)CH2(H2(p-PhOPh)           15.16         -C(O)o-PhNHPh         -C(O)CII2(o-PhNHPh)         -C(O)CH2(H2(p-PhNHPh)           15.17         -C(O)m-PhNHPh         -C(O)CH2(m-PhNHPh)         -C(O)CH2(H2(m-PhNHPh)           15.18         -C(O)p-PhNHPh         -C(O)CH2(p-PhNHPh)         -C(O)CH2(H2(p-PhNHPh)           15.19         -C(O)o-PhSPh         -C(O)CH2(p-PhNHPh)         -C(O)CH2(H2(p-PhSPh)           15.20         -C(O)m-PhSPh         -C(O)CH2(m-PhSPh)         -C(O)CH2(H2(m-PhSPh)           15.21         -C(O)m-PhSPh         -C(O)CH2(p-PhSPh)         -C(O)CH2(H2(p-PhSPh)           15.22         -C(O)a-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2(H2(p-PhCH2SPh)           15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2CH2(m-PhCH2SPh)           15.24         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(H2(p-PhCH2SPh)           15.25         -C(O)m-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2(H2(p-PhCH2SPh)           15.26         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2(P2(predentyl)         -C(O)CH2(P2(predentyl)           15.26         -C(O)adamantyl         -C(O)CH2(clohexyl)         -C(O)CH2(p2(clohexyl)         -C(O)CH2(p2(clohexyl)           15		<u> </u>	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
15.16         -C(O)c-PhNHPh         -C(O)CI12(o-PhNHPh)         -C(O)CH2(H2(o-PhNHPh)           15.17         -C(O)m-PhNHPh         -C(O)CH2(m-PhNHPh)         -C(O)CH2(H2(m-PhNHPh)           15.18         -C(O)p-PhNHPh         -C(O)CH2(p-PhNHPh)         -C(O)CH2(H2(p-PhNHPh)           15.19         -C(O)o-PhSPh         -C(O)CH2(p-PhNHPh)         -C(O)CH2(H2(p-PhSPh)           15.20         -C(O)m-PhSPh         -C(O)CH2(m-PhSPh)         -C(O)CH2(H2(m-PhSPh)           15.21         -C(O)p-PhSPh         -C(O)CH2(p-PhSPh)         -C(O)CH2(H2(p-PhSPh)           15.22         -C(O)a-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2(H2(p-PhCH2SPh)           15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(H2(m-PhCH2SPh)           15.24         -C(O)p-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(H2(p-PhCH2SPh)           15.25         -C(O)a-PhCH2SPh         -C(O)CH2(Pp-PhCH2SPh)         -C(O)CH2(H2(p-PhCH2SPh)           15.26         -C(O)a-PhCH2SPh         -C(O)CH2(adamantyl)         -C(O)CH2(H2(p-PhCH2SPh)           15.26         -C(O)a-PhCH2SPh         -C(O)CH2(cyclopentyl)         -C(O)CH2(Cyclopentyl)           15.27         -C(O)a-PhCH2(pyridin-PhCH2SPh)         -C(O)CH2(PCH2(cyclopentyl)         -C(O)CH2(PCH2(cyclopentyl)           15.28         -C(O)CH2(O(cyclope	15.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
15.17         -C(O)m-PhNHPh         -C(O)CH2(m-PhNHPh)         -C(O)CH2(H2(m-PhNHPh)           15.18         -C(O)p-PhNHPh         -C(O)CH2(p-PhNHPh)         -C(O)CH2(H2(H2(p-PhNHPh))           15.19         -C(O)o-PhSPh         -C(O)CH2(p-PhSPh)         -C(O)CH2(H2(P-PhSPh))           15.20         -C(O)m-PhSPh         -C(O)CH2(m-PhSPh)         -C(O)CH2(H2(H2(p-PhSPh))           15.21         -C(O)p-PhSPh         -C(O)CH2(p-PhSPh)         -C(O)CH2(H2(P-PhSPh))           15.22         -C(O)o-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(H2(m-PhCH2SPh))           15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(H2(m-PhCH2SPh))           15.24         -C(O)p-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(H2(m-PhCH2SPh))           15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2CH2(p-PhCH2SPh)           15.26         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2CH2(adamantyl)           15.27         -C(O)adamantyl         -C(O)CH2(cyclopentyl)         -C(O)CH2(cyclopentyl)           15.28         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2(Cyclopentyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2(D(Cyclopentyl)         -C(O)CH2(Cyclopentyl)           15.29         -C(O)CH2O(cyclopentyl)<			-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
15.18         -C(O)p-PhNHPh         -C(O)CH2(p-PhNIIPh)         -C(O)CH2(P-PhNHPh)           15.19         -C(O)o-PhSPh         -C(O)CH2(o-PhSPh)         -C(O)CH2CH2(o-PhSPh)           15.20         -C(O)m-PhSPh         -C(O)CH2(m-PhSPh)         -C(O)CH2CH2(m-PhSPh)           15.21         -C(O)p-PhSPh         -C(O)CH2(p-PhSPh)         -C(O)CH2CH2(p-PhSPh)           15.22         -C(O)o-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2CH2(m-PhCH2SPh)           15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2CH2(m-PhCH2SPh)           15.24         -C(O)p-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2CH2(m-PhCH2SPh)           15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2CH2(p-PhCH2SPh)           15.26         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2CH2(adamantyl)           15.27         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2CH2((cyclopentyl)           15.28         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2(cyclopentyl)         -C(O)CH2CH2(cyclopentyl)           15.29         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclopentyl)         -C(O)CH2C(cyclopentyl)         -C(O)CH2C(cyclopentyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2(pyridin-3-yl) <td></td> <td>-C(O)o-PhNHPh</td> <td>-C(O)CII<sub>2</sub>(o-PhNHPh)</td> <td>-C(O)CH2CH2(o-PhNHPh)</td>		-C(O)o-PhNHPh	-C(O)CII <sub>2</sub> (o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
15.19         -C(O)o-PhSPh         -C(O)CH2(o-PhSPh)         -C(O)CH2(d-PhSPh)           15.20         -C(O)m-PhSPh         -C(O)CH2(m-PhSPh)         -C(O)CH2(d-PhSPh)           15.21         -C(O)p-PhSPh         -C(O)CH2(m-PhSPh)         -C(O)CH2(d-PhSPh)           15.22         -C(O)o-PhCH2SPh         -C(O)CH2(d-PhCH2SPh)         -C(O)CH2(d-PhSPh)           15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(d-PhCH2SPh)           15.24         -C(O)p-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(d-PhCH2SPh)           15.25         -C(O)p-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(d-PhCH2SPh)           15.26         -C(O)p-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(d-PhCH2SPh)           15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2(d-PhCH2SPh)           15.26         -C(O)cyclopentyl         -C(O)CH2(adamantyl)         -C(O)CH2(cyclopentyl)           15.27         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2(cyclopentyl)         -C(O)CH2(cyclopentyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclopentyl)         -C(O)CH2(cyclopentyl)         -C(O)CH2(cyclopentyl)           15.29         -C(O)CH2O(cyclopentyl)         -C(O)CH2(pyridin-2-yl)         -C(O)CH2(pyridin-2-yl)         -C(O)CH		-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
15.20   -C(O)m-PhSPh   -C(O)CH2(m-PhSPh)   -C(O)CH2(m-PhSPh)     15.21   -C(O)p-PhSPh   -C(O)CH2(p-PhSPh)   -C(O)CH2(p-PhSPh)     15.22   -C(O)o-PhCH2SPh   -C(O)CH2(o-PhCH2SPh)   -C(O)CH2CH2(o-PhCH2SPh)     15.23   -C(O)m-PhCH2SPh   -C(O)CH2(m-PhCH2SPh)   -C(O)CH2CH2(m-PhCH2SPh)     15.24   -C(O)p-PhCH2SPh   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2CH2(p-PhCH2SPh)     15.25   -C(O)adamantyl   -C(O)CH2(adamantyl)   -C(O)CH2CH2(adamantyl)     15.26   -C(O)cyclopentyl   -C(O)CH2(cyclopentyl)   -C(O)CH2CH2((cyclopentyl)     15.27   -C(O)cyclohexyl   -C(O)CH2(cyclohexyl)   -C(O)CH2CH2(cyclohexyl)     15.28   -C(O)CH2O(cyclopentyl)   -C(O)CH2NH(cyclohexyl)   -C(O)CH2S(cyclohexyl)     15.29   -C(O)CH2O(cyclohexyl)   -C(O)CH2NH(cyclohexyl)   -C(O)CH2S(cyclohexyl)     15.30   -C(O)pyridin-2-yl   -C(O)CH2(pyridin-2-yl)   -C(O)CH2CH2(pyridin-2-yl)     15.31   -C(O)pyridin-3-yl   -C(O)CH2(pyridin-3-yl)   -C(O)CH2CH2(pyridin-3-yl)     15.33   -C(O)furan-2-yl   -C(O)CH2(furan-2-yl)   -C(O)CH2CH2(furan-2-yl)     15.34   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2CH2(furan-3-yl)     15.34   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2(furan-3-yl)     15.35   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2(furan-3-yl)     15.36   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2(furan-3-yl)     15.37   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2(furan-3-yl)     15.38   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2(furan-3-yl)     15.39   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2(furan-3-yl)     15.30   -C(O)furan-3-yl   -C(O)CH2(furan-3-yl)   -C(O)CH2(furan-3-yl)     15		-C(O)p-PhNHPh	-C(O)CH <sub>2</sub> (p-PhN11Ph)	-C(O)CH2CH2(p-PhNHPh)
15.21         -C(O)p-PhSPh         -C(O)CH2(p-PhSPh)         -C(O)CH2(p-PhSPh)           15.22         -C(O)o-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2(p-PhCH2SPh)           15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(p-PhCH2SPh)           15.24         -C(O)p-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2(p-PhCH2SPh)           15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2(p-PhCH2SPh)           15.26         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2(cyclopentyl)           15.27         -C(O)cyclohexyl         -C(O)CH2(cyclohexyl)         -C(O)CH2CH2(cyclopentyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclopentyl)         -C(O)CH2S(cyclopentyl)           15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)CH2O(cyclohexyl)         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-3-yl)           15.34 <td></td> <td>-C(O)o-PhSPh</td> <td>-C(O)CH2(o-PhSPh)</td> <td>-C(O)CH2CH2(o-PhSPh)</td>		-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
15.22         -C(O)o-PhCH2SPh         -C(O)CH2(o-PhCH2SPh)         -C(O)CH2(G-PhCH2SPh)           15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2(M-PhCH2SPh)           15.24         -C(O)p-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2CH2(p-PhCH2SPh)           15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2CH2(adamantyl)           15.26         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2CH2((cyclopentyl))           15.27         -C(O)cyclohexyl         -C(O)CH2(cyclohexyl)         -C(O)CH2CH2(cyclohexyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)CH2O(cyclohexyl)         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2(pyridin-4-yl)         -C(O)CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2(furan-3-yl)         -C(O)CH2(furan-3-yl)	15.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
PhCH2SPh   -C(O)CH2(m-PhCH2SPh   -C(O)CH2(m-PhCH2SPh)   -C(O)CH2(CH2(m-PhCH2SPh)   -C(O)CH2CH2(m-PhCH2SPh)   -C(O)CH2CH2(p-PhCH2SPh)   -C(O)CH2CH2(p-PhCH2SPh)   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2(p-PhCH2SPh)   -C(O)CH2CH2(p-PhCH2SPh)   -C(O)C		-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
15.23         -C(O)m-PhCH2SPh         -C(O)CH2(m-PhCH2SPh)         -C(O)CH2CH2(m-PhCH2SPh)           15.24         -C(O)p-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2CH2(p-PhCH2SPh)           15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2CH2(adamantyl)           15.26         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2CH2((cyclopentyl)           15.27         -C(O)cyclohexyl         -C(O)CH2(cyclohexyl)         -C(O)CH2CH2(cyclohexyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2(pyridin-4-yl)         -C(O)CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2(furan-3-yl)         -C(O)CH2(furan-3-yl)	15.22	-C(O)o-PhCl12SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
PhCH2SPh				<u> </u>
15.24         -C(O)p-PhCH2SPh         -C(O)CH2(p-PhCH2SPh)         -C(O)CH2(p-PhCH2SPh)           15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2(CH2(adamantyl))           15.26         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2CH2((cyclopentyl))           15.27         -C(O)cyclohexyl         -C(O)CH2(cyclohexyl)         -C(O)CH2CH2(cyclohexyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(fyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-2-yl)           15.34         -C(O)turan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)	15.23	-C(O)m-PhCH2SPh	-C(O)CH <sub>2</sub> (m-PhCH <sub>2</sub> SPh)	
PhCH2SPh   15.25   -C(O)adamantyl   -C(O)CH2(adamantyl)   -C(O)CH2(CH2(adamantyl)   -C(O)CH2(CH2(adamantyl)   -C(O)CH2(CH2(adamantyl)   -C(O)CH2(CH2(Cyclopentyl)   -C(O)CH2CH2((cyclopentyl)   -C(O)CH2CH2((cyclopentyl)   -C(O)CH2CH2(cyclopentyl)   -C(O)CH2CH2(cyclopentyl)   -C(O)CH2CH2(cyclopentyl)   -C(O)CH2CH2(cyclopentyl)   -C(O)CH2S(cyclopentyl)   -C(O)CH2S(cyclopentyl)   -C(O)CH2S(cyclopentyl)   -C(O)CH2S(cyclopentyl)   -C(O)CH2CH2(pyridin-2-yl)   -C(O)CH2CH2(pyridin-2-yl)   -C(O)CH2CH2(pyridin-3-yl)   -C(O)CH2CH2(pyridin-3-yl)   -C(O)CH2CH2(pyridin-3-yl)   -C(O)CH2CH2(pyridin-4-yl)   -C(O)CH2CH2(pyridin-				<del></del>
15.25         -C(O)adamantyl         -C(O)CH2(adamantyl)         -C(O)CH2(adamantyl)           15.26         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2CH2((cyclopentyl))           15.27         -C(O)cyclohexyl         -C(O)CH2(cyclohexyl)         -C(O)CH2CH2(cyclohexyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclopentyl)         -C(O)CH2S(cyclopentyl)           15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-3-yl)           15.34         -C(O)turan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)	15.24	-C(O)p-PhCH2SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	
15.26         -C(O)cyclopentyl         -C(O)CH2(cyclopentyl)         -C(O)CH2(l(cyclopentyl))           15.27         -C(O)cyclohexyl         -C(O)CH2(cyclohexyl)         -C(O)CH2(l(cyclohexyl))           15.28         -C(O)CH2(O(cyclopentyl))         -C(O)CH2NH(cyclopentyl)         -C(O)CH2S(cyclopentyl)           15.29         -C(O)CH2(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-3-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)			<u> </u>	
15.27         -C(O)cyclohexyl         -C(O)Cl12(cyclohexyl)         -C(O)CH2CH2(cyclohexyl)           15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclopentyl)         -C(O)CH2S(cyclopentyl)           15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-2-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)			<del></del>	-C(O)CH2CH2(adamantyl)
15.28         -C(O)CH2O(cyclopentyl)         -C(O)CH2NH(cyclopentyl)         -C(O)CH2S(cyclopentyl)           15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-2-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)		-C(O)cyclopentyl	-C(O)CH <sub>2</sub> (cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
15.29         -C(O)CH2O(cyclohexyl)         -C(O)CH2NH(cyclohexyl)         -C(O)CH2S(cyclohexyl)           15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-2-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)		<u> </u>	_ <u></u>	-C(O)CH2CH2(cyclohexyl)
15.30         -C(O)pyridin-2-yl         -C(O)CH2(pyridin-2-yl)         -C(O)CH2CH2(pyridin-2-yl)           15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2CH2(furan-2-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2CH2(furan-3-yl)			-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
15.31         -C(O)pyridin-3-yl         -C(O)CH2(pyridin-3-yl)         -C(O)CH2CH2(pyridin-3-yl)           15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2CH2(pyridin-4-yl)           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2(furan-2-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2(furan-3-yl)	15.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
15.32         -C(O)pyridin-4-yl         -C(O)CH2(pyridin-4-yl)         -C(O)CH2(Ed2(pyridin-4-yl))           15.33         -C(O)furan-2-yl         -C(O)CH2(furan-2-yl)         -C(O)CH2(furan-2-yl)           15.34         -C(O)furan-3-yl         -C(O)CH2(furan-3-yl)         -C(O)CH2(furan-3-yl)	15.30		-C(O)CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
15.33 -C(O)furan-2-yl -C(O)CH2(furan-2-yl) -C(O)CH2CH2(furan-2-yl) 15.34 -C(O)furan-3-yl -C(O)CH2(furan-3-yl) -C(O)CH2(furan-3-yl)	15.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
15.34 -C(O)furan-3-yl -C(O)CH2(furan-3-yl) -C(O)CH2CH2(furan-3-yl)	15.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
	15.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
15.35 -C(O)thiophen-2-yl -C(O)CH2(thiophen-2-yl) -C(O)CH2CH2(thiophen-2-yl)	15.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
	15.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)

15.36				5.5.51.51.41.
15.38   C(O)oxazo-2-yl   -C(O)CH2(bioazo-2-yl)   -C(O)CH2(hlioazo-2-yl)   -C(O)CH2(hlioazo-2-y	15.36	-C(O)thiophen-2-yl		-C(O)CH2CH2(thiophen-2-yl)
15.39   C(O)thiotazo-2-yl   -C(O)CH2(thiotazo-2-yl)   -C(O)CH2(thiot	15.37	-C(O)imidazo-2-yl		
15.39	15.38	-C(O)oxazo-2-yl	-C(O)CH <sub>2</sub> (oxazo-2-yl)	
15.40		-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	
15.41   -C(O)benzothiana-3-yl   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(CH2(benzothiana-3-yl)   -C(O)CH2(CH2(benzothiana-3-yl)   -C(O)CH2(CH2(benzothiana-3-yl)   -C(O)CH2(CH2(benzothiana-3-yl)   -C(O)CH2(CH2(benzothiana-3-yl)   -C(O)CH2(CH2(benzothiana-2-yl)   -C(O)CH2(CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C(O)CH2(benzothiana-3-yl)   -C			-C(O)CH <sub>2</sub> (benzofuran-2-yl)	
15.43	15.41	-C(O)benzofuran-3-yl		-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-
15.49   C(O)benzimidazo-2-yl   -C(O)CH2(Denzimidazo-2-yl)   -C(O)CH2(Denzimidazo-2-yl)   -C(O)CH2(Denzoxazo-2-yl)   -C(O)Ph-2-(Iluoren-9-yl)   -C(O)Ph-3-(Iluoren-9-yl)   -C(O)Ph-3-(Iluoren-9-yl)   -C(O)Ph-4-(Iluoren-9-yl)   -C(O)Ph-4-(Iluoren-9-yl)   -C(O)Ph-4-(Iluoren-9-yl)   -C(O)C(CH3)ZNHSO2(naphth-2-yl)   -C(O)C(CH3)ZNHSO2(naphth-2-yl)   -C(O)C(Denzhydroixaphth-2-yl)   -C(O)C(Denzhydroixaphth-3-yl-4-(Ph)   -C(O)Cenzhydroixaphth-3-yl-4-(Ph)   -C(O)Cenzhydroixaphth-3-yl-4-(Ph)   -C(O)Cenzhydroixaphth-3-yl-4-(Ph)   -C(O)CH2(N-benzoxazol-2-one)   -C(O)CH2(N-benzoxiazol-2-one)	15.42	-C(O)benzothiophen-2-yl	-C(O)CII <sub>2</sub> (benzothiophen-2-yl)	-2-yl)
15.44	15.43	-C(O)thiophen-2-vl	-C(O)CH2(thiophen-2-yl)	
15.45			-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	2-yl)
15.47	15.45	-C(O)benzoxazo-2-yl		-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
15.48	15.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	2-yl)
15.48	15.47	-C(O)o-Ph(P(O)Ph3)		
15.49				
15.50		-C(O)N-indolin-2-one		
15.51		-C(O)C(CH3)2NHSO2(naphth-		-C(O)cyclohexyl-2-(Ph)
15.53 -C(O)tetrahydroisoquinolin-1-yl -C(O)tetrahydroisoquinolin-3-yl -C(O)CH2((2-oxo)indolin-3-yl)  15.54 -C(O)CH2(N-benzimidazol-2-one) -C(O)CH2(N-benzoxtazol-2-one)  15.55 -C(O)CH2(N-dihydroimidazol-2-one) -C(O)CH2(N-dihydrooxazol-2-one)  15.56 -CO  CO  N  O  O  N  O  O  N  O  O  N  O  O  N  O  O	15.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-(Ph)	4-(Ph)
15.53   -C(O)Retrahydroisoquinolin-1-yl   -C(O)Retrahydroisoquinolin-3-yl   -C(O)CH2((2-oxo)indolin-3-yl)     15.54   -C(O)CH2((N-benzimidazol-2-one)   -C(O)CH2((N-benzoxazol-2-one)     15.55   -C(O)CH2((N-dihydroimidazol-2-one)   -C(O)CH2((N-dihydroimidazol-2-one)     15.56   CO-	15.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
15.54   -C(O)CH <sub>2</sub> (N-benzimidazol-2-one)   -C(O)CH <sub>2</sub> (N-benzothiazol-2-one)   -C(O)CH <sub>2</sub> (N-benzothiazol-2-one)   -C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)   -C(O)CH			-C(O)tetrahydroisoquinolin-3-yl	<u> </u>
15.55 -C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one) -C(O)CH <sub>2</sub> (N-dihydrooxazol-2- one) -C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one) -C(O)CH <sub>2</sub> (N-dihydrothia	15.54		-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	one)
15.56  15.57  -OC N N O -OC N O -OC N O -OC N O OC N O OC N OC N	15.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol-	,	
15.58 -OC O -OC NHO -OC NHS  15.59 -C(O)N(CH3)CH2Ph -C(O)N(C2H5)CH2Ph -C(O)N(C3H7)CH2Ph	15.56	<del></del>	Q N COO	
15.59 -C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph	15.57	-30 13	·oc N o	-0c N S
1911.5 64.07.46.2.17.6.2	15.58	-oc o		-oc o
1911.5 64.07.46.2.17.6.2	15.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph

15.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
15.62	-C(O)CH2O(o-PhCH2OH)	-C(O)Cl12O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
15.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
15.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
15.65	-C(O)CH2O(o-PhCH2COOH)	-C(O)CH2O(m-PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
15.66	-00 0		
	l 'N <sup>N</sup> O		
	سر ا	I	
			· ·
l	(_)		

Table 16

Formula I : A = -B(pinanediol) ; X = guanidinyl ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  =  $\mathbb{C}\mathbb{H}_3$ 

	.1	.2	.3
16.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
16.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH2SPh
16.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
16.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
16.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
16.6	-C(O)o-PhCII2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
16.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
16.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
16.9	-C(O)o-biphenyl	-C(O)CH <sub>2</sub> (o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
16.10	-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-hiphenyl)	-C(O)CH2CH2(m-biphenyl)
16.12	-C(O)p-biphenyl	-C(O)Cl12(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
16.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
16.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
16.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
16.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
16.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
16.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
16.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
16.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
16.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
16.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (o-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o- PhCH <sub>2</sub> SPh)
16.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (m-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m- PhCH <sub>2</sub> SPh)
16.24	-C(O)p-PhCH2SPh	-C(U)CH2(p-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p- PhCH <sub>2</sub> SPh)
16.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
16.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
16.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
16.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
16.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
16.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)

			(C)
16.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
16.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
16.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
16.34	-C(O)furan-3-yl	-C(O)CH <sub>2</sub> (furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
16.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
16.36	-C(O)thiophen-2-yl	-C(O)CII2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
16.37	-C(O)imidazo-2-yl	-C(O)CH <sub>2</sub> (imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
16.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
16.39	-C(O)thioazo-2-yl	-C(O)CH <sub>2</sub> (thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
16.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2-yl)
16.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
16.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-vl)
16.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
16.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
16.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
16.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-yl)
16.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
16.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
16.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indol-2-yl
16.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
16.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-(Ph)	-C(O)tetrahydrothiophen-3-yl-4-(Ph)
16.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
16.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)terrahydroisoquinolin-3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- yl)
16.54	-C(O)CH <sub>2</sub> ( <i>N</i> -benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)Cll <sub>2</sub> (N-benzothiazol-2-one)
16.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
16.56		Checo-	
16.57	-oc N NH	-oc N 0	-oc N s

16.58	-0ç <u>0</u>	OC 2 2 (2)	-0C N O O
16.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph
16.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
16.61	-C(O)C(CH3)2OPh	-C(O)CH(C <sub>2</sub> H <sub>5</sub> )OPh	-C(O)CH2OCH2Ph
16.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
16.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
16.64	-C(O)CH2O(o-PhCOOCH3)	-C(U)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
16.65	-C(O)CH2O(n-PhCH2COOH)	-C(O)CH2O(m-PhCH2COOH)	-C(O)CH2O(p-PhCH2COOH)
16.66	0 × 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

<u>Table 17</u>.

Formula I : A = -B(pinanediol) ; X = guanidinyl ;  $R^3$  = table below ;  $R^{11}$  = -CH2(p-PhOH).

	.1	.2	
17.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
17.2	-C(O)CH <sub>2</sub> OPh	-C(O)CH2NHPh	-C(O)CH2SPh
17.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
17.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
17.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
17.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
17.7	-C(O)naphth-1-yl	-C(O)Cll2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
17.8	-C(O)naphth-2-yl	-C(O)Cll2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
17.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
17.10	-C(O)m-hiphenyl	-C(O)Cl12(m-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
17.12	-C(O)p-hiphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
17.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
17.14	-C(O)m-PhOPh	-C(O)Cl12(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
17.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
17.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
17.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
17.18	-C(O)p-PhNHPh	-C(O)CH <sub>2</sub> (p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
17.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
17.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
17.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
17.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o- PhCH2SPh)

17.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m- PhCH2SPh)
		Company ( Di Cilla CDb)	-C(O)CH2CH2(p-
17.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	PhCH <sub>2</sub> SPh)
17.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
17.26		-C(O)Cl12(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
17.27	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
17.28	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
17.29	-C(O)pyridin-2-yl	-C(O)Cl12(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
17.30 17.31	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
17.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
17.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
17.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
17.35	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
17.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
17.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
17.38	-C(O)oxazo-2-yl	-C(O)Cl12(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
17.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
17.40	-C(O)benzofuran-2-yl	-C(O)CH2(bcnzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2-
17.40		·	yl)
17.41	-C(O)benzoturan-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
17.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-yl)
17.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
17.44	-C(O)benzimidazo-2-yl	-C(O)Cl12(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-vl)
17.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2- yl)
17.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-vl)
17.47	-C(O)o-Ph(P(O)Ph3)	-C(())m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
17.48	-C(O)Ph-2-(fluoren-9-yl)	-C(())Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-vl)
17.49	-C(O)N-indolin-2-one	-C(())indolin-2-yl	-C(O)indol-2-vl
17.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
17.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-(Ph)	-C(O)tetrahydrothiophen-3-yl-4-(Ph)
17.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
17.53	-C(O)tetrahydroisoquinolin-1-yl	-C.(O)tetrahydroisoquinolin-3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- yl)
17.54	-C(O)Cl12(N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
17.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydroxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)

17.56	CO.	C S S S S S S S S S S S S S S S S S S S	
17.57	OC CODE	O C C C C C C C C C C C C C C C C C C C	-oc N s
17.58		هُ حِيَّے ﴿ ﴾	o No
17.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
17.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
17.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
17.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
17.63	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
17.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
17.65	-C(O)CH2O(o-PhCH2COOH)	-C(O)CH2O(m-PhCH2COOH)	-C(O)CH2O(p-PhCH2COOH)
17.66			

Table 18

Formula I : A = -B(pinanediol) ; X = guanidinyl ;  $R^3$  = table below ;  $R^{11}$  = -CH<sub>2</sub>CH<sub>2</sub>Ph.

T	.1	.2	.3
18.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
18.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
18.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
18.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH <sub>2</sub> OH
18.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
18.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
18.7	-C(O)naphth-1-yl	-C(O)Cll2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
18.8	-C(O)naphth-2-yl	-C(O)Cll2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
18.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
18.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
18.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-hiphenyl)
18.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
18.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
18.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
18.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
18.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
18.18	-C(O)p-PhNHPh	-C(O)Cl12(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
18.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
18.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
18.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
18.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (o-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o- PhCH <sub>2</sub> SPh)
18.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (m-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m- PhCH <sub>2</sub> SPh)
18.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p- PhCH <sub>2</sub> SPh)
18.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
18.26	-C(O)cyclopentyl	-C(O)Cll2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
18.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
18.28	-C(O)CH2O(cyclopentyl)	-C(O)Cll2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
18.29	-C(O)CH2O(cyclohexyl)	-C(O)CII2NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
18.30	-C(O)pyridin-2-yl	-C(O)Cl12(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
18.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
18.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
18.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
18.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
18.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
18.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
18.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (imidazo-2-yl)
18.38	-C(O)oxazo-2-yl	-C(O)Cl12(oxazo-2-yl)	-C(O)Cl12Cl12(oxazo-2-yl)
18.39	-C(O)thionzo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
18.40	-C(O)benzoluran-2-yl	-C(O)CH2(benzoturan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2-yl)

18.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzoturan-3-yl)	-C(O)CH2CH2(benzoturan-3-yl)
18.42	-C(O)benzothiophen-2-yl	-C(O)Cl12(benzothiophen-2-yl)	C(O)CH2CH2(benzothiophen -2-yl)
18.43	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
18.44	-C(O)benzimidazo-2-yl	-C(O)CJ12(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
18.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
18.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-yl)
18.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
18.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
18.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-vl
18.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
18.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)teurahydrofuran-3-yl-4-(Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
18.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
18.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
18.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2- one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
18.55	-C(0)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
18.56		CN CO.	C COC
18.57	OC NHNH -	-00° N	-00 N S
18.58		oc, z, z, (=)	O Z O
18.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C3H7)CH2Ph
18.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH2Ph)
18.61	-C(O)C(CH3)2OPh	-C(U)CH(C2H5)OPh	-C(O)CH2OCH2Ph
18.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
18.63	-C(O)CH2O(o-PhCOOII)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
18.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH <sub>2</sub> O(m-PhCOOCH <sub>3</sub> )	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
18.65			
10.00	-C(O)CH2O(o-PhCH2COOH)	-C(O)CH2O(m-PhCH2COOH)	-C(O)CH2O(p-PhCH2COOH)

WO 95/09634

18.66	-oç <u>o</u>		
	, N <sub>M</sub> O		
	· ·		
		·	
	\ <b></b> /		

 $\frac{Table\ 19}{\text{Formula I: A = -B(pinanediol): X = guanidinyl: }} R^3 = table\ below\ ;\ R^{11} = -Ph.$ 

	.1	.2	.3
19.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
19.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
19.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
19.4	-C(O)o-PhCH2OII	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
19.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
19.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
19.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
19.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
19.9	-C(O)o-hiphenyl	-C(O)CH2(o-hiphenyl)	-C(O)CH2CH2(o-biphenyl)
19.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
19.12	-C(O)p-biphenyl	-C(O)CH2(p-hiphenyl)	-C(O)CH2CH2(p-biphenyl)
19.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
19.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
19.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
19.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
19.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
19.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
19.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
19.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
19.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
19.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
19.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
			PhCH <sub>2</sub> SPh)
19.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
19.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
19.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
19.27	-C(O)cyclohexyl	-C(O)CH <sub>2</sub> (cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
19.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
19.29	-C(0)CH <sub>2</sub> O(cyclohexyl)	-C(O)CII2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
19.30	-C(O)pyridin-2-yl	-C(O)CII2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
19.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
19.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
19.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
19.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
19.35	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-vl)
19.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
19.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
19.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
19.39	-C(O)thioazo-2-yl	-C(O)Cl12(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
19.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2-vl)

J 04

19.41	-C(O)benzofuran-3-yl	-C(O)Cl12(benzoturan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofwan-3-yl)
19.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	C(O)CH2CH2(benzothiophen -2-vl)
19.43	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-vl)
19.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
19.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2- vl)
19.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-yl)
19.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
19.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
19.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
19.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-vl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
19.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3- yl-4-(Ph)
19.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
19.53	-C(O)tetrahydroisoquinolin-1- yl	-C(O)tetrahydroisoquinolin- 3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
19.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
19.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
19.56	CO-	CO-	N4 ~
19.57	oc NH	-oc*N	-oc <sup>-N<sup>2</sup>s</sup>
19.57	OC NOT		-oc, N, 1, o, N, 1, o
19.58	-0C 0	-oc,	O
·	O	-OC,  (N) (N) (-C(O)N(C2H5)CH2Ph (-C(O)Ph-3-(CH2(thiophen-2-	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph -C(O)Ph-3-(CH <sub>2</sub> Ph)
19.58 19.59 19.60	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph -C(O)pyridin-3-yl-5-(Ph)	-OC,  (N) (N) (D) (-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-OC ON
19.58 19.59 19.60 19.61	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph -C(O)pyridin-3-yl-5-(Ph)	-OC, N N -C(O)N(C2H5)CH2Ph -C(O)Ph-3-(CH2(thiophen-2-yl)) -C(O)CH(C2H5)OPh	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph -C(O)Ph-3-(CH <sub>2</sub> Ph)
19.58 19.59 19.60	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph -C(O)pyridin-3-yl-5-(Ph)	-OC, N, N, -C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph -C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-OC O NHO -C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph -C(O)Ph-3-(CH <sub>2</sub> Ph)

205

19.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(p- PhCH <sub>2</sub> COOH)
19.66	OC NOTICE OF THE PROPERTY OF T		

5

Formula I: A = -B(pinanediol); X = guanidinyl;  $R^3$  = table below;  $R^{11}$  = -CH<sub>2</sub>(naphth-2-yl).

			.3
20.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
20.1	-C(O)CH <sub>2</sub> OPh	-C(O)CH2NHPh	-C(O)CH2SPh
20.2	-C(O)cH2OH	-C(O)m-PhOH	-C(O)p-PhOH
20.3	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
20.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
20.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
20.7	-C(O)naphth-1-vl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
20.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
20.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
20.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
20.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
20.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-PhOPh)
20.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhOPh)
20.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
20.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
20.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNIIPh)	-C(O)CH2CH2(m-PhNHPh)
20.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
20.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
20.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
20.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
20.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
20.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
			PhCH <sub>2</sub> SPh)
20.24	-C(O)p-PhCH2SPh	-C(O)Cll <sub>2</sub> (p-PhCll <sub>2</sub> SPh)	-C(O)CH2CH2(p-PhCH2SPh)
20.25	-C(O)adamantyl	-C(O)CH <sub>2</sub> (adamantyl)	-C(O)CH2CH2(adamantyl)
20.26	-C(O)cyclopentyl	-C(O)CH <sub>2</sub> (cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
20.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
20.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
20.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
20.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
20.31	-C(O)pyridin-3-yl	-C(O)Cll2(pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
20.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
20.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
20.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)

		•	
20.35	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
20.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
20.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
20.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
20.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
20.40	-C(O)benzofuran-2-yl	-C(O)CII <sub>2</sub> (benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2-yl)
20.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
20.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-yl)
20.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
20.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-vI)
20.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
20.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-yl)
20.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
20.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
20.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
20.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
20.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3- yl-4-(Ph)
20.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
20.53	-C(O)tetrahydroisoquinolin-1- yl	-C(O)tetrahydroisoquinolin- 3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
20.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
20.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol- 2-one)
20.56	CO-	CO.	COO
20.57	-oc NHNH	-0C N	-oc N s

20.58	·oc o	OC 2 2	oc s
20.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
20.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
20.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
20.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
20.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH <sub>2</sub> O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
20.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH <sub>2</sub> O(m-PhCOOCH <sub>3</sub> )	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
20.65	C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOII)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH2O(p- PhCH2COOH)
20.66			

Formula I: A = -B(pinanediol); X = -CH<sub>2</sub>NH<sub>2</sub>;  $R^3 = table below$ ;  $R^{11} = CH<sub>3</sub>$ 

	.1	.2	
21.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
21.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
21.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
21.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
21.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
21.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
21.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
21.8	-C(O)naphth-2-yl	-C(O)CH <sub>2</sub> (naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
21.9	-C(O)o-biphenyl	-C(O)CII2(o-hiphenyl)	-C(O)CH2CH2(o-bipbenyl)
21.10	-C(O)m-biphenyl	-C(O)Cl12(m-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
21.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
21.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
21.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
21.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
21.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
21.17	-C(O)m-PhNHPh	-C(O)Cl12(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
21.18	-C(O)p-PhNHPh	-C(O)Cl12(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
21.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
21.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
21.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)

21.22	-C(O)o-PhCH <sub>2</sub> SPh		-C(O)CH2CH2(o- PhCH2SPh)
			-C(O)CH2CH2(m-
21.23	-C(O)m-PhCH2SPh	-C(O)CH <sub>2</sub> (m-PhCH <sub>2</sub> SPh)	PhCH <sub>2</sub> SPh)
21.24	-C(O)p-PhCH <sub>2</sub> SPh		-C(O)CH2CH2(p- PhCH2SPh)
21.25	-C(O)adamantyl	-C(O)CH <sub>2</sub> (adamantyl)	-C(O)CH2CH2(adamantyl)
21.26		-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
21.27		-C(O)CII <sub>2</sub> (cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
21.27		-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
21.28		-C(O)CH2NH(cyclohexyl)	-C(O)CH2S(cyclohexyl)
		-C(O)CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
21.30		-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
21.31	-C(O)///11din 5 ):	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
21.32	O(O)/////	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
21.33	-C(O)furan-2-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
21.34	-C(O)furan-3-yl	-C(O)CI12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
21.35	-C(O)thiophen-2-yl	-C(O)C.112(Unophich-2-31)	yl)
22.24	2004	-C(O)CII2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
21.36	-C(O)thiophen-2-yl	-C.(O)C.112(UIK)/MC11-2-31/	vl)
21.37	-C(O)imidazo-2-yl	-C(O)Cll2(imidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (imidazo-2-yl)
21.38	-C(O)oxazo-2-yl	-C(U)Cl12(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
21.39	-C(O)thioazo-2-yl	-C(O)Cl12(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
21.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-
21.40	-C(O)()CH2O(u1an-2-)1		2-yl)
21.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 3-yl)
	0(0)	-C(O)CH2(benzothiophen-2-yl)	•
21.42	-C(O)benzothiophen-2-yl	-(.(O)(.112(tc:120tino)/men-2 )1)	C(O)CH2CH2(benzothiophen
ŀ			-2-vl)
21.43	-C(O)thiophen-2-yl	-C(O)CI12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
21.43	-C(O)dincijinen 2 yr		yl)
21.44	-C(O)henzimidazo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo-
			2-yl)
21.45	-C(O)benzoxazo-2-ÿT	-C(O)Cl12(benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-
			yl)
21.46	-C(O)benzothiazo-2-yl	-C(O)Cl12(benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo-
L			2-yl)
21.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
21.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
21.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
21.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-vl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
21.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-
12		(Ph)	vl-4-(Ph)
21.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
21.53	-C(O)tetrahydroisoquinolin-1-	-C(O)tetrahydroisoquinolin-3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-
	yl		yl)
21.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	
1 '	one)		one)

		00001-01-11	-C(O)CH <sub>2</sub> (N-dihydrothiazol-
21.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol-	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	2-one)
21.56	2-one)		CO-
21.30		O P	
21.57	-oc NH		-00 N S
21.58	-oc o	° 2 2 (2)	(2 co
21.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C3H7)CH2Ph
21.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2- vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
21.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
21.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
21.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
21.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
21.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m-PhCH2COOH)	-C(O)CH <sub>2</sub> O(p- PhCH <sub>2</sub> COOH)
21.66	-oc N		· ·

 $\frac{Table\ 22}{\text{Formula I: A = -B(pinanediol)}};\ X = -CH_2NH_2;\ R^3 = table\ below\ ;\ R^{11} = -CH_2(p-PhOH).$ 

	.1	.2	.3
22.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH <sub>2</sub> CH <sub>2</sub> Ph
22.2	-C(0)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
22.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
22.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
22.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
22.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH <sub>2</sub> COOH
22.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
22.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
22.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
22.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
22.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
22.12	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
		-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
22.14	-C(O)m-PhOPh	-C(U)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
22.15	-C(O)p-PhOPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
22.16	-C(O)o-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhNHPh)
22.17	-C(O)m-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
22.18	-C(O)p-PhNHPh	-C(O)CH2(o-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-PhSPh)
22.19	-C(O)x-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhSPh)
22.20	-C(O)m-PhSPh -C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
22.21		-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(0-
22.22	-C(O)o-PhCH2SPh	- (c)(c)(c)(2(0-1)(c)(2(0-1))	PhCH <sub>2</sub> SPh)
22.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
22.23			PhCH <sub>2</sub> SPh)
22.24	-C(O)p-PhCH2SPh	-C(O)Cl12(p-PhCl12SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-
	0(0),2		PhCH <sub>2</sub> SPh)
22.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
22.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
22.27	-C(O)cyclohexyl	-C(O)Cl12(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
22.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
22.29	-C(0)CH <sub>2</sub> O(cyclohexyl)	-C(O)Cll2NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
22.30	-C(O)pyridin-2-yl	-C(O)Cl12(pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
22.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
22.32	-C(O)pyridin-4-yl	-C(O)Cl12(pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
22.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
22.34	-C(O)furan-3-yl	-C(O)CH <sub>2</sub> (furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
22.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-
			yl)
22.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
			yl)
22.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
22.38	-C(O)oxazo-2-yl	-C(O)Cl12(oxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (oxazo-2-yl)
22.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)

22.40	-C(O)benzofuran-2-yl	-C(O)Cl12(benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoturan- 2-yl)
22.41	-C(O)benzofuran-3-yl	-C(O)Cl12(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoturan- 3-yl)
22.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-vl)
22.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-vl)
22.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
22.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-vl)
22.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-yl)
22.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
22.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
22.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
22.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
22.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydroturan-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3- yl-4-(Ph)
22.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
22.53	-C(O)tetrahydroisoquinolin-1- yl	-C(O)tetrahydroisoquinolin-3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
22.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
22.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
22.56	CO.	C N CO.	C CO.
22.57	-oc N NH	-00° N	-oc N s
22.58	-oc o	-oc, N, N, (1)	-0Ç N
22.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> II <sub>5</sub> )CII <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
22.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2- vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
22.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C <sub>2</sub> H <sub>5</sub> )OPh	-C(O)CH2OCH2Ph
22.62	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)

22.62	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
22.63	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
22.64	-C(O)CH2O(o-PhCH2COOH)	-C(O)CH2O(m-PhCH2COOH)	-C(O)CH2O(p-
22.03	-C(0)CH20(0-1 hCH20001)		PhCH2COOH)
22.66	-0C 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

Table 23 Formula I : A = -B(pinanediol) :  $X = CH_2NH_2$ ;  $R^3 = table below$ ;  $R^{11} = -CH_2CH_2Ph$ .

	.1	.2	.3
23.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
23.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
23.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
23.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH <sub>2</sub> OH
23.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
23.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
23.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
23.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
23.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
23.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
23.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
23.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
23.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhOPh)
23.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
23.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
23.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
23.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
23.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
23.20	-C(O)m-PhSPh	-C(O)CH <sub>2</sub> (m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
23.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
23.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
23.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m- PhCH <sub>2</sub> SPh)
23.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
23.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
23.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
23.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
23.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NII(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
23.29	-C(O)CH2O(cyclohexyl)	-C(O)CII2NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
23.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
23.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
23.32	-C(O)pyridin-4-yl	-C(O)Cll2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
23.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
23.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
23.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
23.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
23.37	-C(O)imidazo-2-vl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
23.38	-C(O)oxazo-2-vl	-C(O)CH2(0xazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
23.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
23.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2- vl)
23.41	-C(O)benzoturan-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)

23.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	C(O)CH2CH2(benzothiophen-
1		yl)	
1			2-yi)
23.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
23.44	-C(O)benzimidazo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-vl)
23.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2- vl)
23.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo-2-yl)
00.47	C(())= Dh(D(())Dha)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
23.47	-C(O)o-Ph(P(O)Ph3)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
23.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)indolin-2-yl	-C(O)indol-2-vl
23.49	-C(O)N-indolin-2-one	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
23.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)		
23.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
23.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )
23.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- yl)
	200000000000000000000000000000000000000	-C(O)CH <sub>2</sub> (N-benzoxazol-2-	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
23.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2- one)	one)	one)
23.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)
		QNS	
23.57	-oc NH	oc N	-oc N s
23.58	OC O	-oc \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	-0C 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
23.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
23.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2- yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
23.61	-C(O)C(CH3)2OPh	-C(O)CII(C <sub>2</sub> 115)OPh	-C(O)CH2OCH2Ph
23.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
23.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
23.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
23.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)

23.66	-oc o	
	ι <sub>ν</sub> μ <sub>Ο</sub>	
ł	7-1	
1		
	<_>	
		 · · · · · · · · · · · · · · · · · · ·

Table 24

Formula I: A = -B(pinanediol);  $X = CH_2NH_2$ ;  $R^3 = table below$ ;  $R^{11} = -Ph$ .

	1	.2	.3
24.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
24.1	-C(O)CH2OPh	-C(O)Cll2NHPh	-C(O)CH <sub>2</sub> SPh
	-C(O)o-PhOH	-C(())m-PhOH	-C(O)p-PhOH
24.3 24.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
24.4	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
24.5	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
24.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
24.7	-C(O)naphth-2-yl	-C(O)Cl12(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
24.9	-C(O)o-biphenyl	-C(O)CH <sub>2</sub> (o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
24.10	-C(O)m-hiphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
24.12	-C(O)p-hiphenyl	-C(O)Cl12(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
24.13	-C(O)o-PhOPh	-C(O)Cl12(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
24.14	-C(O)m-PhOPh	-C(O)Cl12(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
24.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
24.16	-C(O)o-PhNHPh	-C(O)CH <sub>2</sub> (o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
24.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
24.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
24.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
24.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
24.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
24.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
24.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
			PhCH <sub>2</sub> SPh)
24.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
24.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
24.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
24.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
24.28	-C(O)CH2O(cyclopentyl)	-C(O)Cll2Nll(cyclopentyl)	-C(O)CH2S(cyclopentyl)
24.29	-C(O)CH2O(cyclohexyl)		-C(O)CH <sub>2</sub> S(cyclohexyl)
24.30	-C(O)pyridin-2-yl	-C(O)Cl12(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
24.31	-C(O)pyridin-3-yl	-C(O)Cl·l <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
24.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
24.33	-C(O)furan-2-vl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
24.34	-C(O)furan-3-yl	-C(O)Cl12(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
24.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
24.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)

		-C(O)Cll2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
24.37	-C(O)imidazo-2-yl	-C(O)CH2(madaz(#2-yi)	-C(O)CH2CH2(oxazo-2-yl)
24.38	-C(O)oxazo-2-yl	-C(U)CH2(0xaz0-2-yi)	-C(O)CH2CH2(thioazo-2-yl)
24.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(benzofuran-2-
24.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	vi)
24.41	-C(O)benzofuran-3-yl	-C(O)Cl12(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)
24.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	C(O)CH2CH2(benzothiophen- 2-yl)
24.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
24.44	-C(O)benzimidazo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
24.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2- vl)
24.46	-C(O)benzothiazo-2-yl	-C(O)Cl12(benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo-2-yl)
24.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
24.48	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
24.49	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NIISO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
24.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
24.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
24.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
24.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
24.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2- one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)
24.56	CO.	CN CO.	CO.
24.57	-oc NHNH	-oc N	-oc N s
100	1	-oc,	-oc o
24.58	-0C 0		N <sub>H</sub> O
24.58		-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph

2/7

24.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
24.62	-C(O)CH2O(o-PhCH2OII)	-C(O)CH2O(m-PhCH2OH)	-C(U)CH2O(p-PhCH2OH)
24.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
24.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
24.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
24.66			

Table 25

Formula I : A = -B(pinanediol) ; X = -CH<sub>2</sub>NH<sub>2</sub>;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -CH<sub>2</sub>(naphth-2-yl).

	.1	.2	.3
25.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
25.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
25.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
25.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
25.5	-C(O)()-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
25.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
25.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
25.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
25.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
25.10	-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
25.12	-C(O)p-biphenyl	-C(O)CH2(p-hiphenyl)	-C(O)CH2CH2(p-biphenyl)
25.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
25.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
25.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
25.16	-C(O)o-PhNHPh	-C(U)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
25.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
25.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
25.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
25.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
25.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
25.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
25.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
	<u> </u>		PhCH <sub>2</sub> SPh)
25.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
25.25	-C(O)adamantyl	-C(O)CH <sub>2</sub> (adamantyl)	-C(O)CH2CH2(adamantyl)
25.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
25.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
25.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)Cl12S(cyclopentyl)
25.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(())CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
25.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-Ç(O)CH2CH2(pyridin-2-yl)
25.31	-C(O)pyridin-3-vl	-C(O)C112(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)

			200 201 201 4 - 11 - 4 - 11 T
25.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
25.33	-C(O)furan-2-yl	-C(O)CH <sub>2</sub> (furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
25.34	-C(O)furan-3-yl	-C(O)CH2(turan-3-yl)	-C(O)CH2CH2(furan-3-yl)
25.35	-C(O)thiophen-2-yl	-C(O)Cll2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
25.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
25.37	-C(O)imidazo-2-yl	-C(O)CH <sub>2</sub> (imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
25.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
25.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
25.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-2-yl)
25.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH2CH2(benzofuran-3- vl)
25.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen- 2-yl)
25.43	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
25.44	-C(O)benzimidazo-2-yl.	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
25.45	-C(O)benzoxazo-2-yl	-C(O)Cl12(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-yl)
25.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo-2-yl)
25.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
25.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
25.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
25.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
25.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
25.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
25.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- yl)
25.54	-C(O)Cl·l <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
25.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(C)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)
25.56	C CO-	CVN PO	COO
25.57	-oc NHNH	-oc*N	-oc N S

25.58	-0C 2	° 2 2 - (2)	-oc N O
25.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Pb
25.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
25.61	-C(O)C(CH3)2OPh	-C(U)CH(C2H5)OPh	-C(O)CH <sub>2</sub> OCH <sub>2</sub> Ph
25.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
25.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
25.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
25.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
25.66	-oc 0		

Table 26 Formula I : A = -B(pinanediol) ; X = -SC(=NII)NI12 ;  $R^3$  = table below ;  $R^{11}$  =  $CH_3$ 

		.2	.3
26.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
26.2	-C(O)CH <sub>2</sub> OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
26.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
26.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
26.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
26.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
26.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
26.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
26.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
26.10	-C(O)m-biphenyl	-C(O)CH <sub>2</sub> (m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
26.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
26.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
26.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
26.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
26.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
26.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
26.18	-C(O)p-PhNHPh	-C(O)CH <sub>2</sub> (p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
26.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
26.20	-C(O)m-PhSPh	-C(U)CH2(m-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhSPh)
26.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
26.22	-C(O)o-PhCH2SPh	-C(U)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
26.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m- PhCH2SPh)
26.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
26.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
26.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
26.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
26.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
26.29	-C(O)CH2()(cyclohexyl)	-C(O)Cl12NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
26.30	-C(O)pyridin-2-yl	-C(())CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
26.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
26.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
26.33	-C(O)furan-2-yl	-C(O)Cl12(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
26.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
26.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
26.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
26.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
26.38	-C(O)oxazo-2-yl	-C(O)CH <sub>2</sub> (oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
26.39	-C(O)thioazo-2-yl	-C(O)CH <sub>2</sub> (thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
26.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2- vl)
26.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH2CH2(benzofuran-3-yl)

26.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-yl)	C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen- 2-vl)
06.40	O(O)vhioshos 2 vi	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
26.43	-C(O)thiophen-2-vl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo-
26.44	-C(O)benzimidazo-2-yl		2-vI)
26.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
26.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo-2-yl)
26.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
26.48	-C(O)Ph-2-(fluoren-9-vl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-vl)
26.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-vl
26.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-vl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
26.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
26.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )
26.53	-C(O)tetrahydroisoquinolin-1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
26.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2- one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
26.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one)
	CO.	Chis	
26.57	-OC N NH	-0C N 0	-oc-N s
26.58	-0Ç 0	-oc z z	-oc'n
26.59	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C3H7)CH2Ph
26.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2- vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
26.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
26.62	-C(O)CH2O(o-PhCI12OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
26.63	-C(O)CH2O(o-PhCOOH)	-C(O)CII2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
26.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
	-C(O)CH2O(o-PhCH2COOH)	-C(O)CH2O(m-	-C(O)CH2O(p-PhCH2COOH)
26.65	-C(O)CH_O(GFTICH_COOH)	PhCH2COOH)	C(O)CitZO(pri neitZeoOtt)

26.66	-oc o	
	NO	

Table 27

Formula I : A = -B(pinanediol) ; X = -SC(=NH)NH<sub>2</sub> ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -CH<sub>2</sub>(p-PhOH).

	1	.2	.3
27.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
27.2	-C(O)Cli2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
27.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
27.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
27.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
27.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
27.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
27.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
27.9	-C(O)o-biphenyl	-C(())CH <sub>2</sub> (o-hiphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
27.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
27.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
27.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
27.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
27.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhOPh)
27.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
27.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
27.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
27.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
27.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
27.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
27.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
27.23	-C(O)m-PhCl1 <sub>2</sub> SPh-	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m- PhCH2SPh)
27.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
27.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
27.26	-C(O)cyclohexyl	-C(O)Cl12(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
27.27	-C(O)CH2O(cyclopentyl)	-C(O)CI12NII(cyclopentyl)	-C(O)CH2S(cyclopentyl)
27.28		-C(O)Cl12NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
27.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)Cl12(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
27.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
27.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
27.32	-C(O)pyridin-4-yl	-C(O)CH2(furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
27.33	-C(O)furan-2-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
27.34	-C(O)furan-3-yl	-C(O)CH2(thiophen-2-yl)	-C(())CH2CH2(thiophen-2-yl
27.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl
27.36 27.37	-C(O)thiophen-2-yl -C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)

27.38         -C(O)(xxxxx-2-y)         -C(O)(C112(thioazo-2-yl)         -C(O)CH2CH2(thioazo-2-yl)           27.39         -C(O)thioazo-2-yl         -C(O)C12(thioazo-2-yl)         -C(O)CH2CH2(thioazo-2-yl)           27.40         -C(O)benzofuran-2-yl         -C(O)C12(benzofuran-2-yl)         -C(O)CH2CH2(benzofuran-2-yl)           27.41         -C(O)benzofuran-3-yl         -C(O)C12(benzofuran-3-yl)         -C(O)CH2CH2(benzofuran-3-yl)           27.43         -C(O)benzofuran-3-yl         -C(O)CH2(benzofuran-3-yl)         -C(O)CH2(benzofuran-3-yl)			0(())CH-(	-C(O)CH2CH2(oxazo-2-yl)
C(O)CH2CH2(benzofuran-2-yl)	27.38	-C(O)oxazo-2-yl	-C(O)CH <sub>2</sub> (oxazo-2-yl)	
C(O)CH2CH2(benzofuran-3-yl)	27.39			
C(O)CH2CH2(benzothiophen-2-yl   C(O)CH2(CH2(CH2(benzothiophen-2-yl)   C(O)CH2(CH2(CH2(benzothiophen-2-yl)   C(O)CH2(CH2(CH2(benzothiophen-2-yl)   C(O)CH2(CH2(CH2(benzothiophen-2-yl)   C(O)CH2(CH2(benzothiophen-2-yl)   C(O)CH2(CH2(benzothiozo-2-yl)   C(O)Ph-2-(fluoren-9-yl)   C(O)Ph-3-(fluoren-9-yl)   C(O)Ph-4-(fluoren-9-yl)   C(O)Ph-4-(fluoren-	27.40	-C(O)benzoluran-2-yl		yl)
VI	27.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzoturan-3-yl)	
C(O)CH2CH2(Denzimidazo-2-yl)	27.42	-C(O)henzothiophen-2-yl		
C(O)CH2CH2(benzimidazo-2-yl)   C(O)CH2CH2(benzimidazo-2-yl)   C(O)CH2CH2(benzimidazo-2-yl)   C(O)CH2CH2(benzimidazo-2-yl)   C(O)CH2CH2(benzoxazo-2-yl)   C(O)CH2(CH2(benzoxazo-2-yl)   C(O)CH2(CH2(benzoxazo-2-yl)   C(O)CH2(Denzoxazo-2-yl)   C(O)CH2(Denzoxazo-2-yl)   C(O)CH2(Denzoxazo-2-yl)   C(O)CH2(Denzoxazo-2-yl)   C(O)CH2(Denzoxazo-2-yl)   C(O)CH2(Denzoxazo-2-yl)   C(O)CH2(Denzoxazo-1-2-yl)   C(O)CH2(Denzo	27.43	-C(O)thiophen-2-vl	-C(O)CH2(thiophen-2-yl)	
27.46				2-yl)
27.47	27.45	-C(O)benzoxazo-2-yl	-C(O)CI12(benzoxazo-2-yl)	vl)
C(O)Ph-3-(fluoren-9-yl)   C(O)Ph-3-(fluoren-9-yl)   C(O)Ph-4-(fluoren-9-yl)	27.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	yl)
27.48	27.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	
27.49				
27.50		-C(O)N-indolin-2-one		
27.51         -C(O)pyrrolidin-3-yl-4-(Ph)         -C(O)tetrahydrofuran-3-yl-4-(Ph)         -C(O)tetrahydrofuran-3-yl-4-(Ph)         -C(O)tetrahydrofuran-3-yl-4-(Ph)         -C(O)tetrahydronaphth-2-yl-2-(C(O)cH2((2-oxo)indolin-3-yl-2-(D)cyclopropyl-2.2-(Ph2))         -C(O)tetrahydroisoquinolin-3-yl-2-(C(O)th2((2-oxo)indolin-3-yl-2-(D)cyclopropyl-2.2-(Ph2))         -C(O)th2((2-oxo)indolin-3-yl-2-(D)cH2((2-oxo)indolin-3-yl		-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth-		<u> </u>
27.53 -C(O)ctarahydroisoquinolin-1-yl	27.51			4-(Ph)
27.53	27.52	-C(O)tetrahydronaphth-1-vl	-C(O)tetrahydronaphth-2-yl	
27.54 -C(O)CH <sub>2</sub> ( <i>N</i> -benzimidazol-2-one) -C(O)CH <sub>2</sub> ( <i>N</i> -benzoxazol-2-one) -C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol-2-one) -C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol-2-o			I -	
27.55 -C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one) -C(O)CH <sub>2</sub> (N-dihydrothiazol-2-one) -C(O)CH <sub>2</sub>	27.54			one)
27.57	27.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol-	-C(O)CH2(N-dihydrooxazol-2-	
27.58 -OC O O O O O O O O O O O O O O O O O O	27.56		CVN CO-	COO
27.59 -C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph -C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-vl))	27.57	-oc N NH	-oc N	-oc N s
27.60 -C(O)pyridin-3-yl-5-(Ph) -C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl)) -C(O)Ph-3-(CH <sub>2</sub> Ph)	27.58	OC N	oc z z	-oc N
27.60 -C(O)pyridin-3-yl-5-(Ph) -C(O)Ph-3-(CH2(thiophen-2-vl)) -C(O)Ph-3-(CH2Ph)	27 50	-C(O)N(CH2)CH2Ph	-C(O)N(C2115)CH2Ph	-C(O)N(C3H7)CH2Ph
0.00.00.00.00			-C(O)Ph-3-(CH2(thiophen-2-	
	27.61	I-COCCH2)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph

224

27.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
27.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
27.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(U)CH2O(p-PhCOOCH3)
27.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(U)CH <sub>2</sub> U(p-PhCH <sub>2</sub> COOH)
27.66	-oc of octanged and octanged an		

 $\label{eq:Table 28} \mbox{Formula I: A = -B(pinanediol); X = -SC(=NH)NH_2; R^3 = table below; R^{11} = -CH_2CH_2Ph. }$ 

		.2	.3
28.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
28.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH2SPh
28.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
28.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
28.5	-C(O)n-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
28.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PbCH2COOH
28.7	-C(O)naphth-1-yl	-C(O)CH <sub>2</sub> (naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
28.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
28.9	-C(O)o-hiphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
28.10	-C(O)m-biphenyl	-C(O)CII <sub>2</sub> (m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
28.12	-C(O)p-biphenyl	-C(O)CII2(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
28.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
28.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
28.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
28.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNIIPh)	-C(O)CH2CH2(o-PhNHPh)
28.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
28.18	-C(O)p-PhNHPh	-C(O)CH <sub>2</sub> (p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
28.19	-C(O)o-PhSPh	-C(O)Cl12(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
28.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
28.21	-C(O)p-PhSPh	-C(O)Cl12(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
28.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
		_	PhCH <sub>2</sub> SPh)
28.23	-C(O):n-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
			PhCH <sub>2</sub> SPh)
28.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-
			PhCH <sub>2</sub> SPh)
28.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
28.26	-C(O)cyclopentyi	-C(O)Cll2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
28.27	-C(O)cyclohexyl —	-C(O)Cl12(cyclohexyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (cyclohexyl)
28.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
28.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
28.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
28.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
28.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
28.33	-C(O)furan-2-yl	-C(O)CH <sub>2</sub> (furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
28.34	-C(O)furan-3-yl	-C(O)CH <sub>2</sub> (furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
28.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
			yl)
28.36	-C(O)thiophen-2-yl	-C(O)Cll2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
-	<u> </u>	0(0)011-6-11-2-2	yl)
28.37	-C(O)imidazo-2-yl	-C(O)Cl12(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
28.38	-C(O)oxazo-2-yl	-C(O)Cll2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
28.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)

			C(O)CH-CH-(hfumm
28.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 2-yl)
28.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH2CH2(benzofuran- 3-yl)
28.42	-C(O)benzothiophen-2-yl	-C(O)C11 <sub>2</sub> (benzothiophen-2-yl)	C(O)CH2CH2(benzothiophen -2-yl)
28.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
28.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
28.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
28.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-vl)
28.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
28.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-vl)
28.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-vl
28.50	- C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-vl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
28.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3- yl-4-(Ph)
28.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
28.53	-C(O)tetrahydroisoquinolin- 1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
28.54	-C(O)CH <sub>2</sub> ( <i>N</i> -benzimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
28.55	-C(O)CH <sub>2</sub> (N- dihydroimidazol-2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
28.56	CCO.	Chico.	CO.
28.57	-oc~NHNH	-0c-N <sup>0</sup> 0-0c-	-oc N S
28.58	-0¢ 0	OC Z Z	-0¢ N 0
28.59	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
28.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
28.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph

227

28.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
28.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
28.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
28.65	-C(O)CH <sub>2</sub> O(o- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOH)	-C(O)CH2O(p- PhCH2COOH)
28.66	-oc 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

<u>Table 29</u>

 $\label{eq:Formula I: A = -B(pinancdiol): X = -SC(=NII)NII_2; R^3 = table \ below \ ; R^{11} = -Ph.$ 

	.1	.2	.3
29.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(U)CH2CH2Ph
29.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
29.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
29.4	-C(O)o-PhCII2OH	-C(O)m-PhCl12OII	-C(O)p-PhCH2OH
29.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
29.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
29.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
29.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
29.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
29.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CII <sub>2</sub> CH <sub>2</sub> (m-hiphenyl)
29.12	-C(O)p-hiphenyl	-C(())CH <sub>2</sub> (p-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
29.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
29.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)Cl12CH2(m-PhOPh)
29.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhOPh)
29.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
29.17	-C(O)m-PhNHPh	-C(O)CII2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
29.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
29.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
29.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
29.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
29.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
			PhCH <sub>2</sub> SPh)
29.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
	•		PhCH <sub>2</sub> SPh)
29.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-
	_		PhCH <sub>2</sub> SPh)
29.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
29.26	-C(O)cyclopentyl	-C(O)Cl12(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
29.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
29.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
29.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
29.30	-C(O)pyridin-2-vl	-C(O)Cll2(pvridin-2-yl)	-C(O)Cl12CH2(pyridin-2-yl)

		Constitution 2 ml	-C(O)CH2CH2(pyridin-3-yl)
29.31	O(O)///	-C(U)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-4-yl)
29.32	C(O)/1/110	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(furan-2-yl)
29.33	C(O), as (ii)	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-3-yl)
29.34		-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(thiophen-2-
29.35	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	yl)
29.36	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2- yl)
29.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (imidazo-2-yl)
	-C(O)midaz(+2-yi	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
29.38		-C(O)Cl12(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
29.39	-C(O)thioazo-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-
29.40	-C(O)benzofuran-2-yl	- :	2-yl)
29.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 3-vl)
29.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	•
29.42	-C(O)Ochzolim/mch-2-31	yl)	C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen
ļ	·		-2,vl)
29,43	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
29.43	-C(O)micipalen-2-31		yl)
29.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
		-C(O)CH2(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-
29.45	-C(O)benzoxazo-2-yl	-(.(U)(.112(Deli20xa2(+2-yi)	vI)
20.45	C(O)	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo-
29.46	-C(O)benzothiazo-2-yl		2-vI)
29.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
29.47	-C(O)Ph-2-(fluoren-9-yl)	-C(())Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
29.49	-C(O)N-indolin-2-one	-C(())indolin-2-yl	-C(O)indol-2-yl
29.49	-C.(())/V-Ind()III-2-()iic	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
29.30	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-yl)	Ci(O)eyenspeanyr a (a as)	
29.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3- vl-4-(Ph)
29.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(())cyclopropyl-2,2-(Ph <sub>2</sub> )
29.52	-C(O)tetrahydroisoquinolin-	-C(O)tetrahydroisoquinolin-3-	-C(U)Cll2((2-oxo)indolin-3-
29.55	1-yl	yl	v1)
29.54	-C(O)CH <sub>2</sub> (N-benzimidazol-	-C(O)C112(N-benzoxazol-2-	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
	2-one)	one)	one)
29.55	-C(O)CH2(N-	-C(O)CH <sub>2</sub> (N-dihydrooxazol-	-C(O)CH <sub>2</sub> (N-dihydrothiazol-
L	dihydroimidazol-2-one)	2-one)	2-one)
29.56	co-	co-	ွှင့် င
- [	N <sub>v</sub> O	N <sub>O</sub>	N-¥
1	1 ( Y Y )	1 427 4	
1			
1	📞		
20.53		1	0
29.57		l ŭ	, , Ŭe
1	-OC N NH	-0C N 0	-oc N's
	1.00 )~		1 ()
1			

229

29.58	-0C 0 0	0 2 2 2 3	0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
29.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> II <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
29.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2- vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
29.61	-C(O)C(CH3)2OPh	-C(O)CH(C <sub>2</sub> H <sub>5</sub> )OPh	-C(O)CH2OCH2Ph
29.62	-C(O)CH2O(n-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
29.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
29.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
29.65	-C(O)CH <sub>2</sub> U(o- PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH2O(p- PhCH2COOH)
29.66	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

5

Formula I : A = -B(pinancdiol) ; X = -SC(=NII)NII<sub>2</sub> ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -CH<sub>2</sub>(naphth-2-yl).

	.1	.2	.3
30.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
30.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
30.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
30.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
30.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
30.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
30.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
30.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
30.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
30.10	-C(O)m-biphenyl	-C(O)Cll2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
30.12	-C(O)p-biphenyl	-C(O)Cll2(p-hiphenyl)	-C(O)CH2CH2(p-biphenyl)
30.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
30.14	-C(O)m-PhOPh	-C(O)Cll2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
30.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
30.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(U)CH2CH2(o-PhNHPh)
30.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
30.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
30.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
30.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
30.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
30.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o- PhCH <sub>2</sub> SPh)

		The state of the s	C(O)CHaCHa(m
30.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m- PhCH2SPh)
		DI OLL-ODE	-C(O)CH2CH2(p-
30.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH <sub>2</sub> (p-PhCH <sub>2</sub> SPh)	PhCH <sub>2</sub> SPh)
		244 211 / L	-C(O)CH2CH2(adamantyl)
30.25	-C(O/dothitan)	-C(O)CH2(adamantyl)	-C(O)CH2CH2((cyclopentyl)
30.26	C(O)O) Citif Citif	-C(O)Cll2(cyclopentyl)	-C(O)CH2CH2(cyclohexyl)
30.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	
30.28	-C(0)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
30.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
30.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
30.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
30.32	-C(O)pyridin-4-yl	-C(O)CH <sub>2</sub> (pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
30.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
30.34	-C(O)furan-3-yl	-C(O)CH <sub>2</sub> (furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
30.35	-C(O)thiophen-2-yl	-C(U)Cll2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
			yl)
30.36	-C(O)thiophen-2-yl	-C(())CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
			yl)
30.37	-C(O)imidazo-2-yl	-C(())Cll2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
30.38	-C(O)oxazo-2-yl	-C(O)Cll2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
30.39	-C(O)thioazo-2-yl	-C(O)CI12(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
30.40	-C(O)benzofuran-2-yl	-C(O)Cll <sub>2</sub> (benzoluran-2-yl)	-C(O)CH2CH2(benzofuran-
			2-yl)
30.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH2CH2(benzofuran-
		C(C)CUs (because in the page 2	3-yl)
30.42	-C(O)benzothiophen-2-yl	-C(O)Cl12(benzothiophen-2-	C(O)CH2CH2(benzothiophen
	1	yl)	-2-yl)
30.43	-C(O)thiophen-2-yl	-C(U)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
30.43	C(O)anophen 2 y		yl)
30.44	-C(O)benzimidazo-2-yl	-C(())CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo-
			2-yi)
30.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-
			vl)
30.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo-
<u> </u>			2-v1)
30.47	-C(O)o-Ph(P(O)Ph3)	-C(())m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
30.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
30.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl -C(O)cyclohexyl-2-(Ph)
30.50		-C(O)cyclopentyl-2-(Ph)	-C.(O)Cyclonexyl-2-(Fil)
l	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth		
20.51	-2-vl) -C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-
30.51	-C(O)pyrronum-5-y1-4-(171)	(Ph)	vl-4-(Ph)
30.52	-C(O)tetrahydronaphth-1-yl	-C(())tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
30.53	-C(O)tetrahydroisoquinolin-	-C(())tetrahydroisoquinolin-3-	-C(O)CH2((2-oxo)indolin-3-
30.55	1-vl	vi	vI)
30.54	-C(O)CH <sub>2</sub> (N-benzimidazol-	-C(U)Cl12(N-benzoxazol-2-	-C(O)CH <sub>2</sub> (N-benzothiazol-2-
1 30.54	2-one)	one)	one)
30.55	-C(O)C112(N-	-C(O)Cl12(N-dihydrooxazol-	-C(O)CH <sub>2</sub> (N-dihydrothiazol-
	dihydroimidazol-2-one)	2-one)	2-one)

30.56	CC O	CO. CO. CO. CO. CO. CO. CO. CO. CO. CO.	
30.57	-00 C N H	OC NO	-oc N s
30.58	OÇ N	OC Z Z (	0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0
30.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(U)N(C3H7)CH2Ph
30.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
30.61	-C(O)C(CH3)2()Ph	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
30.62	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
30.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH <sub>2</sub> O(p-PhCOOH)
30.64	-C(O)CH <sub>2</sub> O(o-PhCOOCH <sub>3</sub> )	-C(O)CH <sub>2</sub> O(m-PhCOOCH <sub>3</sub> )	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
30.65	-C(O)CH <sub>2</sub> O(o- PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH2O(p- РhCH2COOH)
30.66	o Ko		

5

Formula II :  $A = -B(OH)_2$ ; X = guanidinyl; Y = table below.

	.1	.2	.3
31.1	-C(O)Cl·l <sub>2</sub> ( <i>N</i> -benzimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
31.2	-C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	-C(O)Cll <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol- 2-one)
31.3	co o	CV, CO.	C COO

31.4	-oc N N H	0000	-oc N S
31.5	-0C 0	o z z (	-0C N O
31.6	-oc o	ر گرگی گرگی	

Table 32

Formula II:  $A = -B(OH)_2$ ;  $X = -CH_2Nll_2$ : Y = table below.

	.1	.2	.3
32.1	-C(0)CH <sub>2</sub> (N-benzimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
32.2	-C(0)CH <sub>2</sub> (N- dihydroimidazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol- 2-one)
32.3	CO-	Chico.	COO
32.4	-oc~N <sup>u</sup> NH	-oc N	00° 00° 00° 00° 00° 00° 00° 00° 00° 00°
32.5	-0°( N	· OC Z Z - C	CZZ O
32.6	-oc "-oc o		

5

Formula II :  $A = -B(OH)_2$  :  $X = -SC(=NH)NH_2$  : Y = table below.

	.1	.2	.3
33.1	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)Cl·l <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
33.2	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)Cl1 <sub>2</sub> ( <i>N</i> -dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
33.3	C C C C C C C C C C C C C C C C C C C	Co.	CN CO

33.4	-oc NH	0 0 0 0 0 0 0 0	ous occ
33.5		OC 2 2 (-)	
33.6			

5

Formula II : A = -B(pinanediol) ; X = guanidinyl ; Y = table below.

Γ	1	.2	.3
34.1	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2- one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
34.2	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CI1 <sub>2</sub> ( <i>N</i> -dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
34.3	CO.	Chico.	CNO CNO CNO CNO CNO CNO CNO CNO CNO CNO
34.4	-oc N NH	-0C N O	-oc N s
34.5	-0C 0	00 Z Z Z	-oc o
34.6	-oc o o		

Table 35

Formula II : A = -B(pinanediol);  $X = -CII_2NII_2$ ; Y = table below.

	<del>,</del>	1 2	.3
35.1	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
35.2	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol- 2-one)
35.3	CO.	CN CO.	CO.
35.4	-oc NHNH	.oc N	.00 N S
35.5	OC N	ر عرب کی این استان ا مراجع می استان	-oc NHO
35.6	och		

## Table 36

10

Formula II : A = -B(pinancdiol) ; X = -SC(=NII)NI12 ; Y = table below.

.1	.2	.3
-C(O)CH2(N-benzimidazol-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-	-C(O)CH2(N-benzothiazol-2-
2-one)	one)	one)
	-C(O)CH2(N-dihydrooxazol-	-C(O)Cl12(N-dihydrothiazol-
dihydroimidazol-2-one)	2-one)	2-one)
.co-	.co-	CO-
N.O		\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \
1 FY"Y	1674	N-W
	\ <u>-</u> 2	
	2-one) -C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	2-one) one)  -C(O)CH <sub>2</sub> (N- dihydroimidazol-2-one) 2-one)  CO-  CO-

36.4	-oc NHNH	-00° N	-oc N s
36.5	-0C 0		-0C N O
36.6	-oc 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,		

Table 37 Formula III : A = -B(OH)2 : X = -CN ;  $\mathbb{R}^3$  = table below :  $\mathbb{R}^{11}$  = CH3

	.1	.2	_3
37.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH <sub>2</sub> CH <sub>2</sub> Ph
37.2	-C(O)CH <sub>2</sub> OPh	-C(O)CII2NIIPh	-C(O)CH <sub>2</sub> SPh
	-C(O)ch2Orli -C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
37.3 37.4	-C(O)o-PhCH2OH	-C(O)m-PhC112OH	-C(O)p-PhCH2OH
	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
37.5	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
37.6 37.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
37.8		-C(O)CH2(naphth=2-y)	-C(O)CH2CH2(o-biphenyl)
37.9	-C(O)o-biphenyl	-C(O)CII2(m-hiphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-biphenyl)
37.10	-C(O)m-biphenyl	-C(O)CH2(p-hiphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-biphenyl)
37.12	-C(O)p-hiphenyl		-C(O)CH2CH2(o-PhOPh)
37.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	
37.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
37.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
37.16	-C(O)o-PhNHPh	-C(O)CII2(o-PhNIIPh)	-C(O)CH2CH2(o-PhNHPh)
37.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhNHPh)
37.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNIIPh)	-C(O)CH2CH2(p-PhNHPh)
37.19	-C(O)o-PhSPh	-C(O)CH <sub>2</sub> (o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
37.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhSPh)
37.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
37.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o- PhCH <sub>2</sub> SPh)
37.23	-C(O)m-PhCH <sub>2</sub> SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m- PhCH2SPh)
37.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p- PhCH <sub>2</sub> SPh)
37.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
37.26	-C(O)cyclopentyl	-C(O)Cl12(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
37.27	-C(O)cyclohexyl	-C(O)Cl12(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
37.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
37.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)Cl12S(cyclohexyl)
37.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
37.31	-C(O)pyridin-3-yl	-C(O)Cl12(pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
37.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
37.33	-C(O)furan-2-yl	-C(O)Cl12(furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
37.34	-C(O)furan-3-yl	-C(O)Cl12(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
37.35	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
37.36	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
37.37	-C(O)imidazo-2-yl	-C(O)Cl12(imidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (imidazo-2-yl)
37.38	-C(O)nxazo-2-yl	-C(U)CH2(mada20-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
37.39	-C(O)thioazo-2-yl	-C(O)C112(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
37.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2-
	· · · · · · · · · · · · · · · · · · ·		yl)
37.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-yl)

		C(())CHathanathianhan 2	7
37.42	-C(O)benzothiophen-2-yl	-C(C))CH <sub>2</sub> (benzothiophen-2-	C(O)CH2CH2(benzothiophen
		γl)	-2-vl)
07.43	C(C)thin-han 2 ad	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
37.43	-C(O)thiophen-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo-
37.44	-Č(O)benzimidazo-2-yl		2-yl)
37.45	-C(O)benzoxazo-2-yl	-C(O)Cl12(benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2- vl)
37.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-yl)
37.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
37.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-vl)	-C(O)Ph-4-(fluoren-9-yl)
37.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indol-2-yl
37.50	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
37.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
37.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
37.53	-C(O)tetrahydroisoquinolin-1-	-C(O)tetrahydroisoquinolin-3- yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- vl)
27.54	yl -C(O)CH <sub>2</sub> (N-benzimidazol-2-	-C(O)Cl12(N-henzoxazol-2-	-C(O)Cl12(N-benzothiazol-2-
37.54	one)	one)	one)
37.55	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
		QNS	
37.57	OC NH	-0c N	-00° N S
37.58	°C N	-oc z z	-oç o
37.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3H7)CH2Ph
37.60	-C(O)pyridin-3-yI-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2- yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
37.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
37.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
	<u> </u>	-C(O)Cll2O(m-PhCOOll)	-C(O)CH2O(p-PhCOOH)
37.63	-C(O)CH2O(o-PhCOOH)	-C(())CI()((m-PhCOOCII3)	-C(U)CH2O(p-PhCOOCH3)
37.64	-C(O)CII2O(o-PhCOOCII3)		-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
37.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH2O(p-FHCH2COOH)

37.66	-oc o		
ł	, N, O		
		·	

Table 38

Formula III : A = -B(OH)2 ; X = -CN ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = -CH<sub>2</sub>CH<sub>2</sub>Ph.

	1 .1	2	.3
38.1	-C(O)Ph	-C(O)CII2Ph	-C(O)CH2CH2Ph
38.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
38.3	-C(O)o-PhOII	-C(O)m-PhOH	-C(O)p-PhOH
38.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
38.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
38.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
38.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
38.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
38.9	-C(O)o-biphenyl	-C(O)CH <sub>2</sub> (o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
38.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
38.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
38.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
38.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
38.15	-C(O)p-PhOPh	-C(O)CH <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
38.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
38.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2Cl12(m-PhNHPh)
38.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
38.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
38.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
38.21	-C(O)p-PhSPh	-C(O)CH <sub>2</sub> (p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
38.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-PhCH2SPh)
38.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(U)CH2CH2(m-
			PhCH2SPh)
38.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-PhCH2SPh)
38.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
38.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
38.27	-C(O)cyclohexyl	-C(O)CH <sub>2</sub> (cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
38.28	-C(O)CH <sub>2</sub> O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
38.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
38.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
38.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
38.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
38.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-2-yl)
38.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (furan-3-yl)
38.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
38.36	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
38.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (imidazo-2-yl)
38.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(U)CH2CH2(oxazo-2-yl)

		CVOVCH-(thisway 2 vl)	-C(O)CH2CH2(thioazo-2-yl)
38.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(benzofuran-2-
38.40	-C(O)henzoiuran-2-yl	-C(O)CH2(benzofuran-2-yl)	vi)
38.41	-C(O)benzofuran-3-yl	-C(O)CH2(benzoluran-3-yl)	-C(O)CH2CH2(benzoturan-3-yl)
38.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen- 2-yl)
38.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
38.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
38.45	-C(O)benzoxazo-2-yl	-C(O)Cl12(benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
38.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo-2-yl)
38.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
38.48	-C(O)Ph-2-(fluoren-9-vl)	-C(())Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
38.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indol-2-yl
38.50	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth- 2-yl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
38.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(O)tetrahydrothiophen-3-yl- 4-(Ph)
38.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
38.53	-C(O)tetrahydroisoquinolin-1-yl		-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- yl)
38.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
38.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol-2-one)
38.56	CO-	CN CO.	CO-CO-CO-CO-CO-CO-CO-CO-CO-CO-CO-CO-CO-C
38.57	-oc~NHNH	-oc N 0	-oc^N <sup>H</sup> s
38.58	-0Ç 0	00 Z Z Z - (1)	-oc o
38.59	-C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C3H7)CH2Ph
38.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
38.61	-C(O)C(Cl13)2()Ph	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
38.62	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
38.63	-C.(O)C.fl2O(0-131C.OO11)	L-CRACHISCAM-LINCOCALL	

241

38.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH <sub>2</sub> O(m-PhCOOCH <sub>3</sub> )	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
38.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOII)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
38.66	-0C O		

Table 39

Formula III : A = -B(OH)2 ; X = -CH2NH2 ;  $\mathbb{R}^3$  = table below ;  $\mathbb{R}^{11}$  = CH3

	.1	.2	.3
39.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
39.2	-C(O)CH2OPh	-C(O)CH2NIIPh	-C(O)CH2SPh
39.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
39.4	-C(O)o-PhCH2OH	-C(O)m-PhCH <sub>2</sub> OH	-C(O)p-PhCH2OH
39.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
39.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(U)p-PhCH2COOH
39.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
39.8	-C(O)naphth-2-yl	-C(O)CH <sub>2</sub> (naphth-2-yl	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-2-yl)
39.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
39.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
39.12	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
39.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
39.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
39.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
39.16	-C(O)o-PhNHPh	-C(O)C112(o-PhN11Ph)	-C(O)CH2CH2(o-PhNHPh)
39.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
39.18	-C(O)p-PhNIIPh	-C(U)CII2(p-PhNIIPh)	-C(O)CH2CH2(p-PhNHPh)
39.19	-C(O)o-PhSPh	-C(O)C112(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
39.20	-C(O)m-PhSPh	-C(O)C112(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
39.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
39.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
L			PhCH <sub>2</sub> SPh)
39.23	-C(O)m-PhCl12SPh	-C(O)Cli2(m-PhCli2SPh)	-C(O)CH2CH2(m-
			PhCH2SPh)
39.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-
			PhCH <sub>2</sub> SPh)
39.25	-C(O)adamantyl	C(O)CH2(adamantyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
39.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(0)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
39.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
39.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
39.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
39.30	-C(O)pyridin-2-yl	-C(O)Cll2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
39.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
39.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(U)CH2CH2(pyridin-4-yl)
39.33	-C(O)furan-2-yl	-C(O)C[[2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)

		Occupation 2 vil	-C(O)CH2CH2(furan-3-yl)
39.34	-C(O)furan-3-yl	-C(O)Cl12(furan-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
39.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
39.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
39.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	
39.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
39.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
39.40	-C(O)benzoturan-2-yl	-C(O)CH2(benzoturan-2-yl)	-C(O)CH2CH2(benzofuran-2-
37.40			yl)
39,41	-C(O)benzoluran-3-yl	-C(O)CI12(benzofuran-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran-3-
32,12			yl)
39.42	-C(O)benzothiophen-2-yl	-C(O)CH2(benzothiophen-2-	C(O)CH2CH2(benzothiophen
	, , ,	ył)	·
			-2-vl) -C(O)CH2CH2(thiophen-2-yl)
39.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	
39.44	-C(O)benzimidazo-2-yl	-C(O)Cl12(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo-
			2-yl)
39.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(U)CH2CH2(benzoxazo-2-
33.13	e(O)		yl)
39.46	-C(O)benzothiazo-2-yl	-C(O)Cl12(benzothiazo-2-yl)	-C(O)CII2CH2(benzothiazo-
39.40	-6(0)//2120111235 - 3		2-vI)
39.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(U)p-Ph(P(O)Ph3)
39.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(())Ph-4-(fluoren-9-yl)
39.49	-C(O)N-indolin-2-one	-C(O)indolin-2-vl	-C(O)indol-2-yl
39.50	-	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
1	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth		٠
	-2-yl)	2.14	-C(O)tetrahydrothiophen-3-yl-
39.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(C))tetrahydrofuran-3-yl-4-	4-(Ph)
	<u></u>	(Ph)	-C(())cyclopropyl-2,2-(Ph <sub>2</sub> )
39.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-
39.53	-C(O)tetrahydroisoquinolin-1-	-C(O)tetrahydroisoquinolin-3-	_
1	yl	yl	yl) -C(O)CH2(N-benzothiazol-2-
39.54	-C(O)Cl12(N-benzimidazol-2-	-C(O)CH <sub>2</sub> (N-benzoxazol-2-	<b>I</b>
<u> </u>	one)	one)	one) -C(O)CH <sub>2</sub> (N-dihydrothiazol-
39.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol-	-C(U)Cl12(N-dihydrooxazol-2-	2-one)
<u></u>	2-one)	one)	
39.56	<u> </u>	co-	, ç <sub>0</sub> ;
1	N.O	N <sub>2</sub> O	N-K
1		141 1	
39.57	<del>                                     </del>	0	0
ו כ.עכ	l S	00° N	μ <sub>s</sub>
	-oc~NHNH	1-00 N J	-oc^N^S
	1 300 PK		( )
1			
1			1-00 0
30.58	00.0	-0C.	1 -00
39.58	-oc 0	-OC	-oc o
39.58	-0C 0	-0C <sub>3</sub>	Nuc
39.58	-0¢ 0	(N)	N <sup>H</sup> O
39.58	-0¢ 0	-0C, N	O LN JO
39.58	-0¢ 0	-0C, N	O LN LO

243

39.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(U)N(C3H7)CH2Ph
39.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
39.61	-C(O)C(CH3)2()Ph	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
39.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
39.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
39.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
39.65	-C(O)CH2O(o-PhCH2COOII)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)
39.66	-oc o		

Table 40 Formula III : A = -B(OH)2 ; X = CH2NH2;  $R^3$  = table below ;  $R^{11}$  = -CH2CH2Ph.

		.2	.3
40.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
40.1	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
40.2	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
40.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
40.4	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
40.5	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
40.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (napth-1-yl)
40.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
40.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
40.10	-C(O)m-hiphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
	-C(O)p-biphenyl	-C(O)CH <sub>2</sub> (p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
40.12	-C(O)o-PhOPh	-C(O)Cll2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
40.13		-C(O)Cll2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
40.14	-C(O)m-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
40.15	-C(O)p-PhOPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
40.16	-C(O)o-PhNHPh	-C(C)/CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
40.17	-C(O)m-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
40.18	-C(O)p-PhNHPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
40.19	-C(O)o-PhSPh	-C(O)Cl12(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
40.20	-C(O)m-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
40.21	-C(O)p-PhSPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
40.22	-C(O)o-PhCH2SPh	20,000,112,001,110,112,01,117	PhCH <sub>2</sub> SPh)
40.23	-C(O)m-PhCH2SPh	-C(U)Cl12(m-PhCH2SPh)	-C(O)CH2CH2(m- PhCH2SPh)
40.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p- PhCH2SPh)
40.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
40.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
40.27	-C(O)cyclohexyl	-C(O)Cl12(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
40.27	-C(O)CH2O(cyclopentyl)	-C(O)Cl12NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
40.28	-C(O)CH2O(cyclohexyl)	-C(())CH2NH(cyclohexyl)	-C(O)CH2S(cyclohexyl)
40.29	-C(O)pyridin-2-yl	-C(())Cl12(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
40.30	-C(O)pyridin-3-yl	-C(())CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
40.32	-C(O)pyridin-4-yl	-C(O)Cl12(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
40.33	-C(O)furan-2-yl	-C(O)Cl1 <sub>2</sub> (fumn-2-yl)	-C(O)CH2CH2(furan-2-yl)
40.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
40.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
40.36	-C(O)thiophen-2-yl	-C(O)Cl12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
40.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
40.37	-C(O)oxazo-2-yl	-C(())Cl12(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
40.39	-C(O)thioazo-2-yl	-C(())C112(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
	-C(O)benzoluran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-2-
40.40	-CAOACHAMMan-2-yi	0,0,0,0	vl)
40.41	-C(O)benzofuran-3-yl	-C(O)CH <sub>2</sub> (benzofuran-3-yl)	-C(O)CH2Cl12(benzofuran-3-yl)

WO 95/09634 PCT/US94/11280

40.42	-C(O)benzothiophen-2-yl	-C(O)C112(benzothiophen-2-	f_
40.42	-C(O)oenzounopnen-2-yr	-	C(O)CH2CH2(benzothiophen
		yl)	-2-vI)
40.42	C(O)thingh on 2 of	C(C)CHa(thiaghan 2 nl)	-C(O)CH2CH2(thiophen-2-yl)
40.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	
40.44	-C(O)benzimid:zo-2-yl	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-yl)
40.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
40.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiazo- 2-vl)
40.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
40.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
40.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
40.50	- C(O)C(CII3)2NIISO2(naphth	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
40.51	-2-yl) -C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-yl-
		(Ph)	4-(Ph)
40.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2.2-(Ph <sub>2</sub> )
40.53	-C(O)tetrahydroisoquinolin-1- yl	-C(O)(etrahydroisoguinolin-3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3- vl)
40.54	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2- one)	-C(O)CI1 <sub>2</sub> (N-benzothiazol-2-one)
40.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CI1 <sub>2</sub> ( <i>N</i> -dihydrooxazol-2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
	ON PO	Qng	
40.57	OC NH NH	-oc N	-oc N S
40.58		O Z Z ( )	OC NO CONTRACTOR OF NO
40.59	-C(O)N(CH3)CH2Ph	-C(O)N(C <sub>2</sub> 11 <sub>5</sub> )CH <sub>2</sub> Ph	-C(O)N(C3117)CH2Ph
40.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2- vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
40.61	-C(O)C(CH3)2()Ph	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
40.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
40.63	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
40.64	-C(O)CH <sub>2</sub> O(o-PhCOOCH <sub>3</sub> )	-C(O)CH2O(m-PhCOOCH3)	
40.65	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> ) -C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> COOH)

40.66	-00 0	
	ί <sub>ν</sub> μ <sub>Ο</sub>	
	آبر:	
	. >==	
	(_)	
ı	_	

Table 41

Formula III : A = -B(pinancdiol) ; X = -CN ;  $R^3 = table below$  ;  $R^{11} = CH_3$ 

	.1	.2	.3
41.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
41.2	-C(O)CH <sub>2</sub> OPh	-C(O)CH2NIIPh	-C(O)CH <sub>2</sub> SPh
41.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(())p-PhOH
41.4	-C(O)o-PhCH2OH	-C(O)m-PhCH <sub>2</sub> OH	-C(U)p-PhCH2OH
41.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
41.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(U)p-PhCH2COOH
41.7	-C(O)naphth-1-yi	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
41.7	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-vl	-C(O)CH2CH2(napth-2-yl)
41.6	-C(O)0-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-hiphenyl)
41.10	-C(O)m-biphenyl	-C(O)Cl12(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
41.12	-C(O)p-biphenyl	-C(())Cll2(p-hiphenyl)	-C(O)CH2CH2(p-biphenyl)
41.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
41.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
41.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
41.16	-C(O)o-PhN1IPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
41.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
41.17	-C(O)p-PhNHPh	-C(U)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
41.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
41.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
41.21	-C(O)p-PhSPh	-C(O)CII2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
41.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(U)CH2CH2(o-
41.22			PhCH <sub>2</sub> SPh)
41.23	-C(O)m-PhC112SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH2CH2(m-
77.23	-C(0/m : mc//2		PhCH <sub>2</sub> SPh)
41.24	-C(O)p-PhCH2SPh	-C(O)Cl12(p-PhCH2SPh)	-C(O)CH2CH2(p-
1	(C), 111112		PhCH2SPh)
41.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(U)CH2CH2(adamantyl)
41.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
41.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
41.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(())CH <sub>2</sub> S(cyclopentyl)
41.29	-C(O)CH2O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
41.30	-C(O)pyridin-2-yl	-C(O)CH <sub>2</sub> (pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
41.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
41.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
41.33	-C(O)Juran-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
41.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
41.35	-C(O)thiophen-2-vl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl

41.37	41.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
41.38   -C(O)xxzzo-2-y    -C(O)CH2(CH2(x)zzo-2-y )   -C(O)CH2(CH2(x)zzo-2-y )   -C(O)CH2(CH2(x)zzo-2-y )   -C(O)CH2(CH2(x)zzo-2-y )   -C(O)CH2(CH2(x)zo-2-y )   -C(O)CH2(x)zo-2-y )   -C(O)CH2(x)zo-2-y    -C(O)CH2(x)zo-				
41.39				
41.40		<u> </u>		
41.41				
41.42			-	yl)
41.42	41.41	-C(O)benzofurun-3-yl	-C(O)Cl1 <sub>2</sub> (benzofuran-3-yl)	
41.44		-C(O)benzothiophen-2-yl		- C(O)CII <sub>2</sub> CH <sub>2</sub> (benzothiophen
41.45	41.43		-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
41.45	41.44	-C(O)benzimidazo-2-yl	-C(O)CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo-
41.47   -C(O)o-Ph(P(O)Ph3)   -C(O)m-Ph(P(O)Ph3)   -C(O)p-Ph(P(O)Ph3)   -C(O)Ph(P(O)Ph3)   -C(O)Ph3)				-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2-yl)
41.48				2-yl)
41.49				
41.50		-C(O)Ph-2-(fluoren-9-vl)		-C(O)Ph-4-(fluoren-9-vl)
C(O)C(CH <sub>3</sub> ) <sub>2</sub> NIISO <sub>2</sub> (naphth -2-yl)		-C(O)N-indolin-2-one		-C(O)indol-2-yl
A1.52	41.50		-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
41.52   -C(O)tetrahydronaphth-1-yl   -C(O)tetrahydronaphth-2-yl   -C(O)cyclopropyl-2.2-(Ph2)     41.53   -C(O)tetrahydroisoquinolin-1-yl   -C(O)tetrahydroisoquinolin-3-yl   -C(O)CH2((2-oxo)indolin-3-yl)     41.54   -C(O)CH2(N-benzimidazol-2-one)   -C(O)CH2(N-benzothiazol-2-one)     41.55   -C(O)CH2(N-dihydroimidazol-2-one)   -C(O)CH2(N-dihydrothiazol-2-one)     41.56   -OC N N N N N N N N N N N N N N N N N N N				
41.53   -C(O)tetrahydroisoquinolin-1-yl		-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	
41.54 -C(O)CH <sub>2</sub> (N-benzimidazol-2-one) -C(O)CH <sub>2</sub> (N-benzoxazol-2-one) -C(O)CH <sub>2</sub> (N-benzoxhiazol-2-one) -C(O)CH <sub>2</sub> (N-dihydroxazol-2-one) -C(O)CH <sub>2</sub> (N-dihydroxazol-2-o		yl		-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-
41.55	· .	one)		-C(O)CH2(N-benzothiazol-2-
41.56 $\begin{array}{c ccccccccccccccccccccccccccccccccccc$				-C(O)CH <sub>2</sub> (N-dihydrothiazol-
-oc N N -oc		COO	CN CO.	
	41.57	-0C N NH	-oc NHO	-oc^ N ** S
41.59 -C(O)N(CH <sub>3</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>2</sub> H <sub>5</sub> )CH <sub>2</sub> Ph -C(O)N(C <sub>3</sub> H <sub>7</sub> )CH <sub>2</sub> Ph		-0C 0	° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° °	-oc N O
	41.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(U)N(C3H7)CH2Ph

41.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH2(thiophen-2-yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
41.61	-C(O)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph -C(O)CH2O(p-PhCH2OH)
41.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH) -C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
41.63	-C(O)CH <sub>2</sub> O(o-PhCOOH) -C(O)CH <sub>2</sub> O(o-PhCOOCH <sub>3</sub> )	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
41.64	-C(O)CH <sub>2</sub> O(o-PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOH)	-C(O)CH2O(p-PhCH2COOH)
41.66	-oc 0		

Table 42

Formula III : A = -B(pinanediol) ; X = -CN ;  ${\bf R}^3$  = table below ;  ${\bf R}^{11}$  = -CH<sub>2</sub>CH<sub>2</sub>Ph.

	l i	.2	.3
42.1	-C(O)Ph	-C(O)CH <sub>2</sub> Ph	-C(O)CH2CH2Ph
42.2	-C(O)CH2OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
42.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
42.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
42.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
42.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
42.7	-C(O)naphth-1-yl	-C(O)CH2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
42.8	-C(O)naphth-2-yl	-C(O)CH2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
42.9	-C(O)o-hiphenyl	-C(O)CH2(o-hiphenyl)	-C(O)CH2CH2(o-biphenyl)
42.10	-C(O)m-biphenyl	-C(O)Cll2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
42.12	-C(O)p-biphenyl	-C(O)CH2(p-biphenyl)	-C(O)CH2CH2(p-hiphenyl)
42.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
42.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
42.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
42.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
42.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
42.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhNHPh)
42.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-PhSPh)
42.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-PhSPh)
42.21	-C(O)p-PhSPh	-C(O)Cl12(p-PhSPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-PhSPh)
42.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-
<u> </u>			PhCII <sub>2</sub> SPh)
42.23	-C(O)m-PhCH2SPh	-C(O)CH2(m-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (m-
			PhCH <sub>2</sub> SPh)
42.24	-C(O)p-PhCH2SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (p-
40.05			PhCH <sub>2</sub> SPh)
42.25	-C(O)adamantyl	-C(O)Cl12(adamantyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (adamantyl)
42.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
42.27	-C(O)cyclohexyl	-C(O)Cl12(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
42.28	-C(O)Cl12O(cyclopentyl)	-C(O)Cli2NII(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
42.29	-C(O)CH2O(cyclohexyl)	-C(O)CII2NII(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
42.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-2-yl)
42.31	-C(O)pyridin-3-yl	-C(O)CH <sub>2</sub> (pyridin-3-yl)	-C(O)CH2CH2(pyridin-3-yl)
42.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
42.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
42.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
42.35	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
42.36	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-vl)
42.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
42.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
42.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)

		·	
42.40	-C(O)benzoturan-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzofuran- 2-yl)
42.41	-C(O)benzofuran-3-yl	-C(O)Cl12(benzofuran-3-yl)	-C(O)CH2CH2(benzofuran- 3-yl)
42.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH <sub>2</sub> CH <sub>2</sub> (benzothiophen -2-yl)
42.43	-C(O)thiophen-2-yl	-C(O)CH <sub>2</sub> (thiophen-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
42.44	-C(O)henzimidazo-2-yl	-C(O)Cl12(benzimidazo-2-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (benzimidazo- 2-vl)
42.45	-C(O)benzoxazo-2-yl	-C(O)CH <sub>2</sub> (benzoxazo-2-yl)	-C(U)CH <sub>2</sub> CH <sub>2</sub> (benzoxazo-2- vl)
42.46	-C(O)benzothiazo-2-yl	-C(O)CH2(benzothiazo-2-yl)	-C(O)CH2CH2(benzothiazo- 2-yl)
42.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
42.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
42.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
42.50	- C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
42.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofurain-3-yl-4- (Ph)	-C(U)tetrahydrothiophen-3- yl-4-(Ph)
42.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
42.53	-C(O)tetrahydroisoquinolin- 1-yl	-C(O)tetrahydroisoquinolin-3-yl	-C(O)CH <sub>2</sub> ((2-oxo)indolin-3-yl)
42.54	-C(O)CH <sub>2</sub> ( <i>N</i> -benzimidazol- 2-one)	-C(())Cli <sub>2</sub> (N-benzoxazol-2- one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
42.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
42.56		Q N 20	
42.57	-oc N NH	-oc N	-oc N S
42.58	-0Ç 0	OC 2 2 0	-oç o
42.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	-C(O)N(C3H7)CH2Ph
42.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2- yl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
		-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph

257

42.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
42.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
42.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH <sub>2</sub> O(p-PhCOOCH <sub>3</sub> )
42.65	-C(O)CH <sub>2</sub> O(o- PhCH <sub>2</sub> COOH)	-C(O)CH <sub>2</sub> O(m- PhCH <sub>2</sub> COOII)	-C(O)CH <sub>2</sub> O(p- PhCH <sub>2</sub> COOH)
42.66	-0¢,0,0		

Formula III : A = -B(pinancdiol) : X = -Cl1<sub>2</sub>NH<sub>2</sub> :  ${\bf R}^3$  = table below :  ${\bf R}^{11}$  = CH<sub>3</sub>

	]	.2	.3
43.1	-C(O)Ph	-C(O)CH2Ph	-C(O)CH2CH2Ph
43.2	-C(O)CH <sub>2</sub> OPh	-C(O)CH2NHPh	-C(O)CH2SPh
43.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
43.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)p-PhCH2OH
43.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
43.6	-C(O)o-PhCH2COOH	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
43.7	-C(O)naphth-1-yl	-C(O)CII2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
43.8	-C(O)naphth-2-yl	-C(O)Cl12(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
43.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH2CH2(o-biphenyl)
43.10	-C(O)m-biphenyl	-C(O)CH2(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
43.12	-C(O)p-biphenyl	-C(O)Cl12(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
43.13	-C(O)o-PhOPh	-C(O)CII2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
43.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
43.15	-C(O)p-PhOPh	-C(O)CH2(p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
43.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CII2CH2(o-PhNHPh)
43.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
43.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
43.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
43.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(O)CH2CH2(m-PhSPh)
43.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
43.22	-C(O)o-PhCH <sub>2</sub> SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH2CH2(o-
	•		PhCH <sub>2</sub> SPh)
43.23	-C(O)m-PhCl12SPh	-C(O)Cl12(m-PhCl12SPh)	-C(U)CH2CH2(m-
	•		PhCII2SPh)
43.24	-C(O)p-PhCH <sub>2</sub> SPh	-C(O)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-
	·		PhCH <sub>2</sub> SPh)
43.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
43.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> ((cyclopentyl)
43.27	-C(O)cyclohexyl	-C(O)CH2(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
43.28	-C(O)CH2O(cyclopentyl)	-C(O)CII2NII(cyclopentyl)	-C(O)Cli2S(cyclopentyl)
43.29	-C(O)Cl12O(cyclohexyl)	-C(O)Cl12N11(cyclohexyl)	-C(O)CH <sub>2</sub> S(cyclohexyl)
43.30	-C(O)pyridin-2-yl	-C(O)CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)

			-C(O)CH2CH2(pyridin-3-yl)
43.31	-C(O)pyridin-3-yl	-C(O)CH2(pyridin-3-yl)	
43.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-4-yl)
43.33	-C(O)furan-2-yl	-C(O)CH2(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
43.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
43.35	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(0)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
43.36	-C(O)thiophen-2-yl	-C(O)CII <sub>2</sub> (thiophen-2-yl)	-C(0)CH <sub>2</sub> CH <sub>2</sub> (thiophen-2-yl)
43.37	-C(O)imidazo-2-yl	-C(O)CH2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
43.38	-C(O)oxazo-2-yl	-C(O)Cl12(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
43.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)Cl12CH2(thioazo-2-yl)
43.40	-C(O)benzofuran-2-yl	-C(O)CH <sub>2</sub> (benzofuran-2-yl)	-C(U)CH2CH2(benzofuran- 2-vl)
43.41	-C(O)benzoturan-3-yl	-C(O)C112(benzoturan-3-yl)	-C(U)CH2CH2(benzofuran- 3-yl)
43.42	-C(O)benzothiophen-2-yl	-C(O)CH <sub>2</sub> (benzothiophen-2-yl)	- C(O)CH2CH2(benzothiophen -2-v!)
43.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-yl)
43.44	-C(O)benzimidazo-2-yl	-C(())CH <sub>2</sub> (benzimidazo-2-yl)	-C(O)CH2CH2(benzimidazo- 2-yl)
43.45	-C(O)benzoxazo-2-yl	-C(O)Cl1 <sub>2</sub> (benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-yl)
43.46	-C(O)benzothiazo-2-yl	-C(O)CH <sub>2</sub> (benzothiazo-2-yl)	-C(U)CH2CH2(benzothiazo- 2-yl)
43.47	-C(O)o-Ph(P(O)Ph3)	-C(O)m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
43.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(())Ph-4-(fluoren-9-yl)
43.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-vl
43.50	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth -2-vl)	-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
43.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4- (Ph)	-C(())tetrahydrothiophen-3- vl-4-(Ph)
43.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(O)cyclopropyl-2,2-(Ph <sub>2</sub> )
43.53	-C(O)tetrahydroisoquinolin- 1-yl	-C(O)tetrahydroisoquinolin-3- yl	-C(U)CH <sub>2</sub> ((2-oxo)indolin-3- yl)
43.54	-C(O)CH <sub>2</sub> (N-benzimidazol- 2-one)	-C(())Cll2(N-benzoxazol-2-one)	-C(())CH <sub>2</sub> (N-benzothiazol-2-one)
43.55	-C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	-C(O)Cl12(N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol- 2-one)
43.56	Ginydroimidazoi-2-dile)	CO- No	
43.57	-oc N N H	-oc N	-oc N S

255

43.58	OC OC	°°° × × •	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
43.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2II5)Cll2Ph	-C(O)N(C3H7)CH2Ph
43.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2-yl))	-C(O)l <sup>1</sup> h-3-(CH <sub>2</sub> Ph)
43.61	-C(0)C(CH3)2OPh	-C(O)CH(C2H5)OPh	-C(O)CH2OCH2Ph
43.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH2O(p-PhCH2OH)
43.63	-C(O)CH2O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
43.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CII2O(m-PhCOOCII3)	-C(O)CH2O(p-PhCOOCH3)
43.65	-C(0)CH <sub>2</sub> O(0- PhCH <sub>2</sub> COOH)	-C(O)CH2O(m- PhCH2COOH)	-C(O)CH2O(p- PhCH2COOH)
43.66	-0¢ 0, 0		

Formula III : A = -B(pinanediol) : X = C112N112:  $\mathbb{R}^3$  = table below :  $\mathbb{R}^{11}$  = -CH2CH2Ph.

	.1.	.2	.3
44.1	-C(O)Ph	-C(O)CII <sub>2</sub> Ph	-C(O)CH2CH2Ph
44.2	-C(O)CH <sub>2</sub> OPh	-C(O)CH2NHPh	-C(O)CH <sub>2</sub> SPh
44.3	-C(O)o-PhOH	-C(O)m-PhOH	-C(O)p-PhOH
44.4	-C(O)o-PhCH2OH	-C(O)m-PhCH2OH	-C(O)ր-PhCH2OH
44.5	-C(O)o-PhCOOH	-C(O)m-PhCOOH	-C(O)p-PhCOOH
44.6	-C(O)o-PhCH2COOII	-C(O)m-PhCH2COOH	-C(O)p-PhCH2COOH
44.7	-C(O)naphth-1-yl	-C(O)Cll2(naphth-1-yl)	-C(O)CH2CH2(napth-1-yl)
44.8	-C(O)naphth-2-yl	-C(O)Cll2(naphth-2-yl	-C(O)CH2CH2(napth-2-yl)
44.9	-C(O)o-biphenyl	-C(O)CH2(o-biphenyl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o-biphenyl)
44.10	-C(O)m-biphenyl	-C(O)Cl12(m-biphenyl)	-C(O)CH2CH2(m-biphenyl)
44.12	-C(O)p-biphenyl	-C(O)Cll2(p-biphenyl)	-C(O)CH2CH2(p-biphenyl)
44.13	-C(O)o-PhOPh	-C(O)CH2(o-PhOPh)	-C(O)CH2CH2(o-PhOPh)
44.14	-C(O)m-PhOPh	-C(O)CH2(m-PhOPh)	-C(O)CH2CH2(m-PhOPh)
44.15	-C(O)p-PhOPh	-C(O)CII <sub>2</sub> (p-PhOPh)	-C(O)CH2CH2(p-PhOPh)
44.16	-C(O)o-PhNHPh	-C(O)CH2(o-PhNHPh)	-C(O)CH2CH2(o-PhNHPh)
44.17	-C(O)m-PhNHPh	-C(O)CH2(m-PhNHPh)	-C(O)CH2CH2(m-PhNHPh)
44.18	-C(O)p-PhNHPh	-C(O)CH2(p-PhNHPh)	-C(O)CH2CH2(p-PhNHPh)
44.19	-C(O)o-PhSPh	-C(O)CH2(o-PhSPh)	-C(O)CH2CH2(o-PhSPh)
44.20	-C(O)m-PhSPh	-C(O)CH2(m-PhSPh)	-C(U)CH2CH2(m-PhSPh)
44.21	-C(O)p-PhSPh	-C(O)CH2(p-PhSPh)	-C(O)CH2CH2(p-PhSPh)
44.22	-C(O)o-PhCH2SPh	-C(O)CH2(o-PhCH2SPh)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (o- PhCH <sub>2</sub> SPh)

		DI CHE CINE	-C(U)CH2CH2(m-
44.23	-C(O)m-PhCl12SPh	-C(O)CH2(m-PhCH2SPh)	
	· · · · · · · · · · · · · · · · · · ·		PhCH <sub>2</sub> SPh)
44.24	-C(O)p-PhCH2SPh	-C(U)CH2(p-PhCH2SPh)	-C(O)CH2CH2(p-
			PhCH <sub>2</sub> SPh)
44.25	-C(O)adamantyl	-C(O)CH2(adamantyl)	-C(O)CH2CH2(adamantyl)
44.26	-C(O)cyclopentyl	-C(O)CH2(cyclopentyl)	-C(O)CH2CH2((cyclopentyl)
44.27	-C(O)cyclohexyl	-C(O)Cl12(cyclohexyl)	-C(O)CH2CH2(cyclohexyl)
44.28	-C(O)CH2O(cyclopentyl)	-C(O)CH2NH(cyclopentyl)	-C(O)CH <sub>2</sub> S(cyclopentyl)
44.29	-C(O)CH <sub>2</sub> O(cyclohexyl)	-C(O)CH2NH(cyclohexyl)	-C(())CH <sub>2</sub> S(cyclohexyl)
44.30	-C(O)pyridin-2-yl	-C(())CH2(pyridin-2-yl)	-C(O)CH2CH2(pyridin-2-yl)
44.31	-C(O)pyridin-3-yl	-C(())CII <sub>2</sub> (pyridin-3-yl)	-C(O)CH <sub>2</sub> CH <sub>2</sub> (pyridin-3-yl)
44.32	-C(O)pyridin-4-yl	-C(O)CH2(pyridin-4-yl)	-C(O)CH2CH2(pyridin-4-yl)
44.33	-C(O)furan-2-yl	-C(O)Cl12(furan-2-yl)	-C(O)CH2CH2(furan-2-yl)
44.34	-C(O)furan-3-yl	-C(O)CH2(furan-3-yl)	-C(O)CH2CH2(furan-3-yl)
44.35	-C(O)thiophen-2-yl	-C(O)CI12(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
44.55	-C(O)(mojnicis-2-3)		vI)
44.36	-C(O)thiophen-2-yl	-C(O)C112(thiophen-2-yl)	-C(O)CH2CH2(thiophen-2-
177.50	-C(O)unt/men 2 3.		yl)
44.37	-C(O)imidazo-2-yl	-C(O)Cll2(imidazo-2-yl)	-C(O)CH2CH2(imidazo-2-yl)
44.38	-C(O)oxazo-2-yl	-C(O)CH2(oxazo-2-yl)	-C(O)CH2CH2(oxazo-2-yl)
44.39	-C(O)thioazo-2-yl	-C(O)CH2(thioazo-2-yl)	-C(O)CH2CH2(thioazo-2-yl)
44.40	-C(O)benzofuran-2-yl	-C(O)CH2(benzofuran-2-yl)	-C(O)CH2CH2(benzofuran-
177.10			2-yl)
44.41	-C(O)benzoiuran-3-yl	-C(O)CH2(benzoturan-3-yl)	-C(O)Cl12CH2(benzofuran-
			3-vi)
44.42	-C(O)benzothiophen-2-yl	-C(O)Cll2(benzothiophen-2-	arough out a manking
		yl)	C(O)CH2CH2(benzothiophen
			-2-yl) -C(O)CH2CH2(thiophen-2-
44.43	-C(O)thiophen-2-yl	-C(O)CH2(thiophen-2-yl)	vl)
	000	-C(O)CH2(benzimidazo-2-yl)	-C(O)CH2CH2(henzimidazo-
44.44	-C(O)benzimidazo-2-yl	-(.(O)(.112(t)enzimidaz(+2-yi)	2-vl)
14.45	-C(O)benzoxazo-2-yl	-C(O)CH2(benzoxazo-2-yl)	-C(O)CH2CH2(benzoxazo-2-
44.45	-C(O)nenzoxazo-z-yi	-C(C)(C)(C)(Z((C)(X)X(X,C)-2-y))	vl)
44.46	-C(O)benzothiazo-2-yl	-C(O)Cl12(benzothiazo-2-yl)	-C(U)CH2CH2(benzothiazo-
44.40	-C.(O)(telizoniliaz.t)-2-y1	-0.000.112(00.1120.1111120.1111120.1111120.111111120.11111111	2-v1)
44.47	-C(O)o-Ph(P(O)Ph3)	-C(())m-Ph(P(O)Ph3)	-C(O)p-Ph(P(O)Ph3)
44.48	-C(O)Ph-2-(fluoren-9-yl)	-C(O)Ph-3-(fluoren-9-yl)	-C(O)Ph-4-(fluoren-9-yl)
44.49	-C(O)N-indolin-2-one	-C(O)indolin-2-yl	-C(O)indol-2-yl
44.50		-C(O)cyclopentyl-2-(Ph)	-C(O)cyclohexyl-2-(Ph)
	C(O)C(CH <sub>3</sub> ) <sub>2</sub> NHSO <sub>2</sub> (naphth		1
	-2-vl)		
44.51	-C(O)pyrrolidin-3-yl-4-(Ph)	-C(O)tetrahydrofuran-3-yl-4-	-C(O)tetrahydrothiophen-3-
		(Ph)	yl-4-(Ph)
44.52	-C(O)tetrahydronaphth-1-yl	-C(O)tetrahydronaphth-2-yl	-C(())cyclopropyl-2,2-(Ph <sub>2</sub> )
44.53	-C(O)tetrahydroisoquinolin-	-C(O)tetrahydroisoquinolin-3-	
<u></u>	1-yl	yl	vI)
44.54	-C(O)C112(N-benzimidazol-	-C(O)CH2(N-benzoxazol-2-	-C(O)Cl12(N-benzothiazol-2-
	2-one)	one)	one) -C(O)CH <sub>2</sub> (N-dihydrothiazol-
44.55	-C(O)CH <sub>2</sub> (N-	-C(O)Cl12(N-dihydrooxazol-	-C.(O)C.H2(N-dinydrodhazoi- 2-one)
1	dihydroimidazol-2-one)	2-one)	Z-OHE)

44.56	CO.	CO.	
44.57	-oc N NH	-oc N 0	-oc N S
44.58		° ( 2 2 - ( )	-oç o
44.59	-C(O)N(CH3)CH2Ph	-C(O)N(C2H5)CH2Ph	<sup>2</sup> C(O)N(C3H7)CH2Ph
44.60	-C(O)pyridin-3-yl-5-(Ph)	-C(O)Ph-3-(CH <sub>2</sub> (thiophen-2- vl))	-C(O)Ph-3-(CH <sub>2</sub> Ph)
44.61	-C(O)C(CH <sub>3</sub> ) <sub>2</sub> OPh	-C(U)CII(C2II5)UPh	-C(O)CH2OCH2Ph
44.62	-C(O)CH2O(o-PhCH2OH)	-C(O)CH2O(m-PhCH2OH)	-C(O)CH <sub>2</sub> O(p-PhCH <sub>2</sub> OH)
44.63	-C(O)CH <sub>2</sub> O(o-PhCOOH)	-C(O)CH2O(m-PhCOOH)	-C(O)CH2O(p-PhCOOH)
44.64	-C(O)CH2O(o-PhCOOCH3)	-C(O)CH2O(m-PhCOOCH3)	-C(O)CH2O(p-PhCOOCH3)
44.65	-C(O)CH <sub>2</sub> O(o- PhCH <sub>2</sub> COOH)	-C(O)Cl12O(m- PhCl12COOH)	-C(O)CH2O(p- PhCH2COOH)
44.66	-0C, N, O,		

Formula IV:  $A = -B(OH)_2$ : X = -CN; Y = table below.

	.1	.2	.3
45.1	-C(O)CH <sub>2</sub> (N-benzimidazol- 2-one)	-C(())Cl1 <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
45.2	-C(O)CH <sub>2</sub> (N-dihydroimidazol-2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrooxazol- 2-one)	-C(U)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
45.3	CO-	CN <sub>4</sub> O	CO.

45.4	OC NH NH	-oc N 0	oc N s
45.5		OC N N N	OC ON OCCUPANT OCCUPA
45.6	-0¢, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0,		

Formula IV :  $A = -B(OH)_2$  :  $X = -CH_2NH_2$  : Y = table below.

		.2	.3
46.1	-C(O)CH <sub>2</sub> (N-benzimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)Cl <sub>12</sub> (N-benzothiazol-2-one)
46.2	-C(O)CH <sub>2</sub> (N- dihydroimidazol-2-one)	-C(O)Cl1 <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
46.3	CV, CO.	Chico.	C COC
46.4	-oc_NNH	-0C_N_0	-oc N s
46.5	-0C 0	oc, z, z, c	-0C 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
46.6	-oc 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0		

257

Table 47

Formula IV : A = -B(pinanediol) : X = -CN : Y = uible below.

	T	7	T 3
47.1	-C(O)CH <sub>2</sub> (N-benzimidazol- 2-one)	-C(O)CI12(N-benzoxazol-2-one)	-C(O)CI12(N-benzothiazol-2-one)
47.2	-C(O)CH <sub>2</sub> (N- dihydroimidazol-2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrothiazol- 2-one)
47.3	CCO	Chico.	C C C
47.4	-oc N NH	-oc N	-oc N
47.5	-oc o	-0c/z/z-(1)	OC NO
47.6	-oc 0, 0		

Table 48

Formula IV :  $A = -B(pinanediol) : X = -CH_2NH_2 : Y = table below.$ 

		.2	.3
48.1	-C(O)CH <sub>2</sub> (N-benzimidazol-2-one)	-C(O)CH <sub>2</sub> (N-benzoxazol-2-one)	-C(O)CH <sub>2</sub> (N-benzothiazol-2-one)
48.2	-C(O)CH <sub>2</sub> (N-dihydroimidazol- 2-one)	-C(O)CH <sub>2</sub> (N-dihydrooxazol- 2-one)	-C(O)CH <sub>2</sub> ( <i>N</i> -dihydrothiazol- 2-one)
48.3	CO.	CN CO.	CNO CNO CNO CNO CNO CNO CNO CNO CNO CNO
48.4	-oc N NH	-oc N 0	-oc N s
48.5	-oc o	-oc z z	-oc o
48.6	-oc 0		

Table 49

Formula I: A = -B(pinanediol); X = See Table below;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = -CH_2CH_2Ph$ 

Example No.	.1	.2	.3
49.1	-NH <sub>2</sub>	-NHCH (=NH)H	-CH2NHC (=NH) NH2
49.2	-NHC(=NH)NH <sub>2</sub>	-CH <sub>2</sub> NH <sub>2</sub>	

### Table 50

Formula I: A = -B(pinanediol); X = See Table below;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = -N(CH_3)_2$ 

Example	.1	.2	.3
No.	·		
50.1	-NH <sub>2</sub>	-NHCH(=NH)H	-CH <sub>2</sub> NH <sub>2</sub>
50.2	-CH <sub>2</sub> NHC (=NH) NH <sub>2</sub>	-NHC (=NH) NH <sub>2</sub>	

### Table 51

Formula I: A = -B(pinanediol);  $X = -CH_2NH_2$ ;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See$  Table below

Example	.1	. 2	.3
No.			<u></u>
51.1	-CH <sub>2</sub> (m-PhCF <sub>3</sub> )	-CH <sub>2</sub> (m-PhCH <sub>3</sub> )	

### Table 52

Formula I:  $A = -B(OH)_2$ ;  $X = -CH_2NH_2$ ;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See Table below$ 

Example	.1	.2	.3
No.			
52.1	-CH <sub>2</sub> (m-PhCF <sub>3</sub> )	-CH <sub>2</sub> (m-PhCH <sub>3</sub> )	

Formula I:  $A = -B(OH)_2$ ;  $X = -CH_2NH_2$ ;  $R^3 = hydrocinnamoy1$ ;  $R^{11} = See Table below$ 

Example No.	.1	.2	.3
53.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2- phenyl)ethyl	-(2,2-diethyl- 2-phenyl)ethyl
53.2	-N-phenyl-N- methyl	-N-benzyl-N- methvl	-C (=0) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
53.3	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>	-(2,2- butanediy1-2- phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenyl]ethyl
53.4	-2-(3,5- dimethyl)phenyl ethvl	-cyclopropyl	-cyclohexyl

# Table 54

Formula I: A = -B(pinanediol);  $X = -CH_2NH_2$ ;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See Table below$ 

Example No.	.1	. 2	.3
54.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2- phenyl)ethyl	-(2,2-diethyl- 2-phenyl)ethyl

54.2	-N-phenyl-N-	-N-benzyl-N- methyl	-C(=0)CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
54.3	-C (=O) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>	-(2,2- butanediy1-2- phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenyl]ethyl
54.4	-2-(3,5-dimethyl)phenylethyl	-cyclopropyl	-cyclohexyl

Formula I:  $A = -B(OH)_2$ ;  $X = -NHC(=NH)NH_2$ ;  $R^3 = hydrocinnamoy1$ ;  $R^{11} = See Table below$ 

Example No.	.1	.2	.3
55.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2- phenvl)ethvl	-(2,2-diethyl- 2-phenyl)ethyl
55.2	-N-phenyl-N- methyl	-N-benzyl-N- methyl	-C (=0) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
55.3	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>	-(2,2-butanediy1-2-phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenyl]ethyl
55.4	-2-(3,5- dimethyl)phenyl ethyl	-cyclopropyl	-cyclohexyl

### Table 56

Formula I: A = -B(pinanediol);  $X = -NHC(=NH)NH_2$ ;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See Table below$ 

242

Example	.1	.2	.3
56.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2- phenyl)ethyl	-(2,2-diethyl- 2-phenyl)ethyl
56.2	-N-phenyl-N- methvl	-N-benzyl-N- methyl	-C (=O) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
56.3	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>	-(2,2-butanediy1-2-phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenyl]ethyl
56.4	-2-(3,5-dimethyl)phenylethyl	-cyclopropyl	-cyclohexyl

Table 57

Formula I:  $A = -B(OH)_2$ ; X = -NHC(=NH)H;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See Table below$ 

Example	.1	.2	.3
57.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2- phenyl)ethyl	-(2,2-diethyl- 2-phenyl)ethyl
57.2	-N-phenyl-N- methyl	-N-benzyl-N-	-C (=O) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
57.3	-C (=O)CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>	-(2,2- butanediyl-2- phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenyl]ethyl

57.4	-2-(3,5-	-cyclopropyl	-cyclohexyl
	dimethyl)phenyl		
,	ethy1		

Formula I: A = -B(pinanediol); X = -NHC(=NH)H;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See Table below$ 

Example No.	.1	.2	.3
58.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2- phenyl)ethyl	-(2,2-diethyl- 2-phenyl)ethyl
58.2	-N-phenyl-N- methyl	-N-benzyl-N- methyl	-C (=O) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
58.3	-С (=0) СН <sub>2</sub> СН <sub>2</sub> - СО <sub>2</sub> СН <sub>3</sub>	-(2,2-butanediy1-2-phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenyl]ethyl
58.4	-2-(3,5- dimethyl)phenyl ethyl	-cyclopropyl	-cyclohexyl

# Table 59

Formula I:  $A = -B(OH)_2$ ;  $X = -NH_2$ ;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See Table below$ 

Example No.	.1	.2	.3
59.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2-	-(2,2-diethyl- 2-phenyl)ethyl
		phenvl)ethvl	

59.2	-N-phenyl-N-methyl	-N-benzyl-N- methyl	-C (=O) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
59.3	-C(=0)CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>	-(2,2-butanediy1-2-phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenvl]ethyl
59.4	-2-(3,5-dimethyl)phenylethyl	-cyclopropyl	-cyclohexyl

Formula I: A = -B(pinanediol);  $X = -NH_2$ ;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = See$  Table below

Example No.	.1	.2	.3
60.1	-(2,2-dimethyl- 2-phenyl)ethyl	-(2,2- ethanediyl-2- phenyl)ethyl	-(2,2-diethyl- 2-phenyl)ethyl
60.2	-N-phenyl-N- methyl	-N-benzyl-N- methyl	-C (=0) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H
60.3	-C (=O) CH <sub>2</sub> CH <sub>2</sub> -CO <sub>2</sub> CH <sub>3</sub>	-(2,2- butanediyl-2- phenyl)ethyl	-[2,2-dimethyl- 2-(3,5- dimethyl)- phenyl]ethyl
60.4	-2-(3,5-dimethyl)phenylethyl	-cyclopropyl	-cyclohexyl

## Table 61

Formula I: A = See Table below;  $X = -CH_2NH_2$ ;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = -CH_3$ 

265

Example	.1	.2	.3
No. 61.1	-C(=0)CO <sub>2</sub> H	-C(=O)OCH <sub>2</sub> CH <sub>2</sub> -NHCbz	-C (=0) OCH <sub>2</sub> CH <sub>2</sub> - NH <sub>2</sub>
61.2	-C(=0)OCH3 HCL	-C(=O)OCH <sub>3</sub> free	-C(=O)CH3
61.3	-Сн <sub>2</sub> Он	-CH <sub>2</sub> OH; X = -CH <sub>2</sub> NHCbz	-C(=O)OH; X = -CH2NHCbz
61.4	-C (OH) (OCH <sub>3</sub> ) - C (=O) OCH <sub>3</sub>	-C (=O) NHNH2	

Formula I: A = -B(pinanediol);  $X = See Table below; <math>R^3 = 2-(2-cyanothiophenyl) - benzoyl$ ;  $R^{11} = -CH_3$ 

Example	.1	.2	.3
No. 62.1	-Br	-CH <sub>2</sub> Br	-SC (=NH) NH2
62.2	-N <sub>3</sub>	-CH2SC(=NH)NH2	-CH <sub>2</sub> N <sub>3</sub>
62.3	-NH <sub>2</sub>	-CH2NHC(=NH)NH2	-NHC (=NH)NH <sub>2</sub>
62.4	-NHC(=NH)H	-CH2NHC (=NH)H	-CH <sub>2</sub> NH <sub>2</sub>
62.4	-SCN		

## Table 63

Formula I: A = -B(pinanediol);  $X = See Table below; <math>R^3 = 2-(thiophenyl)benzoyl$ ;  $R^{11} = -CH_3$ 

Example	.1	.2	.3
No. 63.1	-Br	-CH <sub>2</sub> Br	-SC(=NH)NH2
63.2	-N <sub>3</sub>	-CH <sub>2</sub> SC (=NH) NH <sub>2</sub>	-CH <sub>2</sub> N <sub>3</sub>

63.3	-NH <sub>2</sub>	$-CH_2NHC (=NH)NH_2$	-NHC (=NH) NH <sub>2</sub>
63.4	-NHC (=NH) H	-CH <sub>2</sub> NHC(=NH)H	-CH <sub>2</sub> NH <sub>2</sub>
63.5	-SCN		

Formula I: A = -B(pinanediol);  $X = -CH_2NH_2$ ;  $R^3 = See$  Table below;  $R^{11} = -CH_3$ 

Example No.	.1	. 2	3
64.1	2-(benzyl)- benzyl	2-(benzyl)- benzyl; R <sup>11</sup> = -C(=0)CH	2-(pyrrol-1- ylmethyl)benzyl
64.2	2-(2- methylbenzyl)- benzovl	3-(3-trifluoro- methylbenzyl)- benzoyl	3-(3- chlorobenzyl)- benzovl
64.3	2-(2-cyano- benzyl)benzoyl		:

### Table 65

Formula I:  $A = -B(OH)_2$ ;  $X = -CH_2NH_2$ ;  $R^3 = See$  Table below;  $R^{11} = -CH_3$ 

Example No.	.1	.2	.3
65.1	2-(benzyl)- benzyl	2-(benzyl)- benzyl; R <sup>11</sup> = -C(=0)CH	2-(pyrrol-1- ylmethyl)benzyl
65.2	2-(2- methylbenzyl)- benzovl	3-(3-trifluoro- methylbenzyl)- benzovl	3-(3- chlorobenzyl)- benzovl

65.3	2-(2-cyano-	
· ·	benzvl)benzovl	

Table 66

Formula I:  $A = -B(OH)_2$ ;  $X = -CH_2NH_2$ ;  $R^3 = See$  Table below;  $R^{11} = -N(CH_3)CH_2Ph$ 

Example No.	.1	.2	.3
66.1	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
66.2	-C (=0) CH- (NHBOC)- CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH- (NHBOC) CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH- (NHBOC) - CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
66.3	-C (=0) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH <sub>2</sub> NH- S(0) <sub>2</sub> CH <sub>3</sub>	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>
66.4	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	-C (=0) CH <sub>2</sub> N- (CH <sub>3</sub> ) S (0) <sub>2</sub> CH <sub>3</sub>	

# Table 67

Formula I: A = -B(pinanediol);  $X = -CH_2NH_2$ ;  $R^3 = See$  Table below;  $R^{11} = -N(CH_3)CH_2Ph$ 

Example No.	.1	.2	.3
67.1	-C (=O) CH <sub>2</sub> CH <sub>2</sub> - CH <sub>2</sub> CO <sub>2</sub> H	-C(=0)CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C (=O) CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
67.2	-C (=0) CH- (NHBOC) - CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH- (NHBOC) CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH- (NHBOC) - CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
67.3	-C (=O) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C(=0)CH <sub>2</sub> NH- S(0) <sub>2</sub> CH <sub>3</sub>	-C (=O) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>

67.4	-C (=O) CH <sub>2</sub> CH <sub>2</sub> -	-C (=O) CH <sub>2</sub> N-	
	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	(CH <sub>3</sub> )S(O) <sub>2</sub> CH <sub>3</sub>	

Formula I:  $A = -B(OH)_2$ ;  $X = -CH_2NH_2$ ;  $R^3 = See$  Table below;  $R^{11} = -CH_2CH_2Ph$ 

Example	.1	.2	.3
68.1	-С (=0) СH <sub>2</sub> СH <sub>2</sub> - СH <sub>2</sub> СО <sub>2</sub> Н	-C(=0)CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C (=O) CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
68.2	-C (=0) CH- (NHBOC) - CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C(=0)CH- (NHBOC)CH <sub>2</sub> CO <sub>2</sub> H	-C (=O) CH- (NHBOC) - CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
68.3	-C (=O) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C(=0)CH <sub>2</sub> NH- S(0) <sub>2</sub> CH <sub>3</sub>	-C (=O) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>
68.4	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	-C (=0) CH <sub>2</sub> N- (CH <sub>3</sub> ) S (0) <sub>2</sub> CH <sub>3</sub>	

Table 69

Formula I: A = -B(pinanediol);  $X = -CH_2NH_2$ ;  $R^3 = See Table$  below;  $R^{11} = -CH_2CH_2Ph$ 

Example No.	.1	.2	.3
69.1	-C (=0) CH <sub>2</sub> CH <sub>2</sub> - CH <sub>2</sub> CO <sub>2</sub> H	-C (=O) CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C (=O) CH <sub>2</sub> C- (CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
69.2	-C (=0) CH- (NHBOC) - CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C(=0)CH- (NHBOC)CH <sub>2</sub> CO <sub>2</sub> H	-C (=0) CH- (NHBOC) - CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>
69.3	-C (=O) CH <sub>2</sub> CH <sub>2</sub> CO <sub>2</sub> H	-C(=O)CH <sub>2</sub> NH- S(O) <sub>2</sub> CH <sub>2</sub>	-C (=O) CH <sub>2</sub> CH <sub>2</sub> - CO <sub>2</sub> CH <sub>3</sub>

69.4	-C (=O) CH <sub>2</sub> CH <sub>2</sub> -	-C (=0) CH <sub>2</sub> N-	,
	CH <sub>2</sub> CO <sub>2</sub> CH <sub>3</sub>	(CH <sub>3</sub> )S(O) <sub>2</sub> CH <sub>3</sub>	

Formula I: A = See Table below;  $X = -CH_2NH_2$ ;  $R^3 = 2-(benzyl)benzoyl$ ;  $R^{11} = -CH_3$ 

Example	.1	.2	.3
70.1	-C (=0) CO <sub>2</sub> H	-C(=0)OCH <sub>2</sub> CH <sub>2</sub> -NHCbz	-C (=0) OCH <sub>2</sub> CH <sub>2</sub> - NH <sub>2</sub>
70.2	-C(=O)OCH3 HCL	-C(=0)OCH <sub>3</sub> free	-C (=O) CH <sub>3</sub>
70.3	-СH <sub>2</sub> ОН	-CH2OH; X = -CH2NHCbz	-C(=0)OH; X = -CH <sub>2</sub> NHCbz
70.4	-C (OH) (OCH <sub>3</sub> ) - C(=O)OCH <sub>3</sub>	-C(=0)NHNH <sub>2</sub>	

### Table 71

Formula I: A = See Table below;  $X = -CH_2NH_2$ ;  $R^3 = 2-(2-trifluoromethylbenzyl)benzoyl; <math>R^{11} = -CH_3$ 

Example No.	.1	.2	.3
71.1.	-C (=0) CO <sub>2</sub> H	-C(=0)OCH <sub>2</sub> CH <sub>2</sub> - NHCbz	-C (=0) OCH <sub>2</sub> CH <sub>2</sub> -
71.2	-C (=0) OCH <sub>3</sub> HCL	-C(=0)OCH <sub>3</sub> free	-C (=0) CH <sub>3</sub>
71.3	-СН <sub>2</sub> ОН	-CH2OH; X =  -CH2NHCbz	-C(=0)OH; X = -CH <sub>2</sub> NHCbz
71.4	-C.(OH) (OCH <sub>3</sub> )-	-C (=0)NHNH <sub>2</sub>	-

Formula I:  $A = -B(OH)_2$ ; X = See Table below;  $R^3 = hydrocinnamoyl$ ;  $R^{11} = -N(CH_3)_2$ 

Example	.1	.2	.3
No.	-NH <sub>2</sub>	-NHCH (=NH) H	-CH <sub>2</sub> NH <sub>2</sub>
	-CH2NHC (=NH)NH2	-NHC (=NH) NH <sub>2</sub>	

### Table 73

Formula I: A = -B(pinanedio1);  $X = -CH_2NH_2$ ;  $R^3 = -C(=0)CH_2CH_2CO_2H$ ;  $R^{11} = See$  Table below

Example	.1	.2	.3
No.			ļ
73.1	-CH <sub>2</sub> (m-PhCF <sub>3</sub> )	-CH <sub>2</sub> (m-PhCH <sub>3</sub> )	<u> </u>

### Table 74

Formula I: A = -B(pinanediol);  $X = -NHC(=NH)NH_2$ ;  $R^3 = See$  Table below;  $R^{11} = -CH_3$ 

Example	.1	.2	.3
74.1	2-(benzyl)- benzyl	2-(benzyl)- benzyl; R <sup>11</sup> = -C(=0)CH	2-(pyrrol-1- ylmethyl)benzyl
74.2	2-(2- methylbenzyl)- benzovl	3-(3-trifluoro- methylbenzyl)- benzoyl	3-(3- chlorobenzyl)- benzovl

74.3	2-(2-cyano-	
	benzvl)benzovl	· l

Formula I:  $A = -B(OH)_2$ ;  $X = -NHC(=NH)NH_2$ ;  $R^3 = See Table$  below;  $R^{11} = -CH_3$ 

Example No.	.1	.2	.3
75.1	2-(benzyl)- benzyl	2-(benzyl)- benzyl; R <sup>11</sup> = -C(=0)CH	2-(pyrrol-1- ylmethyl)benzyl
75.2	2-(2- methylbenzyl)- benzoyl	3-(3-trifluoro-methylbenzyl)-benzovl	3-(3- chlorobenzyl)- benzoyl
75.3	2-(2-cyano- benzyl)benzoyl		

### Table 76

Formula I:  $A = -B(OH)_2$ ;  $X = -CH_2NHC(=NH)H$ ;  $R^3 = See$  Table below;  $R^{11} = -cyclopropyl$ 

Example	.1	.2	.3
No.			
76.1	hydrocinnamovl	-C(=0)CH <sub>2</sub> 0Ph	-C (=0) CH <sub>2</sub> SPh

### Table 77

Formula I: A = -B(pinanediol);  $X = -CH_2NHC(=NH)H$ ;  $R^3 = See$  Table below;  $R^{11} = -cyclopropyl$ 

Example	.1	.2	.3
No.			
77.1	hvdrocinnamovl	-C(=0)CH20Ph	-C(=0)CH <sub>2</sub> SPh

Formula I:  $A = -B(OH)_2$ ;  $X = -CH_2NH_2$ ;  $R^3 = See$  Table below;  $R^{11} = -cyclopropyl$ 

Example	.1	.2	.3
No.	hvdrocinnamovl	-C(=0)CH <sub>2</sub> 0Ph	-C(=0)CH <sub>2</sub> SPh

Table 79

Formula I: A = -B(pinanediol);  $X = -CH_2NH_2$ ;  $R^3 = See Table$  below;  $R^{11} = -cyclopropyl$ 

Example	.1	.2	.3
No.			
79.1	hvdrocinnamovl	-C(=0)CH20Ph	-C(=0)CH2SPh

#### SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- (i) APPLICANT: Galemmo, et al.
- (ii) TITLE OF INVENTION: ELECTROPHILIC PEPTIDE ANALOGS AS INHIBITORS OF TRYPSIN LIKE ENZYMES
- (iii) NUMBER OF SEQUENCES:
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: THE DUPONT MERCK PHARMACEUTICAL COMPANY
- (B) STREET: 1007 MARKET STREET LEGAL DEPARTMENT
- (C) CITY: WILMINGTON
- (D) STATE: DE
  - (E) COUNTRY: U.S.A.

273

Gly

PAIGEOCHE - MICH DECORDANT I -

19898 (F) ZIP: (v) COMPUTER READABLE FORM: 3.50 INCH DISK (A) MEDIUM TYPE: COMPUTER: APPLE MACINTOSH (B) OPERATING SYSTEM: APPLE MACINTOSH (C) (D) SOFTWARE: MICROSOFT WORD (vi) CURRENT APPLICATION DATA: (A) APPLICATION NUMBER: N/A (B) FILING DATE: HEREWITH (C) CLASSIFICATION: (viii) ATTORNEY/AGENT INFORMATION: (A) NAME: REINERT, NORBERT F. (B) REGISTRATION NUMBER: 18,926 (C) REFERENCE/DOCKET NUMBER: DM-6666 (ix) TELECOMMUNICATION INFORMATION: (A) TELEPHONE: 302-892-8867 TELEFAX: 302-892-8536 (B) INFORMATION FOR SEQ ID NO:1: (2) SEQUENCE CHARACTERISTICS: (A) LENGTH: 14 (B) TYPE: amino acid TOPOLOGY: linear (C) (ii) MOLECULAR TYPE: peptide (vi) ORIGINAL SOURCE: synthetic (ix) FEATURE: (D) OTHER INFORMATION: thrombin inhibitor (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1: Leu Ser Asn Leu Ser Asn Leu Ser Asn Leu Ser Asn

INFORMATION FOR SEQ ID NO:2: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 14 amino acid (B) TYPE: (C) TOPOLOGY: linear (ii) MOLECULAR TYPE: peptide (vi) ORIGINAL SOURCE: synthetic (ix) FEATURE: (D) OTHER INFORMATION: thrombin inhibitor (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2: Leu Ser Asn Leu Ser Asn Leu Ser Asn 5 10 Gly INFORMATION FOR SEQ ID NO:3: (2) (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 14 amino acid (B) TYPE: (C) TOPOLOGY: linear (ii) MOLECULAR TYPE: peptide
(vi) ORIĞINAL SOURCE: synthetic (ix) FEATURE: (D) OTHER INFORMATION: thrombin inhibitor (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3: Leu Ser Asn Leu Ser Asn Leu Ser Asn

5 .

Gly

10

#### Utility

The compounds which are described in the present invention represent a novel class of potent, reversible inhibitors of trypsin-like enzymes. Trypsin-like enzymes are a group of proteases which hydrolyzed peptide bonds at basic residues liberating either a Cterminal arginyl or lysyl residue. Among these are enzymes of the blood coagulation and fibrinolytic system required for hemostasis. They are Factors II, 10 X, VII, IX, XII, kallikrein, tissue plasminogen activators, urokinase-like plasminogen activator, and plasmin. Enzymes of the complement system, acrosin (required for fertilization), pancreatic trypsin are also in this group. Elevated levels of proteolysis by 15 these proteases can result in disease states. example, consumptive coagulopathy, a condition marked by a decrease in the blood levels of enzymes of both the coagulation system, the fibrinolytic system and accompanying protease inhibitors is often fatal. 20 Intervention by a synthetic inhibitor would clearly be valuable. More specifically, proteolysis by thrombin is required for blood clotting. Inhibition of thrombin results in an effective inhibitor of blood clotting. The importance of an effective inhibitor of thrombin is 25 underscored by the observation that conventional anticoagulants such as heparin (and its complex with the protein inhibitor, antithrombin III) are ineffective in blocking arterial thrombosis associated with myocardial infractions and other clotting disorders. However, a low molecular weight thrombin inhibitor, containing a different functionality, was effective in blocking arterial thrombosis (Hanson and Harker, Proc. Natl. Acad. Sci. U.S.A. 85, 3184 (1988). Therefore, we have chosen to demonstrate utility of 35 compounds in the inhibition of thrombin, both as in

WO 95/09634 PCT/US94/11280

buffered solutions and in plasma. Specifically, the compounds have utility as drugs for the treatment of diseases arising from elevated thrombin activity such as myocardial infarction, and as reagents used as anticoagulants in the processing of blood to plasma for diagnostic and other commercial purposes.

Compounds of the present invention are expected to be effective in the control of aberrant proteolysis and a number of accompanying disease states such as inflammation, pancretitis, and heritary angioedema.

The effectiveness of compounds of the present invention as inhibitors of blood coagulation proteases was determined using purified human proteases and synthetic substrates following procedures similar to those described in Kettner et al. (1990).

For these assays, the rate of enzymatic (thrombin, Factor Xa, and Factor VIIa) hydrolysis of chromogenic substrates (S2238 (H-D-Phe-Pip-Arg-pNA), S2222, and S2288, respectively; Kabi Pharmacia, Franklin, OH) was measured both in the absence and presence of compounds of the present invention. Hydrolysis of the substrate resulted in the release of pNA, which was monitored spectrophotometrically by measuring the increase in absorbance at 405 nM. A decrease in the rate of absorbance change at 405 nm in the presence of inhibitor is indicative of enzyme inhibition. The results of this assay are expressed as inhibitory constant, Ki.

Thrombin and Xa determinations were made in 0.10 M sodium phosphate buffer, pH 7.5, containing 0.20 M NaCl, and 0.5 % PEG 8000. VIIa determinations were made in 0.05 M tris buffer, pH 7.6, containing 0.10 M NaCl, 4 mM CaCl<sub>2</sub>, and 0.1% bovine serum albumin. The Michaelis constant, K<sub>m</sub>, for substrate hydrolysis was determined at 25 °C using the method of Lineweaver and Burk.

277

10

30

35

Values of Ki were determined by allowing 0.2 - 0.5 nM human thrombin or human factor Xa (Enzyme Research Laboratories, South Bend, IN), or 50 nM human factor VIIa (BiosPacific, Emeryville, CA) react with the substrate (0.20 mM - 1 mM) in the presence of inhibitor. Reactions were allowed to go for 30 minutes and the velocities (rate of absorbance change vs time) were measured in the time frame of 25-30 minutes. The following relationship was used to calculate Ki values.

$$\frac{v_0-v_s}{v_s} = \frac{I}{\kappa_i (1 + S/K_m)}$$

where:

20

vo is the velocity of the control in the absence of inhibitor;

 $v_{\rm S}$  is the velocity in the presence of inhibitor; I is the concentration of inhibitor;  $K_{\rm i}$  is the dissociation constant of the enzyme:

inhibitor complex;

S is the concentration of substrate;  $K_{\text{m}}$  is the Michaelis constant.

Using the methodology described above,

representative compounds of this invention were
evaluated and found to exhibit a Ki of less 500 µM
thereby confirming the utility of compounds of the
invention as effective inhibitors of human blood
coagulation proteases. The results of these assays are
summarized in Table 80, where +++ indicates a Ki < 500
nM; ++ indicates a Ki < 50,000 nM; and + indicates a
Ki 500,000 < nM; - indicates inactive.

Table 80. Ki values for inhibition of Serine 35 Proteases by compounds of the present invention.

WO 95/09634 PCT/US94/11280

Ex No.	Thrombin	Factor X	a Factor	IC50
			VIIa	Thrombin time
49.1.1	+++	++	inactive	NT
49.1.2	+++	NT	NT	+++
49.1.3	+++	+++	NT	+++
49.2.1	+++	+++	NT	++
50.1.1	NT	NT	NT	NT
50.1.2	+++	NT	NT	+++
50.1.3	+++	NT	NT	NT
51.1.1	+++	+++	++	NT
51.1.2	+++	NT	NT	NT
52.1.1	+++	+++	++	NT
52.1.2	+++	+++	++	+++
53.1.1	+++	+++	+++	+++
53.1.2	+++	+++	+++	+++
53.2.1	+++	NT	NT	++
53.2.2	+++	+++	+++	NT
53.4.3	+++	++	+++	NT
54.1.1	+++	NT	NT	NT
54.1.2	+++	++	+++	+++
54.1.3	+++	++	+++	NT
54.2.1	+++	NT	NT	++
54.2.2	+++	NT	NT	++
54.2.3	+++	++	++	NT
54.3.1	. +++	++	++	NT
54.3.2	+++	++	+++	NT
54.3.3	NT	++	+++	NT
54.4.1	. NT	++	++	NT
54.4.2	+++	++	+++	+++
54.4.3	+++	++	+++	++
55.1.1	+++	+++	+++	NT
56.1.1	+++	+++	+++	NT
56.1.2	+++	+++	+++	NT
56.3.3	3 +++	+++	+++	NT
56.4.3	L NT	NT	NT	NT

			•	
57.1.1	+++	+++	+++;	NT
57.1.2	+++	+++	+++	NT
57.4.2	+++	. ++	NT	+++
58.1.1	+++	+++	+++;	NT
58.3.3	NT	NT	NT !	NT
58.4.1	NT	NT	NT	NT
58.4.2	++	NT	NT	NT
59.1.1	+++	++	+++	NT
59.4.2	+++	++	NT	,+++
60.1.1	+++	++	NT :	NT
60.3.3	+++	++	++	NT
60.4.1	+++	++	++	NT
60.4.2	+++	NT	NT	NT
61.1.1	+++	++	++ {	ŊT
61.1.2	++	Inactive	Inactive	NT
61.1.3	++	++	Inactive	NT
61.2.1	++	++	++	NT
61.2.2	++	Inactive	Inactive	NT
61.2.3	NT	Inactive	Inactive	NT
61.3.1	++.,	Inactive	Inactive	NT
61.3.2	++	Inactive	Inactive	NT
61.3.3	++	++	inactive	NT
61.4.1	+++	++	++	<u>,</u> ++
62.1.1	++	++	++	ΝТ
62.1.2	++	++	++	NT
62.1.3	+++	+++	+++	NT
62.2.1	++	++	++	NT
62.2.2	+++	+++	++	NT
62.2.3	++	++	++	NT
62.3.1	++	++	++	NT
62.3.2	+++	+++	++	NT
62.3.3	+++	+++	+++	· NT
62.4.1	+++	+++	+++	NT
62.4.2	+++	+++	++	NT
62.4.3	+++	+++	++	++


WO 95/09634 PCT/US94/11280

			•	
63.1.3	+++	+++	++	NT
63.3.1	+++'	++	inactive	NT
63.4.1	+++	+++	++	+++
63.4.2	+++	+++	++	++
63.5.1	++	++	inactive	NT
64.1.1	+++	++	++	Inactive
64.1.2	+++	++	+++	NT
64.1.3	+++	NT	NT	++
64.2.2	+++	+++	++	NT
64.2.3	+++	+++	+++	NT
65.1.3	+++	NT	NT	++
66.1.1	+++	+++	++	++
66.1.2	+++	NT	++	NT
66.1.3	+++	NT	++	NT
66.3.2	+++	+++	+++	+++
67.1.2	+++	++	++	NT
67.1.3	+++	+++	++	NT
67.3.2	+++	NT	NT	NT
68.2.1	+++	++	++	NT
68.3.1	+++	++	++	NT
68.3.3	+++	++	++	++
68.4.1	+++	+++	++	++
68.4.2	+++	NT	NT	NT
69.1.1	+++	+++	++	++
69.2.1	+++	NT	+++	NT
69.2.2	+++	NT	+++	NT
69.2.3	+++	· NT	++	NT
69.3.2	+++	+++	+++	+++
69.4.1	+++,	++	++	NT
69.4.2	+++	NT	NT	NT
70.4.2	++	++	inactive	NT
71.4.2	++	NT	NT	NT
72.1.3	+++	NT	NT	+++
73.1.2	+++	+++	++	NT
75.3.1	NT	NT	+++	NT

76.1.1	+++	NT	N.I.	14.1
77.1.1	+++	NT	NT	NT
78.1.2	+++	NT	ΝΤ	+++
78.1.3	+++	NT	NT	+++
79.1.2	+++	NT	NТ	+++
70 1 3	444	יזיע	NT	+++

WHAT IS CLAIMED IS:

1. A compound of formula (I):



5

or a pharmaceutically acceptable salt, hydrate or prodrug thereof, wherein:

10

 $R^1$  is

- a)  $-(C_1-C_{12} \text{ alkyl})-X$ ,
- b)  $-(C_1-C_{12} \text{ alkenyl})-X$ , or

C) -(CH<sub>2</sub>)<sub>q</sub> (CH<sub>2</sub>)<sub>p</sub>X

15

X is

- a) halogen,
- b) -CN,
- 20 c)  $-NO_2$ ,
  - d) -CF3,
  - e) -NH<sub>2</sub>,
  - f) -NHOR<sup>2</sup>,
  - g)  $-NHC (=NH) R^2$ ,
- 25 h) -NHC (=NH) NHOH,
  - i) -NHC(=NH)NHNH2,
  - j) -NHC(=NH)NHCN,
  - )  $-NHC (=NH) NHR^2$ ,
  - j)  $-NHC(=NH)NHCOR^2$ ,

```
k) -C (=NH) NHR<sup>2</sup>
         1) -C (=NH) NHCOR^2
         m) -C(=0)NHR<sup>2</sup>,
         n) -CO_2R^2
         o) -OR^2,
5
         p) -OCF3,
          g) -S(0)_{r}R^{2},
          r) -SC(=NH)NHR^2, or
          s) -SC(=NH)NHC(=0)R^2;
10
     R^2 is
          a) hydrogen, or
          b) C_1-C_4 alkyl;
     R^3 is
15
           a) -C(=0)-aryl
           b) -C(=0)-(CH_2)_p-CR^6R^7-(CH_2)_q-aryl,
           c) -C(=0)-(C_2-C_5 \text{ alkenyl})-aryl,
           d) -C(=O)-W-CR8R9-aryl, with the proviso that W
                 cannot be a bivalent oxygen atom,
20
           e) -C(=0)-CR^8R^9-W-(CH_2)_r-aryl, with the proviso
                  that W cannot be -NR^4- or -NC(=0)R^4-,
            f) -C(=O)-heteroaryl,
           g) -C(=0)-(CH_2)_p-CR^6R^7-(CH_2)_q-heteroaryl,
           h) -C(=0)-(C_2-C_5 \text{ alkenyl})-heteroaryl,
25
            i) -C(=0) -W-CR<sup>8</sup>R<sup>9</sup>-heteroaryl,
            j) -C(=0)-CR^8R^9-W-(CH_2)_r-heteroaryl, with the
                 proviso that W cannot be -NR^4- or -NC(=0)R^4-.
            k) -C(=0)-heterocycle,
            1) -C(=0)-(CH_2)_p-CR^6R^7-(CH_2)_q-heterocycle,
30
            m) -C(=0)-(C_2-C_5 \text{ alkenyl})-heterocycle,
            n) -C(=0) -W-CR<sup>8</sup>R<sup>9</sup>-heterocycle,
            o) -C(=0)-CR^8R^9-W-(CH_2)_r-heterocycle, with the
                  proviso that W cannot be -NR4- or -NCOR4-,
            p) -C(=0)-(CH<sub>2</sub>)t-adamantyl,
35
```

q)  $-C(=0)-(CH_2)_{t}-(C_5-C_7 \text{ cycloalkyl})$ ,

r)  $-C(=O)-(CH_2)_t-W-(C_5-C_7 \text{ cycloalkyl})$ ,

s)

5 limited to phenyl,

H or 
$$R^{13}$$

$$(CH_2)_p\text{-W-}(CH_2)_q\text{-aryl}, \text{ wherein aryl}$$

is limited to phenyl,

10

H or R<sup>13</sup>

15

with the proviso that  $R^{13}$  cannot be  $-N(C_1-C_4$  alkyl)2 when A is  $-C(=0)R^{14}$ ,

10 y) 
$$(CH_2)_p$$
  $(CH_2)_r$   $R^{10}$   $(CH_2)_q$ 

15 aa)

WO 95/09634 PCT/US94/11280

H or R<sup>13</sup>

ee)

5

10

hh)

10

ii)

jj) 
$$-(C(=0)-(CR^8R^9)-NR^{11'})_{v}-R^{11};$$
  
kk)  $-(C(=0)-(CR^8R^9)-NR^{11'})_{v}-C(=0)R^{11};$ 

11)  $-(C(=0)-(CR^8R^9)-NR^{11})_v-C(=S)R^{11};$ 

mm)

nn)

10

00) 
$$-C(=0) - (CR^8R^9) - NHS(0)_{r}R^8$$
;

pp)

15 qq)

 ${\bf R^4}$  and  ${\bf R^5}$  are independently selected at each occurrence from the group consisitng of:

- a) hydrogen,
- b)  $C_1-C_4$  alkyl,
- c)  $-(C_1-C_4 \text{ alkyl}) \text{aryl}$ , or
- d) C5-C7 cycloalkyl;

 $\rm R^6,\ R^7,\ R^8$  and  $\rm R^9$  are independently selected at each occurrence from the group consisting of:

- a) hydrogen,
- b)  $C_1-C_4$  alkyl,
- 10 c)  $C_1$ - $C_4$  alkoxy,
  - d) aryl,
  - e)  $-(C_1-C_4 \text{ alkyl}) \text{aryl}$ ,
  - f)  $-(C_1-C_4 \text{ alkyl})$  -heterocycle,
  - g) -0-aryl,
- 15 h)  $-(CH_2)_p-CO_2R^4$ ,
  - i)  $R^6$  and  $R^7$  can be taken together to form a ( $C_2$ - $C_7$ ) alkyl, or
  - j)  $R^8$  and  $R^9$  can be taken together to form a ( $C_2$ - $C_7$ ) alkyl;

20

25

R<sup>10</sup> is:

phenyl, wherein phenyl is optionally substituted with one to three substituents selected from the group consisting of halogen,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_7$ - $C_{15}$  alkylaryl,  $C_7$ - $C_{15}$  alkoxyaryl, methylenedioxy, -NO<sub>2</sub>, -CF<sub>3</sub>, -SH, -S(O)<sub>r</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), CN, -OH, -NH<sub>2</sub>, -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)<sub>2</sub>, -NHCOR<sup>4</sup>, -(CH<sub>2</sub>)<sub>p</sub>-CO<sub>2</sub>R<sup>4</sup>;

- $30 R^{11} is:$ 
  - a)  $C_1-C_4$  alkyl,
  - b) C<sub>3</sub>-C<sub>6</sub> cycloalkyl,
  - c)  $-OR^4$ ,
  - d)  $-NR^{15}R^{16}$ ,
- 35 e)  $-NC(=0)R^{15}R^{16}$ ,
  - $f) NR^{15}C (=0) OR^4$

g) aryl,

h)  $-(C_1-C_4 \text{ alkyl})-\text{aryl}$ ,

i) heteroaryl,

j)  $-(C_1-C_4 \text{ alkyl})-\text{heteroaryl},$ 

k)  $-(C_1-C_4 \text{ alkyl})-CO_2R^4$ 

heterocycle,

m)  $-(C_1-C_4 \text{ alkyl})$ heterocycle,

n)

10 0)

p)

q)

r)

15

 ${\rm R}^3$  and  ${\rm R}^{11},$  when taken together to form a ring bonded to the nitrogen:

### 5

## 10 c)

### d)

WO 95/09634 PCT/US94/11280

و المرابع الم

5

10

h)

 $\mathbb{R}^4$  and  $\mathbb{R}^{11}$  can also be taken together to form -(CH<sub>2</sub>)<sub>2</sub>-;

- 5 R<sup>11</sup>' is independently selected at each occurrence from the group consisting of:
  - a) hydrogen;
  - b) C<sub>1</sub>-C<sub>4</sub> alkyl
  - $c) OR^4$
- 10
- d)  $-NR^{15}R^{16}$
- e)  $-NC(=0)R^{15}R^{16}$
- $f) -NR^{15}C(=0)OR^4$
- g) aryl,
- h)  $-(C_1-C_4 \text{ alkyl})-\text{aryl}$ ,
- i) heteroaryl,
  - j) -(C<sub>1</sub>-C<sub>4</sub> alkyl)-heteroaryl,
  - k)  $-(C_1-C_4 \text{ alkyl})-CO_2R^4$
  - 1) heterocycle,
  - m) -(C<sub>1</sub>-C<sub>4</sub> alkyl)heterocycle,

20

 ${\bf R}^{13}$  is independently selected at each occurrence from the group consisting of:

- a) hydrogen
- 25
- b) halogen,
- c) C<sub>1</sub>-C<sub>4</sub> alkyl,
- d)  $C_1-C_4$  alkoxy,
- e) methylenedioxy,
- $f) -NO_2$ ,
- 30
- g) -CF3,

```
h) -SH,
           i) -S(0)_r - (C_1 - C_4 \text{ alkyl}),
           j) -CN,
           k) - OH,
           1) - NH_2,
 5
           m) -NH(C_1-C_4 \text{ alkyl}),
           n) -N(C_1-C_4 \text{ alkyl})_2,
           o) -NHC(=0)R^4, or
           p) -(CH_2)_p-CO_2R^4;
10
     R^{14} is:
            a) -CF3,
            b) -CHF2,
            c) -CH2F,
            d) -CH2Cl,
15
            e) -C(=0)OR^4,
            f) -C(=0)NR^{15}R^{16},
            g) -C(=0)R^4,
            h) -C(=0)COOR^4,
            i) -C(=0)C(=0)NR^{15}R^{16},
20
            j) -C(=0)C(=0)R^{4},
            k) -CY^3Y^4COOR^4,
            1) -CY^3Y^4C (=0)NR^{15}R^{16}, or
            m) -CY^3Y^4C (=0)R^4;
25
     \mathrm{R}^{15} and \mathrm{R}^{16} are independently selected at each
     occurrence from the group consisting of:
            a) hydrogen,
            b) C_1-C_4 alkyl,
            c) -(C_1-C_4 \text{ alkyl}) - \text{aryl},
30
            d) C5-C7 cycloalkyl, or
            e) phenyl, unsubstituted or substituted by R^{13};
      {\rm R}^{15} and {\rm R}^{16} taken together to form a ring can also
             include:
35
             a)
```

b)

- 5 R<sup>17</sup> is:
  - a) hydrogen,
  - b)  $C_1-C_4$  alkyl,
  - c) aryl,
  - d)  $-(C_1-C_4 \text{ alkyl}) \text{aryl}$ , or
- e) C5-C7 cycloalkyl;

R<sup>18</sup> is:

- a) hydrogen,
- b)  $-(C_1-C_5)$  alkyl, or
- 15 c)  $-(C_1-C_5)$  haloalkyl,
  - d)  $-(C_1-C_5)$  alkoxy;

R<sup>19</sup> is:

- a) hydrogen,
- 20 b)  $-(C_1-C_5)$  alkyl,
  - c) halo, or
  - d)  $-(C_1-C_5)$  haloalkyl,
  - e) -NO2,
  - f)  $-NR^4R^5$ ,
- 25 g) -CN,
  - h)  $-(C_1-C_5)$  alkoxy;

 $\mathbb{R}^{20}$  is

- a) hydrogen; or
- 30 b)  $-N^2$  with amine protecting;

```
A is:
```

- $\cdot a) -BY^1Y^2$ , or
- b)  $-C(=0)R^{14}$ ,
- c)  $C(OH) R^{14}R^{18}$ ;

#### W is

- a) -0-,
- b)  $-S(0)_{r}^{-}$ ,
- c)  $-NR^4-$ , or
- 10 d)  $-NC(=0)R^{4}-;$

 $y^1$  and  $y^2$  are

- a) -OH,
- b) -F,
- 15 c)  $-NR^4R^5$ ,
  - d) C<sub>1</sub>-C<sub>8</sub> alkoxý, or

when taken together  $Y^1$  and  $Y^2$  form:

- e) a cyclic boron ester where said chain or ring contains from 2 to 20 carbon atoms and, optionally, 1-3 heteroatoms which can be N,
- optionally, 1-3 heteroatoms which can be N, S, or O,
  - f) a cyclic boron amide where said chain or ring contains from 2 to 20 carbon atoms and, optionally, 1-3 heteroatoms which can be N,
- 25 S, or O,
  - g) a cyclic boron amide-ester where said chain or ring contains from 2 to 20 carbon atoms and, optionally, 1-3 heteroatoms which can be N, S, or O;

30

20

 $y^3$  and  $y^4$  are

- a) -OH or
- b) -F;
- 35 n is 0 or 1; p is 0 to 3;

q is 0 to 4; r is 0 to 2; t is 1 to 3; v is 1 to 17;

5

wherein aryl is defined as phenyl, fluorenyl, biphenyl and naphthyl, which may be unsubstituted or include optional substitution with one to three substituents;

- heteroaryl is 2-, or 3-, or 4-pyridyl; 2-or 3-furyl; 2or 3-benzofuranyl; 2-, or 3-thiophenyl; 2- or 3benzo[b]thiophenyl; 2-, or 3-, or 4-quinolinyl; 1-, or
  3-, or 4-isoquinolinyl; 2- or 3-pyrrolyl; 1- or 2- or
  3- indolyl; 2-, or 4-, or 5-oxazolyl; 2-benzoxazolyl;
  2- or 4- or 5-imidazolyl; 1- or 2- benzimidazolyl; 2or 4- or 5-thiazolyl; 2-benzothiazolyl; 3- or 4- or 5isoxazolyl; 3- or 4- or 5-pyrazolyl; 3- or 4- or 5isothiazolyl; 3- or 4-pyridazinyl; 2- or 4- or 5-
- pyrimidinyl; 2-pyrazinyl; 2-triazinyl; 3- or 420 cinnolinyl; 1-phthalazinyl; 2- or 4-quinazolinyl; or 2quinoxalinyl ring; said ring(s) may be unsubstituted or
  include optional substitution with one to three
  substituents.
- 25 heterocycle is tetrahydroisoquinoline,
   tetrahydroquinoline, tetrahydrofuran,
   tetrahydrothiophene, piperidine, piperazine,
   morpholino; said ring(s) may be unsubstituted or
   include optional substitution with one to three
  30 substituents.

the substituents that may be attached to the aryl, heteroaryl and heterocycle ring(s) may be independently selected at each occurrence from the group consisting

of:
halogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, methylenedioxy,

-NO2, -CF3, -SH, -S(O)<sub>r</sub>-(C<sub>1</sub>-C<sub>4</sub> alkyl), CN, -OH, -NH2, -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)<sub>2</sub>, -NHC(=O)R<sup>4</sup>, -(CH<sub>2</sub>)<sub>p</sub>-CO<sub>2</sub>R<sup>4</sup>, phenyl which may be unsubstituted or substituted with R<sup>13</sup>.

5

# 2. A compound of claim 1 wherein:

A is  $-BY^1Y^2$ ;

10

heteroaryl is 2-, 3-, or 4-pyridyl; 2-, or 3-furyl; 2-, or 3-thiophenyl; 2-, 3-, or 4-quinolinyl; or 1-, 3-, or 4-isoquinolinyl which may be unsubstitued or include optional substitution with one to three substituents;

15

heterocycle is 1-, 3-, or 4-tetrahydroisoquinolinyl, 2or 3-pyrrolidinyl, and 2-, 3- or 4-piperidinyl which may be unsubstituted or include optional substitution with one to three substituents.

20

3. A compound of claim 2 selected from the group consisting of:

Hydrocinnamoy1-[N-(Phenethyl)-Gly]-boroOrn-C10H16 HCl

25 Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn(CH=NH)-C10H16 HCl

Hydrocinnamoyl-[N-(Phenethyl)-Gly]-boroOrn(CH=NH)-OH HCl

Hydrocinnamoy1-[N-(Phenethyl)-Gly]-boroArg(CH3)-C10H16

30 HC

Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroLys-C10H16 HCl Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroLys-OH HCl

Hydrocinnamoyl-[N-(N(CH3)2)-Gly]-boroOrn-C10H16 HCl

Hydrocinnamoy1-[N-(N(CH3)2)-Gly]-boroOrn(CH=NH)-C10H16

35 HC

Methanesulfonyl-Sar-[N-(Phenethyl)-Gly]-boroLys-C10H16

```
HC1
    Methanesulfonyl-Sar-[N-(Phenethyl)-Gly]-boroLys-OH HCl
    Methanesulfonyl-Gly-[N-(Phenethyl)-Gly]-boroLys-C10H16
         HCl
   Hydrocinnamoyl-[N-(3-(Trifluoromethyl)-Phenethyl)-Gly]-
5
         boroLys-C10H16 HCl
    Hydrocinnamoy1-[N-(3-(Trifluoromethy1)-Phenethy1)-Gly]-
         boroLys-OH HCl
    Hydrocinnamoy1-[N-(3-(Methy1)-Phenethy1)-Gly]-boroLys-
10
         C10H16 HCl
    Hydrocinnamoy1-[N-(3-(Methy1)-Phenethy1)-Gly]-boroLys-
         OH HCl
    Succinyl-[N-(3-(Methyl)-Phenethyl)-Gly]-boroLys-C10H16
         HC1
    Hydrocinnamoy1-[N-(2,2-(Dimethy1)-Phenethy1)-Gly]-
15
         boroLys-OH HCl
    Hydrocinnamoyl-[N-(2-(Cyclopropyl)-Phenethyl)-Gly]-
         boroLys-C10H16 HCl
    Hydrocinnamoy1-[N-(2-(Cyclopropy1)-Phenethy1)-Gly]-
         boroLys-OH HCl
20
    Hydrocinnamoy1-[N-(2-(Cyclopropy1)-Phenethy1)-Gly]-
         boroOrn(CH=NH)-OH HCl
    Hydrocinnamoyl-[N-(2,2-(Diethyl)-Phenethyl)-Gly]-
         boroLys-C10H16 HCl
    Hydrocinnamoyl-Sar-Lys[C(=0)-C(=0)-OH]
25
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly[CH2)3-Br]-
         C10H16
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly(CH2)4)-
          Br]-C10H16
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroIrg-C10H16 HBr
30
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly[CH2)3-N3]-
         C10H16
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-borohomolrg-C10H16
     (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroGly[CH2)4)-
35
```

N3]-C10H16

```
(2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroOrn-C10H16 HCl
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-borohomoArg-C10H16
         HC1
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroArg-C10H16 HCl
   (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroOrn(CH=NH)-
5
         C10H16 HCl
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroLys(CH=NH)-
         C10H16 HCl
    2-Benzyl-(N-Benzyl)-Sar-boroLys-C10H16 HCl
    2-Thiophenyl-Benzoyl-Sar-boroLys(CH=NH)-C10H16
10
    2-(Thiophenyl)-Benzoyl-Sar-boroIrg-C10H16 HBr
    2-(Thiophenyl)-Benzoyl-Sar-boroOrn-C10H16 HCl
    2-(Thiophenyl)-Benzoyl-Sar-boroOrn(CH=NH)-C10H16 HCl
    Pinanediol N-{N-methyl-N-[2-(Thiophenyl)-Benzoyl]Sar}-
         1-amido-5-thiocyanatobutane boronate
15
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroLys-C10H16 HCl
    Acetyl-Gly[N-(2-(Benzyl))-Benzyl)]-boroLys-C10H16 HCl
    Pinanediol N-{N-methyl-N-[2-(pyrrol-1-ylmethyl)-
         Benzyl]glycyl}-1-amido-5-aminopentaneboronate,
         hydrochloride salt
20
    N-{N-methyl-N-[2-(pyrrol-1-ylmethyl)-Benzyl]glycyl}-1-
         amido-5-aminopentaneboronic acid, hydrochloride
         salt2-(2-(Trifluoromethyl)-Benzyl)-Benzoyl-Sar-
         Lys-C(=O)-NHNH2 2 HC1
    2-(Benzyl)-Benzoyl-Sar-Lys-C(=0)-NHNH2 2 HCl
    [3-(Trifluoromethyl)-Benzyl]-Benzoyl-Sar-boroLys-C10H16
         HC1
    3-(3-(Chloro)-Benzyl)-Benzoyl-Sar-boroLys-C10H16 HCl
    Hydrocinnamoyl-Sar-Lys(Z)-C(=0)-O-(CH2)2-NH(Z)
    Hydrocinnamoyl-Sar-Lys-C(=O)-O-(CH2)2-NH2 2 HCl
    Hydrocinnamoyl-Sar-Lys(Z)-C(=0)-OCH3
    Hydrocinnamoyl-Sar-Lys-C(=0)-OCH3 HCl
     Hydrocinnamoyl-Sar-Lys-C(=O)-CH3 HCl
     Hydrocinnamoyl-Sar-Lys(Z)-H
     Hydrocinnamoyl-Sar-NHCH(CH2OH)(CH2)4-NH(Z)
     Hydrocinnamoyl-Sar-NHCH(CH2OH)(CH2)4-NH2
```

```
Hydrocinnamoyl-Sar-Lys[CH(OH)(OCH3)-C(=0)-OCH3] HCl
   Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn-C10H16 HCl
   Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl
   Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn(CH=NH)-OH
         HCl
5
    Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroOrn(CH=NH)-
         C10H16 HCl
    Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys(CH=NH)-
         C10H16 HCl
    Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys(CH=NH)-OH
10
         HCl
    Phenoxyacetyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl
    Thiophenacetyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl
    Phenoxyacetyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl
    Thiophenacetyl-[N-(Cyclopropyl)-Gly]-boroLys-OH HCl
15
    Hydrocinnamoyl-[N-(N-(Methyl)-Phenyl)-Gly]-boroLys-
         C10H16 HCl
    Hydrocinnamoyl-[N-(N-(Methyl)-Phenyl)-Gly]-boroLys-OH
         HC1
    Hydrocinnamoyl-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-
20
         C10H16 HCl
    Glutaryl-[N-(Phenethyl)-Gly]-boroLys-OH HCl
     Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-
          C10H16
    Methyl Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-
25
         boroLys-C10H16 HCl
     Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-boroLys-OH
     Methyl Glutaryl(3,3-Dimethyl)-[N-(Phenethyl)-Gly]-
          boroLys-OH HCl
     Boc-Asp-[N-(Phenethyl)-Gly]-boroLys-C10H16
30
     Boc-Glu-[N-(Phenethyl)-Gly]-boroLys-C10H16
     Boc-Glu(OCH3)-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl
     Boc-Glu-[N-(Phenethyl)-Gly]-boroLys-OH
     Hydrocinnamoyl-[N-(N-(Methyl)-Benzyl)-Gly]-boroLys-OH
          HC1
35
     Methanesylfonyl-Gly-[N-(N-(Methyl)-Benzyl)-Gly]-
```

```
boroLys-C10H16 HCl
   Methanesulfonyl-Gly-[N-(N-(Methyl)-Benzyl)-Gly]-
         boroLys-OH HCl
   Hydrocinnamoyl-[N-(Succinyl)-Gly]-boroLys-C10H16
5 Hydrocinnamoyl-[N-(Methyl Succinyl)-Gly]-boroLys-C10H16
         HCl
    Succinyl-[N-(Phenethyl)-Gly]-boroLys-OH
    Methyl Succinyl-[N-(Phenethyl)-Gly]-boroLys-OH HCl
    Glutaryl-[N-(Phenethyl)-Gly]-boroLys-C10H16
10 Methyl Glutaryl-[N-(Phenethyl)-Gly]-boroLys-C10H16 HCl
    Methyl Glutaryl-[N-(Phenethyl)-Gly]-boroLys-OH HCl
    [N-(-C(O)(CH_2)_2Ph)-N-(CH_2CH_2Ph)]-Gly-boroLys-OH
    [N-(-C(O)(CH_2)_2Ph)-N-(CH_2C(CH_3)_2Ph)]-Gly-boroLys-OH
    [N-(-C(O)(CH_2)_2Ph)-N-(CH_2C(ethanediyl)Ph)]-Gly-boroLys-
         OH
15
    Hydrocinnamoy1-[N-(2-(Cyclopropyl)-Phenethyl)-Gly]-
         boroArg-C10H16 HCl
    Hydrocinnamoy1-[N-(2-(Cyclopentyl)-Phenethyl)-Gly]-
         boroLys-C10H16 HCl
    (2-(2-Cyano)-Thiophenyl)-Benzoyl-Sar-boroArg-OH HCl
20
    Hydrocinnamoy1-[N-(2,2-(Dimethy1)-Phenethy1)-Gly]-
         boroLys-C10H16 HCl
    Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
         boroOrn-C10H16 HCl
    Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
25
         boroLys-OH HCl
    Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
         boroArg-C10H16 HCl
    Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
         boroOrn (CH=NH) -C10H16 HCl
30
    Hydrocinnamoy1-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
         .boroArg-OH HCl
    Hydrocinnamoy1-[N-(2,2-(Dimethy1)-Phenethy1)-Gly]-
          boroOrn(CH=NH)-OH HCl
    Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]-
35
          boroOrn-C10H16 HCl
```

Hydrocinnamoyl-[N-(2,2-(Dimethyl)-Phenethyl)-Gly]boroLys-C10H16 HCl

- Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-boroOrn-C10H16 HCl
- 5 Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-boroLys-C10H16 HCl
  - Hydrocinnamoy1-{N-[2,2-(Dimethy1)-2-(3,5-dimethylpheny1)-ethyl]-Gly}-boroArg-C10H16 HC1

Hydrocinnamoyl-{N-[2,2-(Dimethy1)-2-(3,5-dimethylpheny1)-ethyl]-Gly}-boroOrn(CH=NH)-C10H16

- dimethylphenyl)-ethyl]-Gly}-boroOrn(CH=NH)-C10H16
  HCl
  - Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-boroArg-C10H16 HCl
  - Hydrocinnamoyl-{N-[2-(3,5-dimethylphenyl)-ethyl]-Gly}-boroOrn(CH=NH)-C10H16 HCl

Hydrocinnamoyl-[N-(Cyclohexyl)-Gly]-boroLys-C10H16 HCl Hydrocinnamoyl-[N-(Cyclopropyl)-Gly]-boroLys-C10H16 HCl Hydrocinnamoyl-[N-(Cyclohexyl)-Gly]-boroLys-OH HCl

- 20 4. A pharmaceutical composition comprising a pharmaceutically suitable carrier and a therapeutically effective amount of a compound of claim 1.
- 5. A pharmaceutical composition comprising a pharmaceutically suitable carrier and a therapeutically effective amount of a compound of claim 2.
  - A pharmaceutical composition comprising a
    pharmaceutically suitable carrier and a therapeutically
    of effective amount of a compound of claim 3.
    - 7. A method of treating a physiological disorder in a warm blooded animal catalyzed by trypsin-like enzymes comprising administering to an animal in need such treatment an effective amount of a compound of claim 1.